


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Details of the Classical and Long-term Experiments Up to 1962

[Full Table of Content](#)



Rothamsted Garden- Clover

Rothamsted Research

Rothamsted Research (1966) *Rothamsted Garden- Clover* ; Details Of The Classical And Long-Term Experiments Up To 1962, pp 31 - 31 - DOI: <https://doi.org/10.23637/ERADOC-1-191>

ROTHAMSTED GARDEN CLOVER, 1854 ONWARDS

First crop sown in spring 1854 on a plot in the kitchen garden of the Manor House, and resown when necessary ever since. Complete resowing or patching is now almost a yearly operation.

Variety: Red Clover.

In 1956 the plot was divided into two to test an annual dressing of muriate of potash at 2 cwt. per acre (4 cwt. per acre in 1961).

In 1960 the sub-plots were halved to test the effect of a foliage spray of molybdenum.

2 or 3 cuts of greenstuff are taken each season.

Whole plot area: 0.0022 acres.

For an account of the history and yields of this plot see Rep. Rothamst. exp. Sta. for 1956, pp. 187-189.

Table 17

CLOVER - ROTHAMSTED GARDEN

Dry matter: cwt per acre

Means over 6 years
1957-62

Muriate of Potash: cwt per acre

None	2*	Mean
18.1	34.6	26.4

NB For 1960-62 the yields from the plots sprayed with sodium molybdate have been excluded.

Means over 3 years
1960-62

Muriate of Potash: cwt per acre	Spray		Mean
	None	Sodium Molybdate	
None	9.3	3.7	6.5
2*	22.7	15.7	19.2
Mean	16.0	9.7	12.8

* 4 cwt in 1961