


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Details of the Classical and Long-term Experiments 1968-73

[Full Table of Content](#)



### R/AG/6 Agdell - Barley, Potatoes, Sugar Beet

#### Rothamsted Research

Rothamsted Research (1977) *R/AG/6 Agdell - Barley, Potatoes, Sugar Beet* ; Details Of The Classical And Long-Term Experiments 1968-73, pp 19 - 23 - DOI: <https://doi.org/10.23637/ERADOC-1-193>

## AGDELL

### RESIDUAL EFFECTS OF P AND K

(R/AG/6)

The cropping and management in 1968 and 1969 continued the pattern set in 1964 when large fresh dressings of P and K were applied to the sub-plots of both grass and fallow areas (*Details 1967*, pp 26-27. Note that the heading in Table 7 p. 27 should read 1920-51 and not 1920-53).

In 1970 a further scheme was introduced with annual applications of P (1970-72) and K (1973-75) to a three-course rotation of sugar beet, barley, potatoes (two crops present each year).

An outline of the layout of the plots from 1958 onwards is set out in the diagram on pages 22 and 23 to show the changes in plot boundaries and the relationship of the successive treatments. Details of treatments between 1958 and 1967 are given on pages 25-27 of *Details 1967*.

#### Detailed treatments 1968-73

##### (1) *P test half plots*

- (a) 1968-69 – Half fallow, half Timothy (S 51 sown 1964) 3 dressings of 100 kg N applied each year to the grass. P and K applied each autumn or early winter to replace removals in grass the previous year, except that plots without fresh P in 1964 continued to receive none to measure the release of P residues accumulated during 1848-1951.

- (b) 1970-72 – Effects of different amounts of soil P were tested in a three-course rotation – sugar beet, barley, potatoes, starting with sugar beet and barley. Fresh P was tested on the sub-plots cumulatively.

|            |                 |
|------------|-----------------|
| Barley     | None v. 27 kg P |
| Sugar beet | None v. 55 kg P |
| Potatoes   | None v. 82 kg P |

Basal manuring. Sugar beet: 190 kg N; 260 kg K as muriate of potash; 60 kg Mg as kieserite.

Barley: 95 kg N, 50 kg K as (25-0-16).

Potatoes: 250 kg N, 210 kg K, 60 kg Mg – materials as for sugar beet.

- (c) 1973 – Rotation ended, barley grown testing residues of P applied in 1964 and 1970-72 with a fresh N test applied in strips of sixty-fourth plots.

|    |         |
|----|---------|
| N1 | 63 kg N |
| N2 | 95 kg N |

##### (2) *K test half plots*

- (a) 1968-70 – Half-fallow, half Timothy (S 51 sown 1964). 100 kg N applied three times in 1968 and 1969, twice in 1970 to the grass.

Balancing P and K applied each autumn or winter to replace removals in grass, except that plots without fresh K in 1964 continued to receive none to measure the recovery of K residues accumulated during 1848-1951 and the release of soil K.

- (b) 1971-72 – The whole area was fallowed in 1971 and cropped with oats in 1972.

Standard applications: Oats – 75 kg N, 14 kg P as (30-13-0)

- (c) 1973 – The three-course rotation commenced with sugar beet and barley.

K treatments applied in first year of rotation.

|            |                  |
|------------|------------------|
| Barley     | None v. 50 kg K  |
| Sugar beet | None v. 257 kg K |

Standard applications: Sugar beet: 190 kg N, 55 kg P as granular superphosphate, 60 kg Mg as kieserite.

Barley: 95 kg N, 18 kg P as (30-13-0).

### (3) *Compensatory dressings of P and K*

Dressings of triple superphosphate and muriate of potash were applied to compensate for the removals in the grass during the years 1964-69 from the P plots and 1964-70 from the K plots. Dressings were normally applied annually to compensate for the removals during the previous season but in the period 1965-67 certain adjustments between years were made. (For details see *Results 1965 to 1970*).

No phosphate was applied to the PO plots and No K to the KO plots throughout the period.

The total removed and replaced (except on the PO and KO plots) were:

| P (kg/ha) |                     |     |     |     |                     |     |     |     |
|-----------|---------------------|-----|-----|-----|---------------------|-----|-----|-----|
|           | Sub-plots testing P |     |     |     | Sub-plots testing K |     |     |     |
|           | P0                  | P1  | P2  | P4  | K0                  | K1  | K2  | K4  |
| Plot 1    | (102)               | 165 | 175 | 186 | 157                 | 194 | 200 | 200 |
| 2         | (68)                | 139 | 158 | 162 | 134                 | 188 | 198 | 198 |
| 3         | (67)                | 147 | 148 | 175 | 148                 | 171 | 176 | 175 |
| 4         | (48)                | 122 | 127 | 151 | 160                 | 189 | 177 | 190 |
| 5         | (38)                | 111 | 128 | 144 | 126                 | 163 | 162 | 167 |
| 6         | (22)                | 117 | 133 | 149 | 125                 | 171 | 169 | 169 |

| K (kg/ha) |                     |      |      |      |                     |      |      |      |
|-----------|---------------------|------|------|------|---------------------|------|------|------|
|           | Sub-plots testing P |      |      |      | Sub-plots testing K |      |      |      |
|           | P0                  | P1   | P2   | P4   | K0                  | K1   | K2   | K4   |
| Plot 1    | 1398                | 1694 | 1664 | 1678 | (689)               | 1453 | 1601 | 1787 |
| 2         | 1020                | 1481 | 1611 | 1532 | (493)               | 1378 | 1631 | 1798 |
| 3         | 974                 | 1525 | 1471 | 1576 | (612)               | 1429 | 1441 | 1619 |
| 4         | 736                 | 1387 | 1313 | 1451 | (707)               | 1461 | 1509 | 1732 |
| 5         | 543                 | 1303 | 1381 | 1449 | (485)               | 1291 | 1423 | 1647 |
| 6         | 371                 | 1311 | 1371 | 1379 | (448)               | 1378 | 1363 | 1549 |

### Liming

In the autumn of 1969 plots 1 and 2 and the south halves of plots 3 (P test) and 4 (K test) received ground chalk at 3 t

### Cropping and Weed Control

#### (1) Varieties

|             |                                                                    |
|-------------|--------------------------------------------------------------------|
| Sugar beet: | Klein E                                                            |
| Barley:     | Julia (dressed with ethirimol 1973)                                |
| Potatoes:   | King Edward, once grown Rothamsted paracrinkle-free seed, chitted. |
| Oats:       | Manod                                                              |

#### (2) Weedkillers

|           |                                                  |
|-----------|--------------------------------------------------|
| Potatoes: | 1971 and 1972: Linuron with paraquat             |
| Oats:     | 1972: Bromoxynil, ioxynil, dichloroprop and MCPA |
| Grass:    | 1968: Ioxynil with mecoprop.                     |

### Other Chemicals applied

|             |                                                     |
|-------------|-----------------------------------------------------|
| Sugar Beet: | 1970 and 1972: Menazon, 1972: pyrethrum             |
| Potatoes:   | 1971 and 1972: Menazon and mancozeb, 1972: captafol |

### Areas harvested

|             |                   |
|-------------|-------------------|
| Grass:      | 0.00081 – 0.00186 |
| Barley:     | 0.00061 – 0.00087 |
| Sugar beet: | 0.00077           |
| Potatoes:   | 0.00069           |


**Soil series** Winchester and shallow Batcombe series

### References

1. Johnston, A.E., Warren, R.G., and Penny, A. (1970)  
The value to arable crops of residues accumulated from superphosphate and from potassium fertiliser.  
*Rothamsted Experimental Station. Report for 1969, Part 2, 39-90.*
2. Johnston, A.E., & Penny, A. (1972)  
The Agdell Experiment 1848-1970.  
*Rothamsted Experimental Station. Report for 1971, Part 2, 38-68.*
3. Johnston, A.E., & Mitchell, J.D.D. (1974)  
Potassium in soils from the Agdell experiment.  
*Rothamsted Experimental Station. Report for 1973, Part 2, 74-97.*

# AGDELL

## Arrangement of sub-plot tests 1958-1973



1959-63 Details of sub-plots on plot 1b (one replicate only shown)

## 1959

| Crop | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | Plot no. |
|------|---|---|---|---|---|---|---|---|---|----|----|----|----------|
| P    | - | 4 | 1 |   |   |   |   |   |   |    |    |    |          |
| S    | - | 1 | 4 |   |   |   |   |   |   |    |    |    |          |
| B    | 4 | 1 | - |   |   |   |   |   |   |    |    |    |          |

288 links

61

## 1960

|   | 1  | 4  | 7 | 10 |  |
|---|----|----|---|----|--|
| S | -1 | -6 | 6 | 4  |  |
| B | -1 | 6  | 4 | -6 |  |
| P | 6  | -4 | 6 | -1 |  |

61

1961 Sub divisions of plots 1b, 3a, 5b only (2a, 4b, 6b fallowed)

|   | 1  | 4  | 7  | 10    |  |
|---|----|----|----|-------|--|
| B | A3 | B6 | B3 | A6 B6 |  |
| P | A6 | C6 | C3 | A3 C3 |  |
| S | B3 | A3 | A6 | B6    |  |


61

1962 Plots 2a, 4b, 6b sub divided as above (1b, 3a, 5b fallowed)

1963 All plots fallowed

(1958-63 G plots remained in grass)

## 1964 The whole experiment



# 1970~73 Details of sub-plots on plot 1

