


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Report 1918-20 With the Supplement to the Guide to the Experimental Plots Containing the Yields per Acre Etc.



[Full Table of Content](#)

Table of Results - the Classical Experiments

Rothamsted Research

Rothamsted Research (1921) *Table of Results - the Classical Experiments* ; Report 1918-20 With The Supplement To The Guide To The Experimental Plots Containing The Yields Per Acre Etc., pp 64 - 85
- DOI: <https://doi.org/10.23637/ERADOC-1-109>

DATES OF SOWING AND HARVESTING (Harvest of 1918).

Field.	Crop.	Variety.	Sowing began.	Sowing finished.	Cutting began.	Carting began.	Carting finished.	Yield per Acre.
Great Knott Wood, east	Oats	Grey Winter	Oct. 24, '17	Oct. 27, '17	Aug. 1	Aug. 15	Aug. 17	37 bush.
" " west	"	"	"	"	"	"	"	"
Little Knott Wood	Wheat	Red Standard (6 acres)	Nov. 8, '17	Nov. 8, '17	Aug. 10	Aug. 18	Aug. 18	40.5 "
Fosters, east	Barley	Red Marvel (7 acres)	Nov. 28, '17	Dec. 1, '17	"	Aug. 27	Aug. 27	33 "
" " west	"	Plumage Archer	Apr. 4, '18	Apr. 10, '18	Sept. 7	Sept. 11	Sept. 12	28 "
West Barnfield	Potatoes	Arran Chief and King Edward	May 3, '18	May 11, '18	Nov. 7	"	"	15 tons* ware & 5 cwts. chats
Long Hoos, east	Wheat	Red Standard	Nov. 6, '17	Nov. 23, '17	Aug. 17	Aug. 24	Aug. 30	40.5 bush.
" " west	"	"	"	"	"	"	"	"
Great Harpenden	Clover	Red	May 8, '17	May 8, '17	June 20	June 27	June 27	1.5 tons†
New Zealand	Swedes	Magnum Bonum and Early White	June 11, '18	June 11, '18	Failed	"	"	"
Stackyard	Mangolds	Yellow Globe	May 28, '18	May 28, '18	Nov. 18	"	"	12.3 tons
Sawpit	Wheat	Red Marvel (6 acres)	Nov. 30, '17	Nov. 30, '17	Aug. 16	Aug. 24	Aug. 24	33 bush.
Broadbalk	"	Red Standard	Oct. 31, '17	Nov. 1, '17	Aug. 12	Aug. 20	Aug. 21	see p. 74
Little Hoos	"	"	Nov. 3, '17	Nov. 5, '17	Aug. 12	Aug. 20	Aug. 21	" 77
Hoos	Barley	Plumage Archer	Mar. 18, '18	Mar. 18, '18	Sept. 2	Sept. 7	Sept. 9	" 76
Barnfield	Mangolds	Sutton's Yellow Globe	Apr. 27, '18	Apr. 27, '18	Nov. 7	"	"	" 69
Agdell	Clover	Red 1st Crop	May 14, '17	May 14, '17	June 22	July 2	July 2	" 67
"	"	" 2nd Crop	"	"	Aug. 31	Sept. 3	Sept. 3	" 67
Greatfield	Grass	Rented out for Grazing	"	"	"	"	"	"
Park	"	1st Crop	"	"	June 24	July 3	July 4	see p. 70
"	"	2nd Crop	"	"	Sept. 16	Sept. 24	Sept. 25	" 70

NOTE.—7 acres of Sawpit were fallow.

* Quantity sold after clamping, see pp. 79 and 83.

† Estimated always on measurements, not weighed.

DATES OF SOWING AND HARVESTING (Harvest of 1919).

Field.	Crop.	Variety.	Sowing began.	Sowing finished.	Cutting began.	Carting began.	Carting finished.	Yield per Acre.
Great Knott Wood, east	Wheat ...	{ Red Standard ...	Nov. 26, '18	Nov. 26, '18	Aug. 23	Sept. 8	Sept. 8	20 bush.
" " west	" ...	{ Red Marvel ...	Mar. 25, '19	Mar. 25, '19	Sept. 18	Oct. 1	Oct. 1	(see p. 83)
Little Knott Wood	" ...	Red Standard ...	Nov. 9, '18	Nov. 9, '18	Aug. 22	Sept. 5	Sept. 5	1 ton
	{ Grass and Clover Ley	{ Red Clover, Alsike } 1st Crop	Apr. 8, '18	Apr. 8, '18	June 12	June 17	June 17	12 cwt.
		{ Timothy, Cocksfoot and Bent ... } 2nd Crop	" "	" "	Aug. 15	Aug. 18	Aug. 18	1 ton
Foster's, east	" ...	" "	May 10, '18	May 10, '18	June 21	June 26	June 30	16 cwt.
" " west	" ...	" "	May 10, '18	" "	June 21	June 26	June 30	35 bush.
West Barnfield	Wheat ...	Red Standard ...	Nov. 19, '18	Nov. 25, '18	Aug. 21	Sept. 1	Sept. 1	28 bush.
Long Hoos, east	Barley ...	Plumage Archer	Apr. 18, '19	Apr. 18, '19	Sept. 8	Sept. 17	Sept. 17	...
	Mangolds	Sutton's Prize Winner, Yellow Globe	May 27, '19	May 27, '19	Oct. 19	...	...	94 tons
" " west	Turnips ...	Sutton's Aberdeen Green Top Turnip	June 21, '19	June 21, '19	Oct. 28	...	...	5½ tons
Great Harpenden	Potatoes	King Edward and Arran Chief	May 20, '19	May 20, '19	Oct. 16	...	...	62 bush
New Zealand	Oats ...	Grey Winter ...	Oct. 3, '18	Oct. 3, '18	July 31	Aug. 12	Aug. 14	36 bush.
Stackyard	Barley ...	Plumage Archer	Apr. 11, '19	Apr. 17, '19	Sept. 2	Sept. 11	Sept. 16	32 bush.
Sawpit	" ...	" "	Apr. 9, '19	Apr. 9, '19	Sept. 1	Sept. 8	Sept. 8	34 bush.
Broadbalk	Wheat ...	Red Standard ...	Oct. 16, '18	Oct. 23, '18	Aug. 15	Aug. 23	Aug. 27	see p. 74
Little Hoos	" ...	" "	Oct. 25, '18	Oct. 26, '18	Aug. 14	Sept. 2	Sept. 5	" 77
Hoos	Barley ...	Plumage Archer	May 6, '19	May 6, '19	Sept. 11	Sept. 22	Sept. 22	" 76
Barnfield	" "	" "	Apr. 8, '19	Apr. 8, '19	Sept. 9	Sept. 16	Sept. 16	" 69
Agdell	Mangolds	Sutton's Yellow Globe	May 14, '19	May 14, '19	Oct. 29	...	...	" 67
Greatfield	Wheat ...	Red Standard ...	Oct. 18, '18	Oct. 18, '18	Aug. 11	Sept. 5	Sept. 5	70
	Grass	Rented out for Grazing	" "	" "	June 16	June 18	June 18	" 70
Park	" "	{ 1st Crop ... } { 2nd Crop ... }	" "	" "	Sept. 23	Sept. 29	Sept. 30	" 70

DATES OF SOWING AND HARVESTING (Harvest of 1920).

Field.	Crop.	Variety.	Sowing began.	Sowing finished.	Cutting began.	Carting began.	Carting finished.	Yield per Acre.
Great Knott Wood, east	Clover ...	Red	May 7, '19	May 7, '19	June 25	July 16	July 16	$\frac{3}{4}$ tons
" " west	Barley ...	Plumage Archer	Mar. 31, '20	Mar. 31, '20	Aug. 20	Aug. 30	Aug. 30	37 bush.
Little Knott Wood	{ Grass and Clover Ley	{ Red Clover, Alsike } 1st Crop Timothy, Cocksfoot } { and Bent } 2nd Crop	Apr. 8, '18	Apr. 8, '18	June 11	June 25	June 25	$1\frac{1}{2}$ tons
Foster's, east	Wheat ...	Yeoman ...	" "	" "	Sept. 2	Sept. 10	Sept. 10	"
" west	{ Grass and Clover Ley	{ Red Clover, Alsike, Timothy, Cocksfoot and Bent }	Sept. 25, '19	Sept. 25, '19	Aug. 9	Aug. 21	Aug. 21	22 bush.
West Barnfield	Oats ...	Grey Winters ...	May 10, '18	May 10, '18	June 28	July 14	July 14	$1\frac{3}{4}$ tons
Long Hoos, east	Potatoes ...	Arran Chief ...	Oct. 3, '19	Oct. 3, '19	Aug. 9	Aug. 17	Aug. 23	*40 bush.
" west	{ Barley Wheat	{ Plumage Archer Yeoman ... }	May 13, '20	May 22, '20	Oct. 25	" "	" "	{ 4 tons 1 cwt.
New Zealand	" "	Red Standard ...	Feb. 23, '20	Feb. 27, '20	Aug. 13	Aug. 26	Aug. 28	36 bush.
Great Harpenden	" "	Red Standard ...	Oct. 17, '19	Oct. 17, '19	Aug. 10	Aug. 21	Aug. 26	32 bush.
Stackyard	" "	Red Standard ...	Oct. 22, '19	Oct. 22, '19	Aug. 11	Aug. 25	Aug. 25	40 bush.
Sawpit	Oats	Red Standard ...	Oct. 18, '19	Oct. 18, '19	Aug. 10	Aug. 23	Aug. 23	32 bush.
Broadbalk	Wheat	Grey Winters ...	Oct. 27, '19	Oct. 27, '19	Aug. 12	Aug. 26	Aug. 26	39 bush.
Little Hoos	Swedes	Red Standard ...	Sept. 29, '19	Oct. 2, '19	Aug. 2	Aug. 16	Aug. 17	*40 bush.
Hoos	Barley	Sutton's Magnum Bonum	Oct. 24, '19	Oct. 25, '19	Aug. 16	Aug. 28	Aug. 31	see p. 74
Barnfield	Mangolds	Plumage Archer	May 18, '20	May 19, '20	Oct. 19	" "	" "	77
Agdell	Swedes	Sutton's Yellow Globe	Mar. 1, '20	Mar. 1, '20	Aug. 14	Aug. 27	Aug. 27	76
Greatfield, north	Grass	Sutton's Magnum Bonum	Apr. 29, '20	Apr. 29, '20	Dec. 5	" "	" "	69
" south	"	"	June 14, '20	June 14, '20	Oct. 19	" "	" "	67
Park	"	{ 1st Crop 2nd Crop }	" "	" "	July 19	Aug. 4	Aug. 18	81
			" "	" "	Aug. 17	Aug. 31	Sept. 9	81
			" "	" "	June 22	June 25	June 26	70
			" "	" "	Sept. 10	Sept. 27	Sept. 28	70

* Sawpit and West Barnfield produce not kept separate this year.

CROP YIELDS ON THE EXPERIMENTAL PLOTS.

NOTE.—In each case the year refers to the harvest, *e.g.*, Wheat harvested in 1920.

1 acre	=	0.404 Hectare	0.963 Feddan
1 bushel (Imperial)	=	0.346 Hectolitre (36.346 litres) ...	0.184 Ardeb.
1 lb. (pound avoirdupois)	=	0.453 Kilogramme	1.009 Rotls.
1 cwt. (hundredweight)	=	50.8 Kilogrammes	113.0 Rotls.
			1.366 Maunds
1 metric quintal ...	=	100.0 Kilogrammes	
		220.46 lb.	
1 bushel per acre ...	=	0.9 Hectolitre per Hectare ...	0.191 Ardeb per Feddan.
1 lb. per acre ...	=	1.12 Kilogramme per Hectare ...	1.049 Rotls per Feddan.
1 cwt. per acre ...	=	125.6 Kilogrammes per Hectare or	117.4 Rotls per Feddan.
		1.256 metric Quintals per Hectare	

In America the Winchester bushel is used = 35.236 litres. 1 English bushel = 1.032 American bushels.

Crops Grown in Rotation. Agdell Field.

PRODUCE PER ACRE.

Year.	CROP.	O. Unmanured		M Mineral Manure.		C. Complete Mineral and Nitrogenous Manure.	
		5.	6.	3.	4.	1.	2.
		Fallow.	Beans or Clover.	Fallow.	Beans or Clover.	Fallow.	Beans or Clover.

EIGHTEENTH COURSE, 1916-19.

1916	Roots (Swedes) cwt.	12.4	1.4	125.2	145.2	285.2	37.8*
	(Barley Grain bush.	9.4	2.5	14.2	15.2	13.1	15.0
1917	Barley Straw ... cwt.	11.6	5.1	16.8	15.6	13.1	19.8
1918	Clover Hay ... cwt.	—	19.5	—	59.5	—	17.0
	(1st and 2nd crops)						
1919	Wheat Grain bush.	8.0	3.4	13.5	7.9	17.5	2.2
	Wheat Straw ... cwt.	14.2	7.4	19.0	17.5	17.2	2.3

PRESENT COURSE (19th), 1920.

1920	Roots (Swedes) cwt.	20.4	2.2	163.8	270.0	262.2	56.4*
------	---------------------	------	-----	-------	-------	-------	-------

In 1920 Rape Cake was omitted from Plots 1 and 2.

In 1916 and 1920, the roots on Plot 2 were badly attacked by finger and toe disease.

RAIN AND DRAINAGE.
MONTHLY MEAN FOR 50 YEARS, 1870—1920.

	Rainfall.	Drainage.			Drainage % of Rainfall.			Evaporation.		
		20-in. Gauge	40-in. Gauge	60-in. Gauge	20-in. Gauge	40-in. Gauge	60-in. Gauge	20-in. Gauge	40-in. Gauge	60-in. Gauge
		Ins.	Ins.	Ins.	Ins.	Ins.	Ins.	Ins.	Ins.	Ins.
September	2.330	0.754	0.717	0.657	32.4	30.8	28.2	1.576	1.613	1.673
October ...	3.233	1.848	1.798	1.669	57.2	55.6	51.6	1.385	1.435	1.564
November	2.795	2.132	2.169	2.047	76.3	77.6	73.2	0.663	0.626	0.748
December	2.869	2.437	2.527	2.414	84.9	88.1	84.1	0.432	0.342	0.455
January...	2.364	1.892	2.078	2.001	80.0	87.9	84.6	0.472	0.286	0.363
February	2.008	1.480	1.584	1.513	73.7	78.9	75.4	0.528	0.424	0.495
March ...	2.103	1.148	1.285	1.215	54.6	61.1	57.8	0.955	0.818	0.888
April ...	2.012	0.653	0.727	0.695	32.5	36.1	34.5	1.359	1.285	1.317
May ...	2.025	0.475	0.537	0.502	23.5	26.5	24.8	1.550	1.488	1.523
June ...	2.375	0.595	0.616	0.595	25.1	25.9	25.0	1.780	1.759	1.780
July ...	2.667	0.680	0.703	0.654	25.5	26.4	24.5	1.987	1.964	2.013
August ...	2.719	0.740	0.741	0.697	27.2	27.3	25.6	1.979	1.978	2.022
Year ...	29.500	14.834	15.482	14.659	50.3	52.5	49.7	14.666	14.018	14.841

All four gauges measure $\frac{1}{1000}$ acre. Drain gauge records start Sept. 1st, 1870. Rain gauge records start Feb., 1853. For purpose of comparison the above figures deal with the same period as the drain gauge records, viz., Sept. 1st, 1870, to Aug. 31st, 1920.

METEOROLOGICAL RECORDS, 1918-20

	Rain.			Drainage through soil.			Bright Sun- shine.	Temperature. (Mean)			
	Total Fall.		No. of Rainy Days. (0.01 inch or more)					Max.	Min.	1 ft. in ground.	Solar Max.
	5-inch Funnel Gauge.	$\frac{1}{1000}$ Acre Gauge.	$\frac{1}{1000}$ Acre Gauge.	20 ins. deep.	40 ins. deep.	60 ins. deep.					
1918	Inches.	Inches.	No.	Inches.	Inches.	Inches.	Hours.	°F.	°F.	°F.	F.
Jan. ...	2.314	2.990	15	2.951	3.059	3.045	57.2	42.6	31.4	37.5	71.7
Feb. ...	1.027	1.232	15	0.537	0.553	0.526	66.3	46.9	36.6	40.8	78.2
Mar. ...	0.861	0.985	8	0.024	0.078	0.073	141.4	49.9	33.2	40.5	94.1
April ...	3.946	4.548	17	3.481	3.537	3.294	97.2	48.8	36.5	43.1	91.4
May ...	2.258	2.471	10	0.487	0.633	0.640	207.5	63.5	45.1	52.3	116.9
June ...	0.862	0.998	12	0.003	0.024	0.027	226.5	64.3	45.5	57.1	123.4
July ...	3.215	3.447	18	0.654	0.698	0.620	200.4	68.0	51.6	60.0	124.5
Aug. ...	1.163	1.331	11	0.004	0.032	0.040	178.9	68.9	52.9	61.4	122.9
Sept. ...	4.974	5.421	24	2.293	2.181	2.044	155.3	60.0	47.6	55.8	111.8
Oct. ...	1.703	1.964	15	1.094	1.140	1.065	78.8	53.5	41.9	49.6	88.1
Nov. ...	2.518	2.674	17	2.165	2.064	1.947	70.8	47.4	35.3	43.9	75.1
Dec. ...	2.839	3.175	26	2.814	2.897	2.754	36.5	48.7	39.8	44.3	65.1
Total or Mean	27.680	31.236	188	16.507	16.896	16.075	1516.8	55.2	41.5	48.9	96.9
1919	Inches.	Inches.	No.	Inches.	Inches.	Inches.	Hours.	°F.	°F.	°F.	F.
Jan. ...	3.840	4.281	25	2.964	3.079	2.980	32.7	40.3	31.5	38.3	60.9
Feb. ...	2.901	3.290	14	3.975	3.961	3.925	48.1	38.7	27.9	35.3	73.1
Mar. ...	3.432	3.747	19	2.796	2.871	2.801	107.3	43.9	33.2	38.7	88.0
April ...	3.311	3.693	16	1.970	2.034	2.020	120.4	51.3	36.2	43.1	106.0
May ...	0.460	0.535	5	0.208	0.359	0.370	257.7	65.1	45.1	52.7	119.0
June ...	1.045	1.159	7		0.009	0.018	230.7	66.6	47.6	59.6	124.3
July ...	2.625	2.767	15	0.330	0.394	0.379	120.1	63.0	49.0	57.7	114.2
Aug. ...	3.239	3.404	12	1.337	1.389	1.346	228.9	70.3	52.2	61.4	124.1
Sept. ...	1.191	1.293	10	0.076	0.118	0.093	158.3	63.4	47.1	57.2	112.2
Oct. ...	0.977	1.073	14	No drainage this month			136.2	51.4	36.5	46.7	93.5
Nov. ...	2.049	2.239	20	1.569	1.331	1.238	48.6	41.4	32.4	40.5	67.4
Dec. ...	5.048	5.573	24	5.717	5.836	5.801	33.4	46.1	35.3	40.0	63.2
Total or Mean	30.118	33.054	181	20.942	21.381	20.971	1522.4	53.5	39.5	47.6	95.5
1920	Inches.	Inches.	No.	Inches.	Inches.	Inches.	Hours.	°F.	°F.	°F.	F.
Jan. ...	2.730	3.015	21	2.548	2.620	2.590	51.0	45.7	34.3	39.6	66.5
Feb. ...	0.432	0.511	10	0.044	0.136	0.108	84.2	48.1	34.8	40.2	79.9
Mar. ...	1.403	1.629	17	0.399	0.405	0.407	141.4	52.3	36.4	42.5	98.6
April ...	4.246	4.585	20	3.167	3.183	3.163	90.3	52.3	40.8	47.1	100.8
May ...	1.208	1.336	13	0.009	0.064	0.061	241.6	61.1	45.1	52.7	118.9
June ...	1.832	1.927	12	0.045	0.079	0.098	233.5	65.6	48.8	59.0	124.9
July ...	4.613	4.780	20	2.036	2.060	1.983	148.2	64.3	50.2	59.4	120.0
Aug. ...	1.256	1.363	8	0.148	0.230	0.211	150.7	63.0	48.8	58.1	118.4
Sept. ...	1.961	2.131	13	0.417	0.388	0.368	110.5	62.9	48.1	55.8	109.3
Oct. ...	1.427	1.530	10	0.592	0.666	0.618	144.8	57.7	42.2	51.9	99.7
Nov. ...	1.687	1.846	9	1.365	1.206	1.129	71.8	48.2	35.3	43.4	77.4
Dec. ...	2.288	2.545	25	2.244	2.362	2.284	37.7	43.0	34.4	40.0	58.7
Total or Mean	25.083	27.198	178	13.014	13.309	13.020	1505.7	55.4	41.6	49.1	97.8

* On January 18 and 19, 1918, the cylinders and tank of 60" gauge were submerged: the figures for the 40" gauge are adopted (2.735") and included in above total.

Mangolds, Barn Field, 1918, 1919, 1920.

Roots since 1856. Mangolds since 1876.
Produce per Acre.

Strip.	Strip Manures.	Cross Dressings.				
		O.	N.	A.	A.C.	C.
		None.	Nitrate of Soda	Ammon. Salts.	Ammon. Salts and Rape Cake.	Rape Cake.
	1918.	Tons.	Tons.	Tons.	Tons.	Tons.
1	Dung only ...	(R. 17 98 (L. 2 79)	33 79 4 17	25 39 3 39	24 45 3 16	24 48 3 69
2	Dung, Super., Potash ...	(R. 25 26 (L. 3 11)	38 58 4 63	34 73 4 45	34 04 5 02	28 30 3 88
4	Complete Minerals ...	(R. 4 61 (L. 0 82)	(R. 28 59 (L. 3 15) (R. 27 65 (L. 3 34)	22 39 2 03	29 65 3 63	16 88 2 11
5	Superphosphate only ...	(R. 5 65 (L. 0 93)	25 18 2 49	12 50 2 56	12 33 2 11	12 50 1 79
6	Super. and Potash ...	(R. 4 58 (L. 0 86)	25 09 2 45	20 30 1 83	25 56 2 94	15 03 1 46
7	Super., Sulphate of Mag., and Sodium Chloride	(R. 4 71 (L. 0 84)	28 81 2 82	23 94 2 41	25 01 3 05	17 21 1 97
8	None ...	(R. 3 18 (L. 0 69)	19 92 2 81	11 05 2 94	10 32 2 53	9 76 1 89
9	Sodium Chloride, Nit. Soda, Sulph. Potash, and Sulph. Mag. ...	(R. 26 56 (L. 2 57)				
	1919					
1	Dung only ...	(R. 9 05 (L. 3 60)	17 49 6 36	14 14 4 55	10 60 4 74	11 28 4 83
2	Dung, Super., Potash ...	(R. 13 97 (L. 5 67)	18 51 8 89	19 87 8 23	15 24 7 60	18 17 6 92
4	Complete Minerals ...	(R. 2 46 (L. 0 97)	(R. 12 98 (L. 6 55) (R. 12 86 (L. 5 69)	7 75 4 43	11 77 6 86	9 57 3 48
5	Superphosphate only ...	(R. 1 97 (L. 0 81)	9 98 4 38	1 13 1 00	3 05 1 86	4 58 1 88
6	Super. and Potash ...	(R. 2 44 (L. 0 91)	14 46 5 81	11 45 4 54	13 14 5 98	12 08 2 84
7	Super., Sulphate of Mag., and Sodium Chloride	(R. 3 13 (L. 0 91)	15 93 5 23	14 48 4 73	14 98 5 31	13 94 2 98
8	None ...	(R. 2 16 (L. 0 82)	7 63 3 63	3 08 1 58	4 06 1 63	6 50 2 57
9	Sodium Chloride; Nit. Soda, Sulph. Potash and Sulph. Mag. ...	(R. 20 38 (L. 5 14)				
	1920.					
1	Dung only ...	(R. 18 99 (L. 3 51)	30 26 4 27	21 38 3 95	23 89 4 62	25 12 5 31
2	Dung, Super., Potash ...	(R. 26 84 (L. 4 78)	37 69 6 74	33 11 6 69	32 67 7 28	28 73 5 94
4	Complete Minerals ...	(R. 4 54 (L. 0 96)	(R. 26 10 (L. 4 68) (R. 21 21 (L. 2 90)	20 81 3 54	26 35 4 75	12 39 2 17
5	Superphosphate only ...	(R. 4 82 (L. 0 95)	20 73 3 33	7 72 2 84	8 31 2 86	8 48 2 13
6	Super. and Potash ...	(R. 4 65 (L. 1 04)	21 50 3 49	18 94 2 84	24 74 4 39	9 89 2 25
7	Super., Sulphate of Mag., and Sodium Chloride	(R. 4 91 (L. 1 28)	21 84 3 27	19 95 3 08	19 05 4 13	11 86 2 41
8	None ...	(R. 3 99 (L. 0 82)	13 81 2 69	5 91 2 19	4 44 1 92	6 17 2 04
9	Sodium Chloride; Nit. Soda; Sulph. Potash and Sulph. Mag. ...	(R. 29 35 (L. 4 86)				

R.—roots. L.—leaves.

Notes: 1918—All Potash, Magnesia, and Rape Cake omitted.

Hay. The Park Grass Plots. 1918, 1919, 1920.

Plot.	Manuring.	Yield of Hay per acre.			Yield of Hay per acre.			Yield of Hay per acre.			Plot.
		1918.			1919.			1920.			
		1st Crop.	2nd Crop.	Total.	1st Crop.	2nd Crop.	Total.	1st Crop.	2nd Crop.	Total.	
1	Single dressing Amm. Salts (=43 lbs. N.) ; (with Dung 8 years, 1856-63)	9.8	8.3	17.1	10.0	7.8	17.8	23.8	10.2	34.0	1
2	Unmanured ; (after Dung 8 years, 1856-63)	17.2	7.6	24.8	10.4	8.5	18.9	24.3	7.8	32.1	2
3	Unmanured	16.5	3.5	20.0	6.4	4.2	10.6	17.5	5.2	22.7	3
4-1	Superphosphate of Lime	16.0	2.5	18.5	6.4	2.0	8.4	19.2	5.5	24.7	4-1
4-2	Superphosphate of Lime and double dressing Amm. Salts (=86 lbs. N.)	14.7	2.8	17.5	4.1	3.7	7.8	15.7	4.1	19.8	4-2
5-1	(N. half) Unmanured ; following double dressing Amm. Salts (= 86 lbs. N.) 1856-97	15.1	3.4	18.5	3.8	2.2	6.0	16.0	4.2	20.0	5-1
5-2	(S. half) Super., Sulphate of Potash ; following double dressing Amm. Salts (= 86 lbs. N.) 1856-97	24.2	4.3	28.5	9.1	4.9	14.0	24.1	6.8	30.9	5-2
6	Complete Mineral Manure as plot 7 ; following double dressing Amm. Salts (= 86 lbs. N.) 1856-68	17.6	3.4	21.0	6.1	1.9	8.0	25.0	6.2	31.2	6
7	Complete Mineral Manure	18.5	5.0	23.5	9.4	1.9	11.3	25.6	5.2	30.8	7
8	Mineral Manure without Potash	26.3	10.5	36.8	17.4	9.0	26.4	31.4	10.0	41.4	8
9	Complete Mineral Manure and double dressing Amm. Salts (= 86 lbs. N.)	12.1	2.8	14.9	4.4	3.8	8.2	10.3	5.4	15.7	9
10	Mineral Manure (without Potash) and double dressing Amm. Salts (= 86 lbs. N.)	24.5	5.8	30.3	13.0	7.3	20.3	20.8	8.8	29.6	10
11-1	Complete Mineral Manure and treble dressing Amm. Salts (= 129 lbs. N.)	41.5	7.9	49.4	23.0	13.1	36.1	35.3	14.8	50.1	11-1
11-2	As plot 11-1 and Silicate of Soda	37.9	7.3	45.2	19.8	13.1	32.9	31.7	14.3	46.0	11-2
		42.3	5.5	47.8	21.1	10.7	31.8	41.8	12.3	54.1	
		21.7	4.2	25.9	9.9	9.8	19.7	19.9	7.4	27.3	
		19.3	4.2	23.5	8.8	6.8	15.6	16.4	7.2	23.6	
		32.5	14.2	46.7	36.5	16.7	53.2	34.3	15.3	49.6	
		46.7	12.8	59.5	50.6	17.1	67.7	42.3	13.1	55.4	
		24.5	9.0	33.5	26.1	8.4	34.5	22.1	10.5	32.6	
		42.8	10.8	53.6	37.1	13.5	50.6	35.3	12.0	47.3	
		44.9	19.0	63.9	35.4	30.1	65.5	41.0	35.4	76.4	
		52.5	17.5	70.0	49.1	18.0	69.1	52.6	21.4	74.0	
		48.0	23.3	71.3	51.8	26.9	78.7	50.9	33.4	84.3	
		49.0	18.6	67.6	57.6	21.5	79.1	52.1	28.1	80.2	

Ground lime was applied to the Southern portion (limed) of the plots at the rate of 2,000 lb. to the acre in the Winter of 1903, 1907, 1915, and at the rate of 2,500 lbs. to the acre in the Winter of 1920, except where otherwise stated. In 1918 all Sulphate of Potash and Sulphate of Magnesia were omitted from the Mineral Manures in plots 6, 7, 9, 11-1, 11-2, 14, 15, 16, and 18; also Potash was omitted from plot 5-2, and Magnesia from plots 8 and 10. In 1919, instead of Sulphate of Potash, an equivalent amount of Muriate of Potash was used. In 1919 and 1920 an equivalent amount of Sulphate of Ammonia was used instead of Muriate of Ammonia. Up to 1914 the limed and unlimed plot results were not separately given in the Annual Report, but the mean of the two was given. From 1915 onwards the separate figures are given. In the reports for 1913, 1914, 1915-17 the manuring of plot 5-2 was incorrectly given as complete Mineral Manure, instead of as Superphosphate and Sulphate of Potash.

9	Complete Mineral Manure, and double Amm. Salts	Limed ...	1st	97.81	—	2.18	95.85	.09	4.05	Rumex acetosa	...	...	9
9	" " " "	" Not limed	2nd	99.26	—	.73	98.40	.97	.61	"	...	...	
10	Mineral Manure (without Potash) and double Amm. Salts	Limed ...	1st	79.91	—	20.10	85.00	—	15.00	Rumex acetosa	...	...	9
10	" " " "	" " " "	2nd	97.34	—	2.65	94.75	—	5.24	"	...	...	
10	" " " "	" " " "	1st	99.68	—	.31	99.59	—	.41	Rumex acetosa	...	...	10
10	" " " "	" " " "	2nd	100.00	—	—	99.77	—	.23	"	...	...	
11-1	Complete Mineral Manure and treble Amm. Salts	Not limed	1st	87.36	—	12.13	92.60	—	7.40	Rumex acetosa	...	...	10
11-1	" " " "	" " " "	2nd	99.59	—	.41	97.36	—	2.63	"	...	...	
11-1	Complete Mineral Manure and treble Amm. Salts	Limed ...	1st	99.63	—	.37	99.84	—	.16	Rumex acetosa	...	...	11-1
11-1	" " " "	" " " "	2nd	100.00	—	—	99.49	—	.52	"	...	...	
11-1	" " " "	Not limed	1st	97.50	—	2.50	98.86	—	1.14	Rumex acetosa	...	...	11-1
11-2	As plot 11-1 and Silicate of Soda	Limed ...	1st	100.00	—	—	99.56	—	.43	"	...	...	
11-2	" " " "	" " " "	2nd	99.59	.06	.35	100.00	—	—	"	...	...	11-2
11-2	" " " "	" " " "	1st	100.00	—	—	99.86	—	.13	"	...	...	
11-2	" " " "	Not limed	2nd	98.54	—	1.47	99.62	—	.38	Rumex acetosa	...	...	11-2
12	Unmanured	" " " "	1st	99.82	—	.18	99.75	—	.25	"	...	...	
12	Dung in 1905 and every 4th year since (omitted in 1917); Fish Guano in 1907 and every 4th year since	Whole plot	2nd	—	—	—	54.80	5.34	39.87	Plantago lanceolata and Conopodium denudatum	...	...	12
13	" " " "	Limed ...	1st	—	—	—	50.56	8.19	41.24	"	...	...	
13	" " " "	" " " "	2nd	—	—	—	86.89	.82	12.28	Rumex acetosa and Conopodium denudatum	...	...	13
13	" " " "	Not limed	1st	—	—	—	2.51	5.37	5.37	"	...	...	
13	" " " "	" " " "	2nd	—	—	—	79.83	.07	20.09	Rumex acetosa and Conopodium denudatum	...	...	13
14	Complete Mineral Manure and double Nitrate of Soda	Whole plot	1st	—	—	—	96.23	.26	3.51	"	...	...	
14	" " " "	" " " "	2nd	—	—	—	92.92	2.41	4.68	Anthriscus sylvestris	...	...	14
15	As plot 7, following double Nitrate of Soda, 1858-75	Whole plot	1st	75.32	6.99	17.70	69.25	5.39	25.35	Rumex acetosa and Achillea millefolium	...	...	15
15	" " " "	" " " "	2nd	73.26	3.15	23.58	77.14	6.54	16.31	"	...	...	
16	As plot 7, and single Nitrate of Soda	Limed ...	1st	—	—	—	89.60	.78	9.62	Anthriscus sylvestris	...	...	16
16	" " " "	" " " "	2nd	—	—	—	90.69	.89	8.42	"	...	...	
16	" " " "	Not limed	1st	—	—	—	85.98	1.17	12.84	Taraxacum vulgare and Plantago lanceolata	...	...	16
16	" " " "	" " " "	2nd	—	—	—	86.60	1.26	12.14	"	...	...	
17	Single Nitrate of Soda	Whole plot	1st	—	—	—	58.53	.43	41.04	Plantago lanceolata and Centaurea nigra	...	...	17
17	" " " "	" " " "	2nd	—	—	—	66.55	.24	33.21	"	...	...	
18	Potash, Sulph. Soda, Magnesia and double Sulph. Amm., 1905 and since	Whole plot	1st	—	—	—	68.93	—	31.07	Rumex acetosa and Conopodium denudatum	...	...	18
18	" " " "	" " " "	2nd	—	—	—	80.04	—	19.95	"	...	...	
19	Farmyard Dung, 1905, and every 4th year since (omitted in 1917)	Whole plot	1st	72.80	6.59	20.61	75.21	6.17	18.61	Rumex acetosa and Ranunculus spp.	...	...	19
19	" " " "	" " " "	2nd	83.97	7.54	8.50	83.43	9.34	7.22	"	...	...	
20	Dung in 1905 and every 4th year since (omitted in 1917). Each intervening year Sulph. Potash, Super. and Nitrate of Soda	Whole plot	1st	78.44	9.26	12.29	81.85	4.72	13.44	Rumex acetosa, Anthriscus sylvestris, Ranunculus spp.	...	...	20
20	" " " "	" " " "	2nd	88.10	4.29	7.61	91.93	4.13	3.94	"	...	...	

WHEAT. BROADBALK FIELD, 1918, 1919, 1920.

1918.										1919.										1920.											
Top portion.					Bottom portion.					Top portion.					Bottom portion.					Top portion.					Bottom portion.						
Dressed Grain.		Straw			Dressed Grain.		Straw			Dressed Grain.		Straw			Dressed Grain.		Straw			Dressed Grain.		Straw			Dressed Grain.		Straw			Average for 61 years, 1852-1912.	
Yield per Acre.	Weight per Bushel.	Yield per Acre.	Weight per Bushel.	per Acre.	Yield per Acre.	Weight per Bushel.	Yield per Acre.	Weight per Bushel.	per Acre.	Yield per Acre.	Weight per Bushel.	Yield per Acre.	Weight per Bushel.	per Acre.	Yield per Acre.	Weight per Bushel.	Yield per Acre.	Weight per Bushel.	per Acre.	Yield per Acre.	Weight per Bushel.	Yield per Acre.	Weight per Bushel.	per Acre.	Yield per Acre.	Weight per Bushel.	Yield per Acre.	Weight per Bushel.	Dressed Grain per Acre.	Straw per Acre.	
Bushels	lb.	cwt.	lb.	cwt.	Bushels	lb.	cwt.	lb.	cwt.	Bushels	lb.	cwt.	lb.	cwt.	Bushels	lb.	cwt.	lb.	cwt.	Bushels	lb.	cwt.	lb.	cwt.	Bushels	lb.	cwt.	lb.	cwt.	Bushels	cwt.
30.8	65.3	38.8	65.1	39.9	36.1	65.1	62.7	19.0	21.6	27.1	62.3	21.6	61.5	38.4	28.3	61.5	38.4	8.1	60.9	33.2	62.0	42.6	35.2	34.8	35.2	42.6	62.0	42.6	35.2	34.8	
11.5	64.1	9.9	64.6	9.1	10.0	64.6	7.0	61.0	7.4	9.8	60.4	7.4	61.0	8.8	9.4	61.0	8.8	8.8	60.9	8.1	60.9	7.8	12.6	10.3	12.6	60.9	7.8	12.6	10.3	12.6	10.3
9.1	64.6	10.0	64.9	10.1	9.0	64.9	60.3	7.8	7.0	10.0	61.5	7.0	61.3	8.3	8.2	61.3	8.3	8.3	62.1	8.2	62.1	8.0	14.5	12.1	14.5	62.1	8.0	14.5	12.1	14.5	12.1
16.5	64.8	23.5	64.9	15.4	19.5	61.1	15.7	60.3	13.9	18.4	61.6	13.9	61.5	16.1	15.3	61.5	16.1	15.1	61.2	15.1	61.2	14.9	23.2	21.4	23.2	61.2	14.9	23.2	21.4	23.2	21.4
25.2	64.6	35.1	65.0	33.8	23.0	65.0	21.4	21.4	21.4	33.3	61.6	24.7	61.3	25.7	21.6	61.3	25.7	30.9	60.9	30.9	60.9	36.5	32.1	32.9	32.1	60.9	36.5	32.1	32.9	32.9	32.9
27.7	65.1	45.6	65.3	44.2	29.6	61.7	29.3	29.3	29.6	37.4	61.7	29.6	61.8	45.4	28.8	61.8	45.4	28.8	60.4	28.8	60.4	42.3	36.6	41.1	36.6	60.4	42.3	36.6	41.1	41.1	41.1
21.5	64.9	30.7	65.4	25.9	21.0	60.2	17.5	60.2	18.8	23.9	61.0	18.8	61.1	25.7	23.6	61.1	25.7	26.8	61.6	26.8	61.6	27.4	—	—	—	61.6	27.4	—	—	—	—
13.8	64.5	16.9	64.7	18.8	16.4	59.8	13.6	59.8	13.1	15.2	58.0	13.1	59.5	19.0	13.0	59.5	19.0	13.0	60.0	13.0	60.0	18.0	20.0	18.4	20.0	60.0	18.0	20.0	18.4	18.4	18.4
17.9	64.5	21.7	64.7	21.1	10.7	57.6	16.1	57.6	15.7	12.4	57.0	15.7	59.7	19.1	13.6	59.7	19.1	9.6	58.9	9.6	58.9	21.5	22.9	22.3	22.9	58.9	21.5	22.9	22.3	22.3	22.3
20.8	64.6	25.8	63.9	27.4	17.3	58.2	16.2	58.2	19.9	18.5	57.8	19.9	60.3	27.9	19.3	60.3	27.9	18.4	60.8	18.4	60.8	28.1	29.1	28.0	29.1	60.8	28.1	29.1	28.0	28.0	28.0
20.3	65.0	27.2	64.7	31.6	29.3	60.6	25.9	60.6	24.8	27.4	60.8	24.8	61.4	29.6	24.9	61.4	29.6	28.9	61.1	28.9	61.1	39.0	31.0	31.5	31.0	61.1	39.0	31.0	31.5	31.5	31.5
21.4	64.9	24.2	64.7	32.1	17.7	58.3	19.2	58.3	23.6	27.5	59.3	23.6	59.8	26.1	15.5	59.8	26.1	18.2	59.7	18.2	59.7	28.8	28.8	28.0	28.8	59.7	28.8	28.8	28.0	28.0	28.0
25.9	65.0	33.0	65.1	31.5	22.5	60.3	18.3	60.3	15.8	20.7	60.4	15.8	61.1	24.3	17.6	61.1	24.3	16.1	61.1	16.1	61.1	23.0	29.9	29.7	29.9	61.1	23.0	29.9	29.7	29.7	29.7
29.7	65.4	41.8	65.5	41.6	32.8	60.9	29.6	60.9	29.7	32.6	61.3	29.7	61.4	30.8	25.2	61.4	30.8	25.5	61.2	25.5	61.2	35.0	—	—	—	61.2	35.0	—	—	—	—
17.5	65.0	20.3	65.1	28.8	11.7	62.9	8.1	62.9	9.4	13.8	62.4	9.4	61.3	19.6	19.1	61.3	19.6	20.0	61.5	20.0	61.5	21.1	29.9	29.5	29.9	61.5	21.1	29.9	29.5	29.5	29.5
12.4	65.0	10.1	65.4	12.6	30.2	60.6	21.9	60.6	25.3	30.0	61.0	25.3	58.4	6.9	6.3	58.4	6.9	6.4	59.9	6.4	59.9	7.7	14.9	13.0	14.9	59.9	7.7	14.9	13.0	13.0	13.0
17.0	65.0	13.7	64.9	16.8	13.5	61.5	9.5	61.5	11.2	12.0	61.6	11.2	60.3	10.9	11.8	60.3	10.9	13.0	60.4	13.0	60.4	14.3	*25.4	*25.7	*25.4	60.4	14.3	*25.4	*25.7	*25.7	*25.7
17.3	65.3	19.8	—	—	21.7	61.5	18.6	61.5	—	—	—	—	59.6	18.7	15.4	59.6	18.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...		

* Average for 20 years, 1893-1912.

Average for 20 years, 1893-1912.

1918, 1919, Sulphate of Potash omitted from plots 5, 6, 7, 8, 9, 13, 15, 16, 17 or 18, 20. Sulphate of Soda omitted from plot 12.

Sulphate of Magnesia omitted from plot 14.

1918, 1919, 1920. Rape Cake omitted from plot 19.

Plot 17 received Mineral dressing in 1919. Amm. Salts in 1918, 1920.

Plot	Received mineral dressing in 1919.	Received mineral dressing in 1920.	Received mineral dressing in 1921.
Plot 18	"	"	"
Plot 19	"	"	"
Plot 20	"	"	"
Plot 21	"	"	"
Plot 22	"	"	"
Plot 23	"	"	"
Plot 24	"	"	"
Plot 25	"	"	"
Plot 26	"	"	"
Plot 27	"	"	"
Plot 28	"	"	"
Plot 29	"	"	"
Plot 30	"	"	"
Plot 31	"	"	"
Plot 32	"	"	"
Plot 33	"	"	"
Plot 34	"	"	"
Plot 35	"	"	"
Plot 36	"	"	"
Plot 37	"	"	"
Plot 38	"	"	"
Plot 39	"	"	"
Plot 40	"	"	"
Plot 41	"	"	"
Plot 42	"	"	"
Plot 43	"	"	"
Plot 44	"	"	"
Plot 45	"	"	"
Plot 46	"	"	"
Plot 47	"	"	"
Plot 48	"	"	"
Plot 49	"	"	"
Plot 50	"	"	"
Plot 51	"	"	"
Plot 52	"	"	"
Plot 53	"	"	"
Plot 54	"	"	"
Plot 55	"	"	"
Plot 56	"	"	"
Plot 57	"	"	"
Plot 58	"	"	"
Plot 59	"	"	"
Plot 60	"	"	"
Plot 61	"	"	"
Plot 62	"	"	"
Plot 63	"	"	"
Plot 64	"	"	"
Plot 65	"	"	"
Plot 66	"	"	"
Plot 67	"	"	"
Plot 68	"	"	"
Plot 69	"	"	"
Plot 70	"	"	"
Plot 71	"	"	"
Plot 72	"	"	"
Plot 73	"	"	"
Plot 74	"	"	"
Plot 75	"	"	"
Plot 76	"	"	"
Plot 77	"	"	"
Plot 78	"	"	"
Plot 79	"	"	"
Plot 80	"	"	"
Plot 81	"	"	"
Plot 82	"	"	"
Plot 83	"	"	"
Plot 84	"	"	"
Plot 85	"	"	"
Plot 86	"	"	"
Plot 87	"	"	"
Plot 88	"	"	"
Plot 89	"	"	"
Plot 90	"	"	"
Plot 91	"	"	"
Plot 92	"	"	"
Plot 93	"	"	"
Plot 94	"	"	"
Plot 95	"	"	"
Plot 96	"	"	"
Plot 97	"	"	"
Plot 98	"	"	"
Plot 99	"	"	"
Plot 100	"	"	"

From 1917 onwards, Muriate of Ammonia has been replaced by an equivalent amount of Sulphate of Ammonia.

Hoos Field (formerly Potato Plots). No Maure since 1901.

Plot.	Manuring given prior to 1901.	1918. BARLEY.				1919. BARLEY.			
		Dressed Grain.		Straw per Acre.	Total Produce per Acre.	Dressed Grain.		Straw per Acre.	Total Produce per Acre.
		Yield per Acre.	Weight per Bush.			Yield per Acre.	Weight per Bush.		
Previous Cropping: Potatoes, 1876-1901; Barley, 1902 and 1903; Oats, 1904; Barley, 1905-1911; Oats, 1912; Barley, 1913 and 1914; Oats, 1915; Barley, 1916-17.									
1	Unmanured	...	...	...	Bush.	lb.	cwt.	lb.	Total Produce per Acre.
2	Unmanured 1882 to 1901, previously Dung only	...	...	...	8.4	52.4	4.0	52.3	645
3	Dung 1883 to 1901	...	...	...	11.1	51.8	5.7	53.1	904
4	Dung 1883 to 1901	...	...	...	16.2	54.8	8.6	53.1	1379
		...	...	...	16.3	53.1	7.5	52.9	1475
Previous Cropping: Potatoes, 1876-1901; Barley, 1902-1903; Oats, 1904; Plots 5, 7, 9, Cow Peas (failed), 1905; Plots 6, 8, 10, Red Clover, 1905; Red Clover, 1906-1911; Oats, 1912; Barley, 1913 and 1914; Oats, 1915; Barley, 1916-17.									
5	Ammonium Salts ...	...	...	...	Bush.	lb.	cwt.	lb.	Total Produce per Acre.
6	Nitrate of Soda ...	...	...	...	11.6	54.5	7.0	51.3	1004
7	Ammonium Salts and Mixed Minerals	...	...	...	15.8	54.1	7.1	51.8	1014
8	Nitrate of Soda and Mixed Minerals	...	...	...	21.0	54.1	11.1	53.1	1339
9	Superphosphate ...	...	...	...	19.2	53.5	8.2	53.0	1150
10	Mixed Minerals ...	...	...	...	14.4	54.1	7.6	53.6	965
		...	...	...	14.8	53.4	6.7	53.5	1278

NOTE.—In 1920 these plots were fallowed.

Permanent Barley Plots. Hoos Field, 1918, 1919, 1920. PRODUCE PER ACRE

Plot.	MANURING.	1918			1919			1920			Average 60 years, 1852—1911.	
		Dressed Grain.	Weight per Bushel.	Straw, cwt.	Dressed Grain.	Weight per Bushel.	Straw, cwt.	Dressed Grain.	Weight per Bushel.	Straw, cwt.	Dressed Grain.	Straw, cwt.
1 O	Unmanured	Bush. 18.3	lb. 52.1	7.9	Bush. 5.5	lb. 52.0	3.6	Bush. 6.4	lb. 55.4	3.2	Bush. 14.3	cwt. 8.4
2 O	Superphosphate only	27.7	51.7	11.0	11.6	54.8	5.7	13.2	53.4	7.5	19.7	10.0
3 O	Alkali Salts only	16.9	51.7	7.9	9.0	54.4	5.8	9.1	52.9	4.7	15.2	8.8
4 O	Complete Minerals	24.9	52.2	11.5	14.6	54.1	8.4	13.7	52.6	6.3	19.7	11.1
5 O	Potash and Superphosphate	14.7	51.5	7.3	9.4	55.0	6.1	5.2	54.5	3.4		
1 A	Ammonium Salts only	25.1	50.7	11.5	11.2	52.5	6.5	17.3	51.2	8.1	25.5	14.7
2 A	Superphosphate and Amm. Salts	41.4	50.6	17.7	18.1	51.4	9.1	22.8	51.7	10.7	38.2	22.0
3 A	Alkali Salts and Amm. Salts	23.4	52.1	11.2	15.7	53.4	9.1	16.1	53.1	9.6	28.0	16.9
4 A	Complete Minerals and Amm. Salts	34.9	51.7	15.7	24.5	54.6	12.5	38.5	52.7	14.9	41.5	25.0
5 A	Potash, Super. and Amm. Salts	38.6	52.1	20.3	23.5	54.6	14.0	30.4	53.4	16.3		
1 AA	Nitrate of Soda only	26.7	51.3	14.7	16.1	53.3	9.7	20.2	53.3	12.1	29.3	17.8
2 AA	Super. and Nitrate of Soda	46.4	52.5	22.6	30.1	54.1	14.4	37.3	53.3	16.1	43.1	26.3
3 AA	Alkali Salts and Nitrate of Soda	21.1	52.0	13.3	16.7	54.1	11.3	14.7	53.1	12.3	30.0	19.3
4 AA	Complete Minerals and Nitrate of Soda	43.3	50.6	16.9	28.0	54.1	15.1	31.8	54.1	15.3	42.7	27.3
1 AAS	As Plot 1 AA and Silicate of Soda	29.2	51.6	14.5	20.3	54.0	12.9	27.7	53.8	14.8	32.8 (1)	19.7 (1)
2 AAS	" " 2 AA "	45.1	52.7	21.7	27.8	52.9	12.0	39.4	53.7	15.7	42.3 (1)	26.0 (1)
3 AAS	" " 3 AA "	24.0	52.7	16.1	19.8	54.0	12.6	23.9	52.8	15.8	35.2 (1)	21.7 (1)
4 AAS	" " 4 AA "	39.5	51.3	20.3	20.5	54.8	12.2	30.5	54.3	15.5	43.6 (1)	27.7 (1)
1 C	Rape Cake only	18.7	52.5	9.3	10.7	53.9	5.6	11.9	53.9	5.8	38.3	22.1
2 C	Superphosphate and Rape Cake	21.9	52.0	11.1	11.7	54.6	6.7	12.1	53.4	5.6	40.5	23.6
3 C	Alkali Salts and Rape Cake	16.6	51.3	9.2	8.5	54.5	5.3	9.7	54.0	4.8	36.9	22.3
4 C	Complete Minerals and Rape Cake	17.3	52.6	8.9	10.0	54.1	6.2	10.2	54.5	4.5	40.5	24.5
7-1	Unmanured (after dung 20 years, 1852—71)	26.7	52.4	12.0	12.1	54.5	6.5	17.5	53.6	8.0	24.8 (2)	14.8 (2)
7-2	Farmyard Manure	58.8	53.0	28.7	32.1	55.2	18.2	48.3	54.3	27.3	47.1	29.6
6-1	Unmanured	20.9	52.3	9.2	7.6	53.5	4.3	13.3	53.5	7.3		
6-2	Ashes from Laboratory Furnace	19.5	50.6	8.5	6.8	53.5	3.9	10.2	53.8	5.7		
1 N	Nitrate of Soda only	26.3	50.6	14.0	12.5	53.0	8.6	10.3	53.3	6.4		
2 N	" " "	27.8	52.1	15.9	19.5	54.9	11.4	20.3	53.0	10.5		

(1) 48 years, 1864—1911. (2) 40 years, 1872—1911.

NOTES:—1918. Sulphate of Potash, Sulphate of Magnesia and Rape Cake omitted in all cases.
1919 and 1920. Rape Cake omitted in all cases. Also since 1917 Murrate of Ammonia replaced by equivalent amount of Sulphate of Ammonia.

Little Hoos Field

PLAN OF ROTATION PLOTS

Arranged to test the RESIDUAL VALUE of VARIOUS MANURES in one, two, three, and four years after their application. Produce per acre.

Plot.	Manure per Acre.	Year of Last Dressing.	1918 (15th Season). Wheat.			1919 (16th Season). Barley.			1920 (17th Season). Swedes.		
			Dress'd Grain Bush. per Acre.	Straw cwt. per Acre.	Total Pr'd'ce lb. per Acre.	Dr'ss'd Grain Bush. per Acre.	Straw cwt. per Acre.	Total Pr'd'ce lb. per Acre.	Roots tons per Acre.	Leav's tons per Acre.	Total tons per Acre.
A 1	Control	—	36.3	36.7	6613	10.8	8.7	1669	9.43	2.60	12.03
2	Ordinary Dung, 16 tons	1920	42.5	45.7	8129	27.9	17.2	3554	18.38	3.10	21.48
3		1913	44.3	42.9	7921	27.4	16.1	3390	12.02	2.45	14.47
4		1914	43.2	42.8	7812	25.8	14.7	3128	10.43	2.17	12.60
5		1915	42.1	41.8	7685	29.3	18.0	3758	13.40	2.84	16.24
B 1	Cake fed dung, 16 tons	1920	44.8	46.3	8288	28.6	16.9	3519	21.74	3.73	25.47
2	Control	—	38.2	36.7	6831	11.7	10.5	2075	8.42	2.32	10.74
3	Cake fed dung, 16 tons	1913	46.8	47.3	8550	28.8	15.9	3462	15.20	3.00	18.20
4		1914	44.2	43.5	7974	29.8	16.7	3546	16.89	3.39	20.28
5		1915	44.6	44.0	8024	29.6	17.1	3553	14.80	2.88	17.68
C 1	Shoddy, 308 lb.; Super. 292 lb.; Sulph. of Potash 110 lb. ...	1920	33.6	32.6	6071	11.4	9.0	1715	11.93	3.62	15.55
2	Control	1913	32.9	32.9	6097	14.3	9.8	1985	10.54	3.58	14.12
3		—	36.5	33.5	6344	13.4	8.4	1756	13.66	3.65	17.31
4		1914	34.9	36.6	6635	17.2	10.5	2194	12.62	3.33	15.95
5		1919	37.5	37.2	6765	23.3	14.5	2996	14.57	3.50	18.07
D 1	Guano 352 lb.; Sulph. Amm. 44 lb. ...	1920	38.3	39.2	7119	14.9	10.0	1968	14.44	4.25	18.69
2	Sulph. of Potash 86 lb. ...	1913	34.8	31.8	6041	12.9	8.4	1735	12.71	3.47	16.18
3		1914	35.5	37.9	6839	17.1	10.4	2171	14.43	4.18	18.61
4		—	37.2	41.2	7258	13.6	8.6	1809	9.61	2.99	12.60
5	Guano 352 lb.; Sulph. Amm. 44 lb.; Sulph. of Potash 86 lb. ...	1919	37.4	38.5	7004	24.3	17.0	3412	5.80	2.22	8.02
E 1	Rape Dust 844 lb.; Super. 240 lb. ...	1920	38.1	38.2	7003	13.7	9.5	1911	13.33	4.14	17.47
2	Sulph. of Potash 80 lb. ...	1913	37.1	40.3	7166	15.7	10.5	2162	14.05	3.13	17.18
3		1914	34.3	34.7	6309	14.5	8.7	1881	14.17	3.42	17.59
4		1919	34.4	37.3	6609	22.4	13.5	2871	10.90	2.78	13.68
5	Control	—	39.3	39.5	7207	15.7	11.0	2161	6.43	2.31	8.74
F 1	Control	—	35.6	37.3	6703	10.8	8.4	1629	3.32	1.46	4.78
2	Super. 292 lb.; Sulph. Amm. 196 lb.; Sulph. of Potash 110 lb. ...	1920	36.7	36.2	6675	12.3	8.9	1805	16.30	3.46	19.76
3		1913	31.8	33.0	5995	11.7	8.1	1673	9.31	2.55	11.86
4		1914	35.5	37.5	6751	12.4	8.2	1696	7.84	2.15	9.99
5		1919	36.1	37.7	6788	23.9	16.3	3276	9.23	2.75	11.98
G 1	Bone Meal 160 lb.; Super. 110 lb.; Sulph. Amm. 188 lb. ...	1920	34.9	33.9	6321	14.8	9.6	1952	8.63	3.10	11.73
2	Control	1913	33.0	32.7	6025	16.5	9.9	2060	6.27	2.01	8.28
3		—	34.6	35.2	6374	17.6	9.6	2118	3.60	1.11	4.71
4		1914	37.5	36.8	6714	16.9	10.2	2131	6.00	1.74	7.74
5		1919	36.0	37.7	6780	23.2	14.6	3022	7.12	2.31	9.43
H 1	Basic Slag 520 lb.; Super. 110 lb.; Sulph. Amm. 196 lb. ...	1920	41.6	37.4	7016	24.4	13.3	2867	15.77	3.58	19.35
2	Control	1913	42.3	41.1	7485	24.0	13.0	2816	11.48	2.87	14.35
3		1914	39.6	40.6	7303	23.2	12.8	2741	11.48	2.85	14.33
4		1919	40.6	37.0	6934	31.2	17.4	3750	10.72	2.58	13.30
5	Control	—	36.8	38.0	6837	19.8	11.3	2431	4.87	1.54	6.41

NOTES AS TO MANURES.

In 1919 a new system of manuring was begun. The manure for each plot (except of series A and B) was rationed at 40 lbs. Nitrogen, 100 lbs. Calcium Phosphate, and 50 lbs. Potash per acre. Each plot was supplied with as much of its particular manure (shoddy, guano, &c.) as possible without exceeding the receipt in any of the three rationed ingredients. Any deficit in either of these three was then made good by adding the necessary quantity of Sulphate of Amm., Superphosphate, or Sulphate of Potash. Series A and B left as before. No manure was applied in 1917 or 1918. For manures 1904-17 see Report for 1915-16-17.

Figures in italics denote unmanured plots. The yields on the plots to which the manure was applied in a given year are printed in heavy type.

RED CLOVER grown year after year on rich Garden Soil, Rothamsted Garden.

Hay, Dry Matter, and Nitrogen per Acre, 1913 to 1920.

Year.	No. of Cuttings.	As Hay.	Dry Matter.	Nitrogen.	Seed Sown.
		lbs.	lbs.	lbs.	
1913	2	4211	3509	98	1912. April, mended
1914	2	2041	1701	46	" " "
1915	1	1304	1087	26	" " "
1916	1	1724	1437	51	1916. April 21st, re-sown
1917	3	3351	2793	81	1917. April 23rd, mended
1918	2	2262	1885	50	1918. April 6th, re-sown
1919	2	898	748	22	1919. April 27th, mended
1920	3	4400	3667	114	1920. May 5th, mended
Averages:					
25 years, 1854—1878		7664	6387	179	
25 years, 1879—1903		3924	3270	101	
50 years, 1854—1903		5794	4829	140	
15 years, 1904—1918		2888	2407	70	

Wheat after Fallow (without Manure 1851, and since). Hoos Field, 1918, 1919, 1920.

	1918.	1919.	1920.	Average 61 years 1856-1916.
Dressed Grain { Yield—Bushels per Acre	15.3	11.8	9.4	15.6
{ Weight per Bushel—lbs.	61.7	59.9	62.8	59.5
Straw—cwt. per Acre	14.1	9.6	8.9	13.4
Total Produce—lbs. per Acre	2611	1848	1642	2477

DRESSED SEED EXPERIMENT, 1919. Barley. Little Hoos Field.

Description of Plot.	Dressed Grain.				Straw per Acre.		Total Produce per Acre.	
	Yield per Acre.		Weight per Bushel.					
	Single Strength.	Double Strength.	Single Strength.	Double Strength.	Single Strength.	Double Strength.	Single Strength.	Double Strength.
	Bushels.	Bushels.	lbs.	lbs.	cwt.	cwt.	lbs.	lbs.
Heavy Oil ... {	23.5	20.0	54.0	54.5	13.2	11.8	2880	2580
	24.3	—	54.3	—	14.1	—	3055	—
Bone Oil ... {	19.3	17.5	54.5	54.0	11.6	10.4	2465	2235
	24.5	—	54.5	—	14.5	—	3110	—
Creosote ... {	21.5	11.3	53.3	53.0	12.3	8.8	2695	1750
	22.6	—	53.0	—	14.1	—	2920	—
Acetone Tar {	21.1	12.5	54.5	53.0	12.5	9.1	2695	1770
	19.9	—	52.5	—	12.9	—	2605	—
Gas Tar ... {	23.0	14.1	53.0	54.0	13.2	7.9	2810	1743
	13.1	—	53.5	—	12.9	—	2260	—
	22.0	14.7	55.0	53.0	12.9	9.1	2810	1885
Control ... {	18.9	—	54.0	—	10.0	—	2300	—
	23.7	—	54.0	—	13.8	—	2935	—
	18.4	—	54.5	—	11.8	—	2460	—

Single Strength represents 1 pint of dressing to 4 bushels of seed.

TOP DRESSING EXPERIMENT.

Oats (Grey Winter). Great Harpenden Field, 1919.

Manures per Acre.	Dressed Grain.						Straw per Acre.			Total Produce per Acre.		
	Yield per Acre.			Weight per Bushel.								
	1st Expt.	2nd Expt.	3rd Expt.	1st Expt.	2nd Expt.	3rd Expt.	1st Expt.	2nd Expt.	3rd Expt.	1st Expt.	2nd Expt.	3rd Expt.
	Bush.	Bush.	Bush.	lbs.	lbs.	lbs.	cwt.	cwt.	cwt.	lbs.	lbs.	lbs.
Sulphate Amm. $1\frac{1}{2}$ cwt., Super 3 cwt. ...	79.9	62.6	62.3	42.0	42.8	41.8	40.4	34.2	32.4	8206	6850	6675
Nitrate Soda 2 cwt., Super 3 cwt. ...	71.9	68.9	67.6	42.8	41.9	42.1	41.7	38.0	37.5	8169	7500	7284
Nitrate Amm. $\frac{3}{4}$ cwt., Super 3 cwt. ...	74.1	68.9	57.4	42.5	42.4	44.0	37.7	34.4	32.6	7700	7119	6544
Nitrolim 2 cwt., Super 3 cwt. ...	67.1	58.4	60.3	42.0	42.6	42.0	33.9	29.0	30.1	6900	6069	6706
Guandine Nitrate 84 lbs., Super 3 cwt. ...	75.7	—	63.1	41.4	—	42.9	37.3	—	32.6	7547	—	6678
Guandine Sulphate 94 lbs., Super 3 cwt. ...	73.3	56.7	53.0	41.3	45.0	43.5	32.4	27.7	27.0	6900	5972	5706
Guandine Carb. 75 lbs., Super 3 cwt. ...	68.2	61.5	52.5	42.1	42.8	43.3	30.6	28.4	26.6	6638	6200	5547
Super 3 cwt. ...	64.1	49.0	—	42.0	42.9	—	30.4	25.5	—	6388	5269	—
Control ...	68.1	49.7	48.6	42.9	44.3	43.0	33.3	24.3	26.3	6981	5238	5444
Control ...	58.9	47.8	47.4	43.6	41.6	43.0	28.4	23.4	23.7	6056	4781	4975
Control ...	—	48.1	39.0	—	46.8	42.5	—	24.3	22.8	—	5256	4637

Wheat (Red Standard). Great Harpenden Field, 1920.

Date of Applying Dressing.	Dressed Grain.				Straw per Acre.		Total Produce per Acre.	
	Yield per Acre.		W'ght per Bush		Single Dressing.	Double Dressing.	Single Dressing.	Double Dressing.
	Single Dressing.	Double Dressing.	Single Dressing.	Double Dressing.				
	Bush.	Bush.	lbs.	lbs.	cwt.	cwt.	lus.	lbs.
Early : Feb. 10th ...	28'7	35'9	63'6	63'6	26'9	35'9	4960	6456
Medium : March 16th ...	29'8	—	63'8	—	31'1	—	5522	—
Late : May 10th ...	31'6	32'6	62'9	62'7	33'6	36'9	6020	6490
Control ...	28'9	—	63'9	—	24'2	—	4683	—

Single dressing represents 100 lbs. Sulphate Amm. and 100 lbs Super.

SUBSOILING EXPERIMENT.

Potatoes (King Edward). West Barnfield, 1918.

Treatment of Plots.								Yield per Acre.	
								East. cwt.	West. cwt.
Subsoiled in 1914 ...	...	...	...	...	...	...	...	75.1	127.3
Not Subsoiled ...	...	...	...	...	...	...	...	90.7	133.7

VARIETY EXPERIMENT.

Wheat. Great Harpenden Field, 1920.

Variety .				Dressed Grain.		Straw per Acre.	Total Produce per Acre.
				Yield per Acre.	Weig't per Bush		
				Bushels.	lbs.	cwt.	lbs.
Red Standard	...	...	26.2	63.6	32.2	5475	
Yeoman	...	...	27.0	63.3	29.7	5333	
Fenman	...	...	28.6	62.8	34.3	5845	

FLUE DUST EXPERIMENTS. Mangolds. Stackyard Field, 1918.

Plot.	Manures per Acre.	Weight of	"Best Rows." ¹	
		Roots per Acre.	Number of Rows.	Weight per Acre.
		Tons.		Tons.
10	Superphosphate 4 cwt., Salt 2 cwt. ...	16.7	8	19.6
12A		16.9	7	25.1
14		17.9	5	21.9
1	Superphosphate 4 cwt., Salt 2 cwt., Sulphate	19.3	5	26.8
9		17.9	9	20.9
11	Superphosphate 4 cwt., Salt 2 cwt., Nitrate	20.7	8	25.2
15	Ammonia 145 lbs. ...	25.9	all	25.9
2	Super. 4 cwt., Salt 2 cwt., Sulphate Amm.			
	2 cwt., Flue Dust, grade 1, 3.1 cwt. ...	14.2	4	22.2
3	Ditto, Flue Dust, grade 2, 7.5 cwt. ...	11.9	2	23.6
4	Ditto, Flue Dust, grade 3, 5 cwt. ...	15.5	4	21.4
5	Ditto, Extracted Flue Dust, 6.5 cwt. ...	19.9	all	19.9
6	Ditto, Sulphate of Potash, 1 cwt. ...	18.6	7	20.2
7	Ditto, Flue Dust, grade 2, 7.5 cwt. (Inter- mediate application) ...	15.5	4	28.9
8	Ditto, Flue Dust, grade 2, 7.5 cwt. (late application) ...	17.3	5	23.1
12	Super. 4 cwt., Salt 2 cwt., Dried Sewage Sludge 2 tons ...	20.2	8	24.4
13	Super. 4 cwt., Salt 2 cwt., Cordite 12 cwt.	21.5	7	23.9
16	No Artificials and no Chalk ...	18.3	10	21.3
17		18.4	9	19.9
19	No Chalk, Manure as for farm ...	18.3	7	20.2
20	Chalked, but no Artificials ...	15.8	all	15.8
21		15.8	all	15.8

¹ There were gaps in these plots. "Best Rows" are rows of full length with all plants growing.

Potatoes. West Barnfield, 1918.

Plot.	Manures per Acre.		Weight per Acre Tons.
1	Superphosphate 4 cwt., Sulphate of Ammonia 2 cwt. ...	...	7.5
2	Super. 4 cwt., Sulphate Amm. 2 cwt., Flue Dust, grade 1, 20.7 cwt. ...	...	7.5
3	Ditto ditto	Flue Dust, grade 2, 7.4 cwt.	8.4
4	Ditto ditto	Flue Dust, grade 3, 3.7 cwt.	8.2
5	Ditto ditto	Flue Dust extracted, 6.4 cwt.	8.3
6	Ditto ditto	Sulphate of Potash, 1 cwt.	8.4
7	Ditto ditto	Flue Dust, grade 2, 7.4 cwt. (Intermediate application)	8.4
8	Ditto ditto	Flue Dust, grade 2, 7 cwt. (Late application)	9.0
9	Ditto ditto	...	8.8
10	Ditto ...	...	8.7
11	Ditto, Nitrate of Amm. 145 lbs. ...	...	8.9
12	Ditto, Sewage Sludge, 2 tons ...	...	8.6
12A	Ditto ...	...	7.8
13	Ditto ...	...	7.3
14	Ditto, Nitrate Amm. 145 lbs. ...	...	8.5
15	No Artificials ...	...	7.2

Flue Dust, grade 1, contains 2.21 p.c. Potash.

Flue Dust, grade 3, contains 8.90 p.c. Potash.

" 2, " 5.85 " extracted, " 7.37 "

Sulphate of Potash contains 50.24 p.c. Potash.

The quantities applied were calculated on the basis of 1 cwt. Sulphate of Potash
(49 p.c. Potash) per Acre.

NOTE.—All Plots received a dressing of Dung at the rate of 10 tons per acre.

SLUDGE EXPERIMENTS, 1920.

Hay. Great Field Pasture, 1920.

Plot.	Manures per Acre.	Yield per Acre.
1 North	Wet Sludge, 61·7 cwt.	cwt. 29·3
2	Control	22·0
3 South	Wet Sludge, 61·7 cwt.	22·6
4 North	Sulphate of Ammonia, 1½ cwt.	35·4
5	Control	22·2
6 South	Sulphate of Ammonia, 1½ cwt.	31·0
7 North	Slag, 10 cwt.	25·0
8	Control	21·2
9 South	Slag, 10 cwt.	26·3
10 North	Superphosphate, 6 cwt.	26·3
11	Control	22·9
12 South	Superphosphate, 6 cwt.	22·7
13 North	Nitrate of Ammonia, 114 lbs.	39·6
14	Control	24·9
15 South	Nitrate of Ammonia, 114 lbs.	36·5
16	Nitrolim, 234 lbs.	34·6
17	Control	26·6
18	Nitrolim, 234 lbs.	30·1

Potatoes. Long Hoos Field, 1920.

		tons
1	Activated Sewage Sludge, 13·3 tons; Super., 6 cwt.	11·8
4		8·8
3	Farmyard Dung, 15 tons; Super., 6 cwt.; Nitrate of Ammonia, 1 cwt.	10·8
6		9·6
2	Control; Super., 6 cwt.; Nitrate of Ammonia, 1 cwt.	7·8
5		8·3
7		8·9
8		7·9

Barley. Long Hoos Field, 1920.

Plot.	Manures per Acre.	Dressed Grain.		Straw per Acre.	Total Produce per Acre.
		Yield per Acre.	Weight per Bushel.		
		Bushels	lbs.	cwt.	lbs.
1	Activated Sewage Sludge, 2·7 tons	36·2	55·5	20·4	4363
4		26·3	56·5	21·1	3897
7		46·3	55·8	28·7	5894
2	Sulphate of Ammonia, 1·45 cwt.	45·1	56·3	25·1	5444
5		38·8	56·3	29·1	5513
3		37·0	55·8	21·8	4557
6	Control	36·5	55·5	23·1	4701
8		39·3	55·5	24·7	5057

Wheat. Little Knott Wood, 1918.

These plots were top dressed on March 12th with $1\frac{1}{4}$ cwt. Super. and 1 cwt. Sulphate Amm. per Acre.

Barley. Stackyard Field, 1919.

STRAW EXPERIMENTS.

NOTE.—Manures as follows were applied to the West Barnfield potato plots viz., 2½ cwt. Super., 1½ cwt. Sulphate of Amm., and ½ bush. Bone Meal per acre.

PROFESSOR BLACKMAN'S ELECTRO CULTURE EXPERIMENTS, 1919-20.

Clover. Foster's Field.

Description of Plots.	1919			1920.
	1st Crop.	2nd Crop.	1st and 2nd Crops.	1st Crop.
	Weight per Acre.	Weight per Acre.	Weight per Acre.	Weight per Acre.
	cwts.	cwts.	cwts.	cwts.
Electro Plot ...	34.8	17.1	51.9	23.9
Control 1* ...	23.1	14.0	37.1	24.1
Control 2* ...	36.5	—	48.1	23.0
Control 3 ...	—	11.0	—	—

* Control 2 could not be used for second crop of 1919; Control 3 was therefore added. Control 1 is the same for both 1919 crops, but was wired off for the second crop. Controls 1 and 2 both different in 1920 from 1919.
NOTE.—2nd crop 1920 was ploughed in.

WHEAT AND BARLEY.

Description of Plots.		Dressed Grain.		Straw. Yield per Acre. cwts.	Total Produce. Yield per Acre. lbs.	
		Yield per Acre. Bushels.	Weight per Bushel. lbs.			
Winter Wheat. Great Knott Wood Field, 1919.						
Electro Plots	...	{ E 1	21.2	60.4	15.6	3210
		{ E 2	21.9	61.0	18.1	3596
		{ C 1	13.9	61.9	11.6	2338
Control Plots	...	{ C 2	16.8	62.3	13.5	2833
	Cage Plot ...	...	8.6	60.5	12.0	2108
Spring Wheat. Great Knott Wood Field, 1919.						
Electro Plots	...	{ E 3	8.2	56.0	10.7	1845
		{ E 4	8.2	56.2	11.3	1988
		{ C 3	10.6	56.6	10.3	1951
Control Plots	...	{ C 4	8.7	55.0	11.4	1978
		{ C 5	6.9	55.1	9.8	1676
Wheat (Yeoman). Foster's Field, 1920.						
Electro Plots	...	{ E 1	18.8	62.2	19.0	3448
		{ E 2	18.4	62.2	19.7	3536
Control Plots	...	{ C 1	20.4	62.5	20.5	3697
		{ C 2	18.2	62.1	17.4	3245
Barley. Foster's Field, 1918.						
Electro Plots	...	{ E 1	44.7	51.5	22.4	4890
		{ E 2	47.4	54.0	25.1	5458
		{ E 3	46.4	53.4	24.4	5284
Control Plots	...	{ C 1	36.4	52.6	22.1	4456
		{ C 2	52.7	53.5	29.1	6162
		{ C 3	36.3	54.0	22.3	4525
Cage Plot ...	...	C 4	44.0	54.9	26.3	5453
Barley. Great Knott Wood Field, 1920.						
Electro Plots	...	{ E 3	31.7	53.1	17.3	3672
		{ E 4	33.0	53.2	18.5	3889
Control Plots	...	{ C 3	29.5	53.4	16.0	3410
		{ C 4	25.2	53.3	12.5	2779

SOIL STERILISING EXPERIMENT.

Wheat after Barley. Long Hoos Field, 1918.

Plot.	Treatment.	Dressed Grain.		Straw per Acre.	Total Produce per Acre.
		Yield per Acre.	Weight per Bushel.		
		Bushels.	lbs.	cwts.	lbs.
1	Cresylic Acid	35.0	63.5	37.9	6745
2	Control	27.7	63.0	30.4	5387
3	Naphthaline	34.5	62.5	38.1	6645

NOTE.—Dressings on sawdust applied November 2nd, 1917, on Barley Stubble and ploughed in at once.
These plots were top dressed as farm, viz. 1 cwt. Sulph. Amm., 1½ cwt. Super. per acre.

MISCELLANEOUS EXPERIMENTS.

Barley. Hoos Field. Leguminous Strips, 1918, 1919, 1920.

Description of Plot.	Manuring per Acre.	1918.				1919.				1920.			
		Dressed Grain.		Straw per Acre.	Total Pr'duce per Acre.	Dressed Grain.		Straw per Acre.	Total Pr'duce per Acre.	Dressed Grain.		Straw per Acre.	Total Pr'duce per Acre.
		Bush. per Acre.	Weight per Bushel lbs.			Bush. per Acre.	Weight per Bushel lbs.			Bush. per Acre.	Weight per Bushel lbs.		
After Lucerne ...	Sulphate Amm. 1½ cwt. ...	20.1	55.3	10.7	2340	15.8	53.1	7.9	1790	27.3	52.8	20.5	3837
	S. Amm. 1½ cwt. Super. 3 cwt.	27.4	54.8	11.1	2777	18.4	54.1	8.0	1939	46.3	53.2	20.0	4799
	Sulphate Amm. 1½ cwt. ...	19.4	54.8	9.6	2170	12.5	53.3	6.9	1493	16.3	53.1	15.9	2719
After Red Clover	S. Amm. 1½ cwt.; Super. 3 cwt.	22.9	54.4	9.1	2282	16.1	54.6	6.6	1651	33.5	52.3	16.1	3630
	Sulphate Amm. 1½ cwt. ...	17.5	54.2	8.5	1930	10.6	53.6	6.7	1375	15.5	53.4	15.8	2657
After Alsike ...	S. Amm. 1½ cwt.; Super. 3 cwt.	21.4	53.8	8.7	2185	15.4	54.0	6.8	1621	38.0	52.1	18.6	4116

Leguminous crops ploughed in November, 1911. For crop yields, see previous Reports.
In 1915 the land was fallow; in 1916 and 1917, barley with clover: no separate weighings
were kept, however.

Mangolds. Long Hoos Field, 1919.

Plot.	Description of Plot.	Manuring per Acre.	Weight of Roots per Acre. Tons.
A	Bouted	20 tons of Dung, 3 cwt. Super., 1½ cwt. Sulphate Ammonia, applied on May 23 and a further 2 cwt. Sulphate Ammonia, applied July 28 ...	11.0
B	Flat		6.5
C	Bouted		9.0

Hoos Field. Barley sown with Clover, 1920.

(Formerly Barley after Alsike, p. 84). Clover cut with Barley and weighed

Plot.	Manures per Acre.	Clover. Yield per Acre. cwts.	Barley. Yield per Acre. cwts.	Total Produce per Acre. cwts.
1	8 cwt. Slag, 10 cwt. Lime	6.7	31.7	38.4
2	5 cwt. Super., 10 cwt. Lime, 14 tons Dung	15.2	31.2	46.4
3	10 cwt. Lime	4.9	27.7	32.6
4	5 cwt. Super., 1.5 cwt. Sulph. Potash, 10 cwt. Lime	9.8	28.1	37.9
5	5 cwt. Super., 10 cwt. Lime	7.1	28.6	35.7
6	10 cwt. Lime	4.5	30.4	34.8
7	10 cwt. Lime, 14 tons Farmyard Dung ...	8.5	41.5	50.0
8	8 cwt. Slag	5.4	21.9	27.3
9	5 cwt. Super., 14 tons Farmyard Dung ...	14.7	21.9	36.6
10	Control	3.6	21.9	25.5
11	5 cwt. Super., 1.5 cwt. Sulph. Potash ...	12.5	24.1	36.6
12	5 cwt. Super.	9.4	24.1	33.5
13	Control	6.3	22.3	28.6
14	14 tons Farmyard Dung	13.0	27.2	40.2
15	10 cwt. Lime, 14 tons Horse Manure ...	11.2	14.7	25.9
16	Control	4.9	14.3	19.2
17	14 tons Horse Manure	11.2	13.8	25.0
18	5 cwt. Super.	4.0	22.8	26.8
19	10 cwt. Lime, 14 tons Cattle Manure ...	8.5	29.5	38.0
20	Control	4.5	26.3	30.8
21	14 tons Cattle Manure	7.6	30.4	38.0

Manures sown March 13th, 1920. Horse, Cattle and Farmyard (Mixed) Manure put on Feb. 20th and 21st, 1920. Barley Seed sown March 19th, 1920. Clover Seed drilled between Barley rows, May 1st, 1920.

Wheat after Clover in 1917. Little Hoos Field, 1918.

Plot.	Manures per Acre.	Dressed Grain. Yield per Acre. lbs.	Straw per Acre. cwts.	Total Produce per Acre. lbs.
1	Control	2195	34.9	6323
6		2325	38.7	6937
3		2493	39.5	7190
8	Superphosphate 2 cwt.	2197	39.3	6905
2		2630	45.8	8058
5		2585	42.1	7655
4	Super. 2 cwt., Sulphate Amm. 1 cwt. ...	2823	44.2	8000
7		2400	44.0	7710