


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Report 1921-22 With the Supplement to the Guide to the Experimental Plots Containing the Yields per Acre Etc.



[Full Table of Content](#)

Table of Results - Later Experiments

Rothamsted Research

Rothamsted Research (1923) *Table of Results - Later Experiments* ; Report 1921-22 With The Supplement To The Guide To The Experimental Plots Containing The Yields Per Acre Etc., pp 90 - 104 - DOI: <https://doi.org/10.23637/ERADOC-1-110>

ROTATION PLOTS.

Little Hoos Field, 1921 and 1922.

Arranged to test the RESIDUAL VALUE of VARIOUS MANURES in year of application and one, two, and three years after.

Produce per acre.

Plot.	Manure per Acre from 1919 onwards.	Year of Dressing.	1921 (18th Season), Barley.					1922 (19th Season), Barley.				
			Dressed Grain.			Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.	Dressed Grain.			Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.
			Yield per Acre.	Weight per Bush.	Offal Grain per Acre.			Yield per Acre.	Weight per Bush.	Offal Grain per Acre.		
A 1	Control	—	Bush. 19.6	lb. 57.1	lb. 88	lb. 640	cwt. 10.0	Bush. 20.9	lb. 51.6	lb. 63	lb. 820	cwt. 10.6
2	...	1920	39.4	58.8	76	1512	19.9	[38.4]	52.9	94	1404	[19.2]
3	...	1921	37.7	58.9	57	1388	18.5	35.7	53.0	82	1284	16.5
4	Ordinary Dung, 16 tons	1922	25.9	58.0	67	844	13.5	38.2	52.7	90	1372	17.8
5	...	1915	31.2	57.4	85	1020	16.6	32.9	53.1	75	1232	15.3
B 1	Cake fed dung, 16 tons	1920	42.5	58.2	95	1804	21.3	38.1	52.7	84	1384	18.8
2	Control ...	—	18.8	56.6	71	688	10.4	26.9	52.8	76	1001	13.1
3	...	1921	40.4	58.6	66	1628	20.0	[39.7]	52.8	97	1484	[20.2]
4	Cake fed dung, 16 tons	1922	32.2	57.4	93	1236	16.3	[44.2]	53.2	109	1660	[22.5]
5	...	1915	30.9	57.8	72	1196	15.7	35.7	53.0	86	1364	17.3
C 1	Shoddy; Superphosphate; Sulphate of Potash ...	1920	22.0	56.0	118	940	12.1	16.6	51.9	95	696	9.9
2	...	1921	30.8	56.3	166	1276	16.0	19.2	52.5	79	816	11.2
3	Control ...	—	19.7	56.0	113	880	11.2	22.8	52.5	72	918	11.8
4	...	1922	21.1	56.0	117	1164	13.8	39.9	52.5	75	1448	18.4
5	Shoddy; Superphosphate; Sulphate of Potash ...	1919	23.2	56.4	102	1068	13.2	28.4	52.9	86	1132	14.5

D	1	Guano; Sulphate of Ammonia; Sulphate of	1920	28.4	56.0	118	1252	15.4	99.3	17.9	52.5	80	732	10.7	85.0
	2	Potash	1921	27.9	55.7	188	1348	17.1	90.8	21.0	52.6	71	924	12.1	86.9
	3	...	1922	18.7	56.3	112	888	11.6	89.9	35.6	53.0	113	1408	19.0	93.9
	4	Control	—	<i>14.7</i>	<i>54.5</i>	<i>122</i>	<i>772</i>	<i>9.9</i>	<i>83.5</i>	<i>17.9</i>	<i>51.8</i>	<i>87</i>	<i>788</i>	<i>12.2</i>	<i>74.5</i>
	5	Guano; Sulphate of Ammonia; Sulphate of Potash	1919	19.1	54.9	131	992	12.6	83.7	18.4	52.1	84	824	11.9	78.0
E	1	...	1920	25.5	55.8	111	1092	13.4	102.5	19.4	50.6	83	872	11.0	86.6
	2	Rape Dust; Superphosphate; Sulphate of Potash	1921	36.1	54.7	152	1524	17.3	109.8	25.5	53.6	70	1068	13.8	93.1
	3	...	1922	13.1	56.0	123	588	7.8	97.6	34.8	52.6	85	1328	17.0	100.8
	4	Control	1919	15.4	55.8	127	688	9.4	93.3	16.5	50.9	75	764	10.2	79.9
	5	...	—	<i>21.4</i>	<i>55.9</i>	<i>134</i>	<i>872</i>	<i>11.9</i>	<i>99.8</i>	<i>19.7</i>	<i>53.1</i>	<i>63</i>	<i>824</i>	<i>11.8</i>	<i>84.2</i>
F	1	Control	—	<i>10.3</i>	<i>52.3</i>	<i>84</i>	<i>476</i>	<i>7.1</i>	<i>78.1</i>	<i>20.3</i>	<i>52.3</i>	<i>87</i>	<i>1048</i>	<i>13.1</i>	<i>77.8</i>
	2	...	1920	22.1	55.5	94	896	11.9	98.5	23.6	52.6	73	960	12.1	96.9
	3	Superphosphate; Sulphate of Ammonia; Sulphate of Potash	1921	33.5	55.9	177	1480	17.8	102.9	23.4	53.8	68	992	12.8	92.5
	4	...	1922	17.6	53.0	115	808	10.5	89.5	37.8	52.5	97	1616	19.7	94.5
	5	...	1919	16.4	55.5	122	896	11.7	78.7	19.6	52.0	70	892	12.0	81.3
G	1	Bone Meal; Sulphate of Ammonia; Sulphate of	1920	20.4	53.7	115	844	11.6	93.5	22.6	53.7	96	1076	14.3	81.9
	2	Potash	1921	24.1	54.6	123	1208	14.1	90.8	28.0	52.8	76	1228	15.1	91.7
	3	Control	—	<i>22.9</i>	<i>54.9</i>	<i>93</i>	<i>1068</i>	<i>13.9</i>	<i>86.4</i>	<i>29.2</i>	<i>52.8</i>	<i>69</i>	<i>1292</i>	<i>15.3</i>	<i>93.9</i>
	4	Bone Meal; Sulphate of Ammonia; Sulphate of	1922	22.8	54.5	91	1004	12.6	94.1	36.2	53.8	103	1696	20.6	88.6
	5	Potash	1919	20.8	53.8	100	944	13.2	82.3	25.5	52.3	79	1208	14.6	86.3
H	1	...	1920	30.8	56.8	88	1184	14.8	111.2	28.1	52.7	87	1148	14.4	97.1
	2	Basic Slag; Sulphate of Ammonia; Sulphate of	1921	34.1	55.8	150	1672	20.1	91.0	29.5	53.2	82	1292	15.6	94.4
	3	Potash	1922	24.6	56.5	91	1040	13.9	94.9	40.3	53.2	100	1844	22.0	90.8
	4	...	1919	30.3	55.9	88	1236	15.2	104.5	30.4	53.8	64	1252	15.1	100.7
	5	Control	—	<i>27.8</i>	<i>55.4</i>	<i>106</i>	<i>1304</i>	<i>16.3</i>	<i>90.3</i>	<i>20.8</i>	<i>53.4</i>	<i>82</i>	<i>1020</i>	<i>12.5</i>	<i>85.6</i>

NOTES.—Since 1919 the manure for each plot (except of series A and B) has been rationed at 40 lb. Nitrogen, 100 lb. Calcium Phosphate, and 50 lb. Potash per acre. Each plot has been supplied with as much of its particular manure (shoddy, guano, &c.) as possible without exceeding the receipt in any of the three rationed ingredients.

Any deficit in either of these three has then been made good by adding the necessary quantity of Sulphate of Ammonia, Superphosphate, or Sulphate of Potash.

Figures in italics denote unmanured plots. The yields on the plots to which the manure was applied in a given year are printed in heavy type.

Figures in square brackets are estimated yields.

STRAW EXPERIMENT, 1921. Potatoes (Arran Chief). Sawpit Field.

Manure per Acre.	Yield per Acre.		
	1st Plot	2nd Plot	3rd Plot
8 tons Rotted Straw Manure—Single Nitrogen ...	Tons 2'30	Tons 2'18	Tons 1'96
16 " " " " " " " ...	2'48	2'63	2'16
32 " " " " " " " ...	1'73	2'39	2'29
2 cwt. Sulphate of Ammonia	1'20	1'48	1'13
4 " " " " " " " ...	1'66	1'57	1'48
8 " " " " " " " ...	1'52	1'71	1'38
16 " " " " " " " ...	1'41	1'55	1'27
8 tons Rotted Straw Manure—Double Nitrogen ...	2'09	2'20	1'86
16 " " " " " " " ...	3'32	2'59	2'50
32 " " " " " " " ...	2'16	2'68	2'04
Control—No Manure	1'39	1'61	1'41
" " " " " " " ...	1'52	1'45	1'39

Single Nitrogen represents 1 cwt. Sulphate of Ammonia added to 1 ton of straw.

Double Nitrogen represents 2 cwt. Sulphate of Ammonia added to 1 ton of straw.

RESIDUAL VALUE OF SLUDGE, 1921. Long Hoos Field.

Treatment of Plots in 1920. Manure per Acre.	Dressed Grain.				Offal Grain per Acre.		Straw per Acre.				Proportion of Total Grain to 100 of Total Straw.	
	Yield per Acre. Bush.		Weight per Bushel. lb.		lb.		Straw. lb.		Total Straw. cwt.			
	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.
1921, Wheat (Red Standard) after Potatoes (1920).*												
Activated Sewage Sludge, 13'3 tons	29'8	27'9	64'0	64'1	371	406	2925	2624	32'7	30'8	62'2	63'6
Farmyard Dung 15 tons ...	34'8	31'6	64'0	64'1	296	371	2461	2600	30'3	29'7	74'5	72'0
Control	26'0	26'9	63'3	63'0	342	325	2299	1997	26'2	26'6	67'6	67'7
1921, Wheat (Red Standard) after Barley (1920).†												
Sulph. of Ammonia 1'45 cwt. ...	24'1		63'0		387		2738		31'1		54'6	
Activated Sewage Sludge, 2'7 tons	30'1		63'0		351		2857		31'4		64'1	
Control	27'2		62'5		405		2738		29'4		63'9	
Control	27'4		63'0		435		2333		30'3		63'7	

* In 1920 this set received a basal dressing of 6 cwt. Super. and 1 cwt. Nitrate of Ammonia per acre. No manure was given in 1921.

† In 1921 this set was manured as farm, viz., 1 cwt. Sulphate of Ammonia and 1 cwt. Superphosphate per acre.

TOP DRESSING EXPERIMENTS.

Treatment of Plots and Quantities per Acre.	Dressed Grain						Official Grain per Acre.			Straw per Acre.						Proportion of Total Grain to 100 of Total Straw.		
	Yield per Acre Bushels.			Weight per Bushel lb.			lb.			lb.			cwt.					
	Plot			Plot			Plot			Plot			Plot			Plot		
	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
Oats (Grey Winter).*																		
Great Knott Field, 1921.																		
Super. 1½ cwt.	52.1	57.4	54.7	44.1	43.0	43.6	420	490	460	1900	2600	2540	31.8	37.5	40.2	76	71	63
Super. 1½ cwt.; Sul./Amm. 1½ cwt., applied March 4	45.0	43.6	44.2	43.3	45.0	43.0	380	485	380	2020	1920	2620	31.6	35.5	37.9	66	61	54
Super. 1½ cwt.; Mur./Amm. 144 lb., applied March 4	48.4	57.3	44.0	44.0	44.0	44.6	435	590	485	2280	2980	2480	37.1	47.0	36.4	62	59	60
No Manure	47.1	48.6	47.9	43.6	44.0	43.5	410	475	420	2060	2340	2200	31.1	36.1	33.4	71	65	67
Wheat (Red Standard).																		
Great Harpenden Field, 1921.																		
Super. 200 lb.; Sul./Amm. 100 lb., applied March 2-3	19.4	16.9	—	62.5	62.5	—	565	430	—	2350	1660	—	30.0	22.9	—	53	58	—
Super. 200 lb.; Sul./Amm. 200 lb., applied March 2-3	20.3	15.4	—	62.5	62.5	—	540	443	—	2060	1900	—	27.1	25.6	—	60	49	—
Super. 200 lb.; Sul./Amm. 100 lb., applied March 2-3 and Sul./Amm. 100 lb., applied May 2	21.3	13.8	—	62.5	62.5	—	495	350	—	2080	1450	—	27.2	20.7	—	60	52	—
Super. 200 lb.; Mur./Amm. 86 lb., applied March 2-3	17.7	15.7	—	62.0	62.0	—	430	370	—	1960	1700	—	24.4	22.8	—	56	53	—
Super. 200 lb.; N./Soda 140 lb., applied March 2-3	15.2	16.9	—	62.5	62.0	—	355	415	—	1750	1980	—	21.4	24.0	—	54	55	—
Super. 200 lb.; Sul./Amm. 100 lb., applied May 2	23.1	16.1	—	62.3	62.0	—	500	385	—	2660	1800	—	30.4	21.9	—	57	56	—
Super. 200 lb.; Sul./Amm. 200 lb., applied May 2	23.8	17.4	—	62.5	62.5	—	500	485	—	2620	1900	—	30.6	22.6	—	58	62	—
Super. 200 lb.; Mur./Amm. 86 lb., applied May 2	17.5	18.2	23.6	62.0	62.0	63.0	338	405	465	1700	1710	2500	21.3	22.6	27.8	60	61	63
Super. 200 lb.; Sul./Amm. 100 lb., applied April 4	17.1	—	—	62.5	—	—	365	—	—	1780	—	—	24.0	—	—	53	—	—
No Manure	13.5	17.6	—	63.5	61.5	—	375	368	—	1320	1880	—	17.6	22.2	—	63	58	—
Super. 200 lb.	19.0	12.2	22.0	62.5	62.5	62.5	415	385	525	1830	1260	2540	22.7	17.1	28.1	63	60	60
Super. 200 lb.	—	16.6	—	—	61.0	—	—	360	—	—	1840	—	—	22.6	—	—	54	—
Wheat (Red Standard).†																		
Foster's Field, 1922.																		
1 cwt. Sul./Amm. applied March 18	14.4	18.4	18.5	59.5	60.0	60.0	425	391	400	1425	1650	1650	17.0	19.2	19.9	67	70	68
1 cwt. Sul./Amm., applied April 20	18.3	21.3	15.8	59.5	60.3	60.0	394	394	353	1750	1850	1475	21.2	23.0	17.6	62	65	66
1 cwt. Sul./Amm., applied May 18	17.2	21.9	16.4	60.0	60.5	60.3	372	409	406	1625	1750	1425	19.4	21.4	17.9	65	72	70
93 lb. Mur./Amm., applied March 18	17.8	19.6	16.4	60.0	60.3	60.5	362	400	300	1550	1875	1375	19.2	22.8	18.0	66	62	64
186 lb. Mur./Amm., applied March 18	19.7	21.2	20.2	59.5	60.0	60.3	350	409	362	1700	1650	1800	21.0	20.8	21.7	65	72	65
2 cwt. Sul./Amm., applied March 18	20.0	20.8	18.3	60.3	60.0	60.0	438	503	369	1800	1950	1725	22.5	25.2	22.1	65	62	59
2 cwt. Sul./Amm., applied April 20	20.1	18.4	13.2	60.8	60.0	55.5	394	525	369	1700	1775	1475	21.4	24.3	19.0	67	60	52
2 cwt. Sul./Amm., applied May 18	19.4	17.9	17.2	60.0	60.5	60.0	425	506	441	1700	1675	1525	21.2	23.0	20.1	67	62	65
Basal Manuring only	13.4	15.0	11.9	59.8	60.0	60.0	328	391	297	1250	1325	1225	15.2	17.4	14.7	67	66	61
No Manure	12.9	16.7	13.2	60.0	60.3	59.5	378	328	331	1325	1525	1350	15.6	18.5	16.5	66	64	60
Barley (Plumage Archer).																		
Long Hoos, 1922.																		
2 cwt. Super.; 1 cwt. Sul./Amm., applied May 9	34.5	30.1	—	51.3	51.8	—	281	238	—	1650	1325	—	22.5	18.0	—	81	89	—
2 cwt. Super.; 2 cwt. Sul./Amm., applied May 9	31.2	35.3	—	51.8	51.6	—	369	287	—	1600	1775	—	21.4	23.0	—	83	82	—
2 cwt. Super.; 98 lb. Mur./Amm., applied May 9	27.7	30.4	—	51.0	51.3	—	256	263	—	1475	1500	—	20.5	21.4	—	73	76	—
2 cwt. Super.; 1 cwt. Mur./Amm., applied May 9	25.7	32.3	—	52.0	54.3	—	250	350	—	1175	1575	—	18.5	23.0	—	77	82	—
2 cwt. Super.; 51½ lb. Urea, applied May 9	32.1	34.7	—	51.8	52.3	—	287	356	—	1575	1675	—	21.0	22.5	—	83	86	—
2 cwt. Super.	23.8	26.6	—	51.0	52.3	—	184	200	—	1175	1250	—	16.3	17.2	—	77	83	—
No Manure	23.6	31.2	—	52.0	51.3	—	200	244	—	1100	1650	—	16.5	22.1	—	77	75	—

* A luxuriant crop of mustard was ploughed in previous to the sowing of the Oats; hence the high yields on these plots.
† All plots (except the unmanured), received a basal dressing of 2 cwt. Super.; 1 cwt. S./Pot. per Acre.

Top Dressing Experiments—*contd.*

Root Crops. Great Harpenden Field, 1922.

Manuring per Acre.	Yield per Acre.	
	1st Plot. Tons.	2nd Plot. Tons.
Potatoes (Kerr's Pink).		
Dunged Series: 15 tons Farmyard Dung per Acre—		
Super. 4 cwt., Sul./Pot. 1½ cwt.	6·73	5·41
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 3 cwt. (half as Top Dressing)	7·92	9·17
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 1½ cwt. ...	7·91	8·06
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul. Amm. 4½ cwt.		
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 3 cwt. ...	10·54	9·62
(1½ cwt. as Top Dressing)	10·08	9·37
Super. 4 cwt., Sul./Pot. 1½ cwt., Mur./Amm. 290 lb.	10·66	10·74
Undunged Series:		
Super. 6 cwt., Sul./Pot. 2 cwt.	6·10	4·90
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 3 cwt. (half as Top Dressing)	7·99	7·89
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 1½ cwt. ...	6·98	7·75
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 4½ cwt. (1½ cwt. as Top Dressing)	9·60	8·36
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 3 cwt. ...	8·72	9·22
Super. 6 cwt., Sul./Pot. 2 cwt., Mur./Amm. 290 lb. ...	9·21	8·50
Swedes (Hurst's Monarch).		
589 lb. Slag,* 1 cwt. Sul./Pot.	{ R 25·13 L 3·04	{ 28·24 4·29
589 lb. Slag,* 1 cwt. Sul./Pot., 2 cwt. Sul./Amm. (as Top Dressing)	{ R 27·48 L 3·82	{ 30·65 4·87
589 lb. Slag,* 1 cwt. Sul./Pot., 10 tons Farmyard Dung	{ R 28·75 L 4·22	{ 32·37 4·12
589 lb. Slag,* 1 cwt. Sul./Pot., 10 tons Farmyard Dung, 2 cwt. Sul./Amm. (as Top Dressing)	{ R 32·61 L 4·60	{ 32·43 4·71

* Equivalent to 5 cwt. Super.

R = Roots.

L = Leaves.

Slag Experiments—contd.
Clover. West Barnfield, 1921 and 1922.

No. of Plot.	Treatment of Plots and Quantities per Acre.	1921				1922			
		Yield per Acre.		Dry Matter per Acre.		Yield per Acre.		Dry Matter per Acre.	
		Series A	Series B	Series A	Series B	Series A	Series B	Series A	Series B
1	High Grade Slag No. 12, 1170 lb.	cwt. 40.8	cwt. 40.4	lb. 3521	lb. 3567	cwt. 10.5	cwt. 16.1	lb. 941	lb. 1418
2	Open Hearth, High Soluble Slag No. 13, 1925 lb.	43.0	38.7	3629	3470	18.5	13.7	1644	1183
3	Open Hearth, Low Soluble Slag No. 14, 1930 lb.	42.4	40.4	3720	3567	18.6	15.6	1679	1374
4	Gafsa Phosphate, 750 lb.	41.7	39.4	3654	3502	17.6	18.3	1604	1681
C	No Manure	43.6	39.4	3812	3593	16.3	18.3	1486	1630
C	"	40.8		3563		17.1		1490	

Barley (Plumage Archer). Long Hoos Field, 1922.

Treatment of Plots.	Dressed Grain.				Offal Grain per Acre.				Straw per Acre.				Proportion of Total Grain to 100 of Total Straw.			
	Yield per Acre in Bushels.		Weight per Bushel in lb.		lb.		lb.		lb.		lb.		lb.		lb.	
	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.
	No. 20.	No. 2.	No. 1.	No. 20.	No. 2.	No. 1.	No. 20.	No. 2.	No. 1.	No. 20.	No. 2.	No. 1.	No. 20.	No. 2.	No. 1.	No. 20.
Basal Manuring, Slag, full quantity	36.0	26.0	28.7	51.3	51.4	51.8	197	162	231	1375	1175	1250	18.5	16.9	19.3	99
Basal Manuring, Slag, half quantity	35.1	31.7	25.5	52.8	52.0	51.0	200	172	213	1375	1238	1238	19.3	18.0	19.1	95
Gafsa Phosphate, 87 lb.	29.9	25.5	32.5	51.9	51.8	51.5	194	169	241	1238	1113	1438	18.5	16.9	20.3	84
	26.2	36.4	29.2	51.3	51.4	51.3	163	181	200	*875	1500	1275	*11.7	19.6	18.3	*115
Basal Manuring, Gafsa Phosphate, 174 lb.	34.8	25.2	27.5	51.6	52.3	52.0	231	162	209	1488	1088	1213	19.6	16.5	18.2	92
Basal Manuring only	25.7	33.7	34.1	52.0	52.0	51.5	178	178	203	1063	1363	1388	15.8	18.3	18.9	85
No Manure	30.1	24.7	27.0	51.8	50.8	51.1	188	203	228	1363	1100	1263	19.2	15.9	18.8	81

Basal Manuring is 1 cwt. Sulphate of Potash; 1 cwt. Sulphate of Ammonia and 436 lb. Slag No. 1 per acre. Full Quantity Slag represents 636 lb. Slag No. 20, 602 lb. Slag No. 2

*There was a high wind blowing when this plot was threshed, hence the low figure for the yield of straw.

Slag Experiments—*contd.*

Swedes (Hurst's Monarch) Produce per Acre.
Great Harpenden Field, 1922.

Manuring per Acre.	Roots.			Leaves.		
	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 2.	Slag No. 1.
	Tons.	Tons.	Tons.	Tons.	Tons.	Tons.
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag full quantity ...	25·92	27·92	30·40	4·89	3·82	4·16
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag full quantity ...	32·08	30·31	30·40	4·01	5·04	4·20
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag half quantity, Gafsa Phosphate, 175 lb. ...	27·19	28·04	31·88	4·18	3·53	4·10
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag half quantity, Gafsa Phosphate, 175 lb. ...	28·21	29·78	28·82	4·28	4·16	4·27
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., No. 7 Nauru Phos- phate, 262½ lb. ...	30·96	26·43	26·50	4·49	4·00	3·98
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., No. 3 Gafsa Phos- phate, 350 lb. ...	27·83	31·12	28·46	3·95	4·58	4·66
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt. ...	27·21	31·45	25·74	4·16	5·02	3·99
No Manure ...	25·67	27·23	22·70	3·54	3·67	3·19

NOTE.—"Full Quantity" Slag is No. 20, 1275 lb. per Acre.

No. 2, 1225 ,,

No. 1, 875 ,,

Description of Slags Used.

No.	Type.	Total Phosphate as $\text{Ca}_3(\text{PO}_4)_2$	Solubility %
1	Open Hearth, L.G., H.S. ...	25·0	90·4
2	" " L.G., L.S. ...	18·0	35·7
8	Phosphate, Slag Mixture ...	53·1	25·5
12	Talbot Process, H.G., H.S. ...	37·3	80·7
13	Open Hearth, L.G., H.S. ...	22·7	91·5
14	" " L.G., L.S. ...	22·6	29·0
15	Talbot Process, H.G., H.S. ...	40·0	72·5
16	Open Hearth, L.G., H.S. ...	21·3	88·3
17	Bessemer, H.G., H.S. ...	42·5	77·2
18	Open Hearth, L.G., H.S. ...	20·8	67·0
19	" " L.G., L.S. ...	20·2	21·0
20	" " L.G., H.S. ...	17·2	78·8

L.G. = Low Grade. L.S. = Low Soluble.

H.G. = High Grade. H.S. = High Soluble.

POTASH EXPERIMENTS.

Manuring per Acre.	Dry Matter per Acre.			Yield per acre.		
	1st Plot	2nd Plot	3rd Plot	1st Plot	2nd Plot	3rd Plot
Clover. West Barn Field, 1922.						
Control	lb. 1369	lb. 1273	lb. 1507	cwt. 15'2	cwt. 15'7	cwt. 18'6
Sulphate of Potash, 210 lb.	1533	1929	2123	18'6	25'0	26'4
Cement Works' Dust, 511 lb.	1381	1710	1729	17'5	21'8	21'4

Potatoes (Arran Chief). Sawpit Field, 1921.

With Dung, 12 tons per Acre.

	Tons.	Tons.	Tons.
3 cwt. Super., 1½ cwt. Sulphate Ammonia, 470 lb. Sylvenite ...	3'57	*3'15	3'71
3 cwt. Super., 1½ cwt. Sulphate Ammonia, ...	3'55	*3'18	3'72
3 cwt. Super., 1½ cwt. Sulphate Ammonia, 1½ cwt. Sulphate Potash	3'67	4'27	3'88
3 cwt. Super., 1½ cwt. Sulphate Ammonia, 1½ cwt. Sulphate Potash, 95 lb. Sulphate Magnesium	*3'07	3'92	3'87
No Manure. Control	*2'28	3'48	3'18
3 cwt. Super., 1½ cwt. Sul. Amm., 1½ cwt. Muriate Potash	*2'31	4'24	3'97
3 cwt. Super., 1½ cwt. Sul. Amm., 1½ cwt. Muriate Potash, 84 lb. Sul. Magnesium	*2'43	3'90	4'15

Without Dung.

4 cwt. Super., 2 cwt. Sul. Amm., 625 lb. Sylvenite	3'49	4'04	3'11
4 cwt. Super., 2 cwt. Sul. Amm.	1'43	1'48	1'15
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Sul. Potash	3'48	4'28	3'52
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Sul. Pot., 127 lb. Sul. Mag.	3'85	4'26	3'25
No Manure. Control	1'24	1'72	1'65
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Muriate Potash	4'15	4'20	4'00
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Muriate Potash, 111 lb. Sul. Magnesium	4'27	3'95	3'63

Potatoes (Arran Chief). Sawpit Field, 1921.

4 cwt. Super., 2 cwt. Sulphate Ammonia, 232 lb. Sul. Potash ...	3'00	2'46	2'82
4 cwt. Super., 2 cwt. Sulphate Ammonia	1'16	0'98	0'89
4 cwt. Super., 2 cwt. Sulphate Ammonia, 5'4 cwt. Sylvenite ...	*1'93	3'36	3'04
Control. No Manure.	*0'73	1'10	1'16

Potatoes (Kerr's Pink). Great Harpenden Field, 1922.

With Dung 15 tons per Acre.

Basal Manuring (=Super. 4 cwt., Sul. Amm. 1'5 cwt. per Acre) ...	8'78	7'72	7'60
Sulphate Potash 183 lb. + Basal Manuring	9'49	9'72	9'45
Muriate Potash 148 lb. + Basal Manuring	9'22	9'60	8'82
Muriate Potash 148 lb. + Salt 497 lb. + Basal Manuring	9'84	9'49	9'14

Without Dung.

Basal (=Super. 6 cwt., Sulphate Ammonia 2 cwt. per Acre) ...	2'11	2'75	2'57
Sulphate Potash 244 lb. + Basal	7'88	8'96	8'06
Muriate Potash 197 lb. + Basal	8'62	8'73	7'62
Muriate Potash 197 lb. + Salt 662 lb. + Basal	8'45	8'27	8'43
Muriate Potash 197 lb. Sulphate Magnesium, 344 lb. + Basal ...	8'68	8'90	7'62
Muriate Potash 197 lb. Salt 662 lb. + Basal	8'66	8'02	7'51
No Manure	3'23	2'87	2'83
Sulphate Potash 244 lb. Sulphate Magnesium 344 lb. + Basal ...	9'25	8'79	7'11
Cement Works' Dust 614 lb. + Basal	7'47	6'66	6'38
Sylvenite 541 lb. + Basal	8'38	7'92	6'90

* On these plots the bouts were badly broken down due to extra hoeing on account of the growth of Wheatbind.

Mangolds (Prizewinner Yellow Globe).

Great Harpenden Field, 1922.

Produce per Acre.

Manuring per Acre.	Roots.		Leaves.	
	1st Plot Tons.	2nd Plot Tons.	1st Plot Tons.	2nd Plot Tons.
No. 9 Slag 4 cwt., Sulphate Ammonia 2 cwt., Sulphate Potash 2 cwt.	17'64	14'12	5'57	5'13
No. 9 Slag 4 cwt., Sulphate Ammonia 2 cwt.	10'45	11'61	4'73	4'94
No. 9 Slag 4 cwt., Sulphate Ammonia 2 cwt., Cement Works' Dust	18'75	18'25	5'61	5'96
No Manure	10'88		4'25	

POTATOES.

Relative Effects of Sulphates and Chlorides on different varieties.

Great Harpenden Field, 1922.

Variety.	Dunged Series.						Undunged Series.					
	Actual Weight of Potatoes.			Average Weight per Plant.			Actual Weight of Potatoes.			Average Weight per Plant.		
	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.
	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.
Ajax ...	16	12 $\frac{3}{4}$	19 $\frac{3}{4}$	3'20	2'55	2'82	13 $\frac{1}{2}$	17 $\frac{3}{4}$	4	2'25	2'54	1'00
	24	21 $\frac{1}{4}$	12 $\frac{1}{4}$	4'00	3'04	1'75	17 $\frac{1}{4}$	18	4 $\frac{1}{2}$	2'46	3'00	0'64
	27	16 $\frac{3}{4}$	33	3'86	4'13	4'71	7 $\frac{1}{4}$	23	4	2'42	3'29	0'67
Arran Comrade	15 $\frac{3}{4}$	11 $\frac{3}{4}$	7 $\frac{1}{4}$	2'25	1'96	2'42	11 $\frac{1}{4}$	8 $\frac{3}{4}$	3 $\frac{1}{4}$	2'25	1'46	0'65
	10 $\frac{1}{4}$	10 $\frac{3}{4}$	13	2'56	2'15	2'17	11 $\frac{1}{4}$	11 $\frac{1}{4}$	1 $\frac{1}{2}$	1'88	2'25	0'30
	15 $\frac{1}{2}$	10 $\frac{1}{2}$	13	2'58	2'10	2'17	15	18	1 $\frac{3}{4}$	2'14	2'57	0'29
British Queen	19 $\frac{1}{4}$	19	19 $\frac{1}{4}$	3'21	2'71	2'75	10 $\frac{1}{4}$	13 $\frac{3}{4}$	7 $\frac{3}{4}$	1'46	1'96	1'11
	19 $\frac{3}{4}$	18 $\frac{3}{4}$	19 $\frac{1}{4}$	2'82	2'68	2'75	16 $\frac{3}{4}$	11 $\frac{1}{2}$	8 $\frac{1}{4}$	2'36	1'92	1'18
	26 $\frac{3}{4}$	25	23 $\frac{1}{4}$	3'82	4'17	3'32	11 $\frac{3}{4}$	15 $\frac{3}{4}$	3 $\frac{3}{4}$	1'96	2'63	0'75
Duke of York ...	7 $\frac{3}{4}$	11	11 $\frac{1}{4}$	1'11	1'57	1'61	9	8	1	1'80	1'60	0'33
	8 $\frac{3}{4}$	14	14	1'25	2'00	2'00	9 $\frac{1}{4}$	6 $\frac{1}{2}$	2 $\frac{1}{2}$	1'54	0'93	0'63
	13 $\frac{1}{2}$	10 $\frac{1}{2}$	14 $\frac{3}{4}$	2'25	1'75	2'46	6 $\frac{1}{4}$	10 $\frac{1}{4}$	1 $\frac{1}{4}$	1'04	1'46	0'42
Epicure ...	16 $\frac{1}{2}$	14 $\frac{3}{4}$	10	2'36	2'11	1'43	12 $\frac{1}{4}$	9 $\frac{3}{4}$	1 $\frac{3}{4}$	2'04	1'63	0'35
	11 $\frac{1}{2}$	13 $\frac{1}{2}$	13 $\frac{1}{2}$	1'64	1'93	2'25	13 $\frac{1}{2}$	13 $\frac{3}{4}$	3 $\frac{3}{4}$	1'93	1'96	0'54
	16	18 $\frac{1}{2}$	19 $\frac{1}{2}$	2'29	2'64	2'79	11 $\frac{1}{4}$	12 $\frac{1}{4}$	1	1'68	1'79	0'25
Great Scott ...	13 $\frac{1}{2}$	19 $\frac{1}{2}$	21 $\frac{1}{2}$	3'38	2'79	3'07	21 $\frac{1}{2}$	17 $\frac{1}{4}$	4 $\frac{1}{2}$	3'07	2'46	0'75
	21 $\frac{1}{2}$	24 $\frac{1}{4}$	19 $\frac{1}{2}$	3'07	3'54	3'25	11 $\frac{3}{4}$	12 $\frac{3}{4}$	1 $\frac{1}{4}$	1'96	2'13	0'42
	27 $\frac{1}{4}$	29	24 $\frac{1}{2}$	3'89	4'14	3'50	14 $\frac{1}{4}$	13 $\frac{1}{4}$	1	2'38	2'65	0'50
Iron Duke ...	24	20	21	3'43	3'33	3'50	16 $\frac{3}{4}$	19 $\frac{1}{4}$	4 $\frac{3}{4}$	2'79	2'75	0'68
	21	18 $\frac{1}{2}$	16 $\frac{1}{4}$	3'00	3'08	2'32	10 $\frac{3}{4}$	20 $\frac{1}{4}$	4	1'79	2'89	1'00
	23 $\frac{3}{4}$	23 $\frac{1}{4}$	23	3'96	3'32	3'29	20	13	4 $\frac{1}{2}$	2'86	3'25	0'64
K. of K. ...	26	23 $\frac{3}{4}$	20 $\frac{1}{4}$	3'71	3'39	2'89	21 $\frac{1}{2}$	18 $\frac{1}{2}$	7 $\frac{1}{4}$	3'07	3'08	1'04
	28 $\frac{1}{2}$	27 $\frac{3}{4}$	21	4'07	4'63	4'20	—	—	—	—	—	—
	29 $\frac{1}{2}$	29 $\frac{1}{2}$	30 $\frac{1}{4}$	4'21	4'21	4'32	19 $\frac{1}{4}$	15 $\frac{1}{2}$	5 $\frac{3}{4}$	2'75	3'10	0'82
Kerr's Pink ...	18 $\frac{1}{4}$	20 $\frac{3}{4}$	12	3'04	2'96	2'00	18 $\frac{3}{4}$	20 $\frac{3}{4}$	6 $\frac{1}{2}$	2'68	2'96	0'93
	25	22 $\frac{1}{4}$	15	3'57	3'18	3'00	11 $\frac{1}{2}$	19 $\frac{1}{2}$	3 $\frac{1}{4}$	1'92	3'90	0'46
	26 $\frac{3}{4}$	30 $\frac{1}{4}$	15 $\frac{1}{2}$	3'82	4'32	3'88	24 $\frac{1}{4}$	22	5	3'46	3'14	0'71
Nithsdale ...	18	14 $\frac{1}{4}$	30 $\frac{3}{4}$	2'57	2'04	1'96	9	9 $\frac{1}{2}$	1 $\frac{1}{4}$	1'29	1'58	0'42
	15 $\frac{1}{2}$	20 $\frac{1}{2}$	20	2'21	2'93	2'86	12	15	1	2'00	2'14	0'33
	21 $\frac{1}{2}$	26	14 $\frac{1}{4}$	3'58	3'71	3'56	14 $\frac{1}{4}$	14 $\frac{1}{2}$	2 $\frac{1}{2}$	2'04	2'07	0'63
Tin Perfection	20 $\frac{3}{4}$	17	12 $\frac{3}{4}$	3'46	2'83	2'55	20	19 $\frac{1}{2}$	7 $\frac{1}{2}$	2'86	2'79	1'07
	21 $\frac{3}{4}$	20 $\frac{3}{4}$	23 $\frac{3}{4}$	3'11	2'96	3'39	18 $\frac{3}{4}$	17 $\frac{3}{4}$	8 $\frac{3}{4}$	2'68	2'54	1'21
	17 $\frac{1}{2}$	19 $\frac{1}{4}$	23 $\frac{1}{2}$	2'50	3'21	3'36	21 $\frac{3}{4}$	17	7	3'11	2'83	1'00
Up-to-Date ...	25 $\frac{3}{4}$	23 $\frac{3}{4}$	25 $\frac{1}{4}$	4'29	3'39	4'21	26 $\frac{3}{4}$	20 $\frac{1}{4}$	9 $\frac{1}{4}$	3'82	2'89	1'32
	20 $\frac{1}{2}$	25 $\frac{1}{2}$	25 $\frac{1}{2}$	2'93	3'68	3'64	20 $\frac{1}{2}$	14 $\frac{1}{4}$	8 $\frac{1}{4}$	2'93	2'38	1'18
	29 $\frac{3}{4}$	28 $\frac{3}{4}$	28 $\frac{3}{4}$	4'25	4'07	4'11	21 $\frac{3}{4}$	21	11	3'11	3'00	1'83

NOTE.—7 Plants were set in each Row.

Manures were:—Dunged Series: Basal Row: Super. 4 cwt.; Sulphate of Ammonia 1 $\frac{1}{2}$ cwt.;
Dung 15 tons per Acre.Sulphate Row: Basal Manuring; Sulphate of Potash 184 lb.
per Acre.Chloride Row: Basal Manuring; Muriate of Potash 147 lb.
per Acre.Undunged Series: Basal Row: Super. 6 cwt.; Sulphate of Ammonia 2 cwt.
per Acre.Sulphate Row: Basal Manuring; Sulphate of Potash 244 lb.
per Acre.Chloride Row: Basal Manuring; Muriate of Potash 197 lb.
per Acre.

Potatoes. Great Harpenden Field, 1922.
Comparison of Varieties.

	Ajax.	Arran Comrade.	British Queen.	Duke of York.	Epicure.	Great Scott.	Iron Duke.	K. of K.	Kerr's Pink.	Nithsdale.	Tin Perfection.	Up-to-Date
Average weight of Potatoes lifted per row	16'21	10'54	16'08	8'86	11'88	16'58	16'89	21'62	17'63	13'49	17'49	21'47
Average weight per plant ...	2'70	1'90	2'43	1'50	1'81	2'79	2'67	3'28	2'73	2'23	2'62	3'17

Comparison of Manurial Treatment.

	Dunged Series.			Undunged Series.		
	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.
Average weight of Potatoes lifted per row	20'12	19'82	18'62	15'17	15'41	4'40
Average weight per plant ...	3'08	3'03	2'94	2'36	2'37	0'80

PROFESSOR BLACKMAN'S
ELECTRO CULTURE EXPERIMENTS.

Clover. Great Knott Field, 1921.

Plots.									Yield per Acre
Electro-Culture	...	...	...	...	...	...	...	...	cwt. 42'0
Control	...	...	...	...	...	...	...	...	41'2

Cereal Crops.

Plots.	Dressed Grain.		Ofal Grain per Acre.	Straw per Acre.		Proportion of Total Grain to 100 of Total Straw.		
	Yield per Acre.	Weight per Bush.		Straw.	Total Straw.			
	Bushels.	lb.	lb.	lb.	cwt.			
Oats (Grey Winter). Foster's Field, 1921.								
Electro-Culture	...	...	40·7	43·4	241	1543	19·3	93·0
Control I.	...	...	33·1	42·0	298	1220	14·9	101·4
Control II.	...	...	31·6	42·2	234	1102	14·6	96·0
Wheat (Red Standard). Foster's Field, 1922.								
Electro-Culture	...	...	15·4	61·4	234	1229	15·8	66·9
Control, North East	...	...	16·5	60·6	249	1272	15·5	72·1
Control, South East	...	...	17·2	61·8	231	1196	14·2	81·5
Barley (Plumage Archer). Great Knott Field, 1922.								
Electro-Culture	...	...	34·1	49·1	273	1808	22·2	78·2
Control	...	...	32·4	48·6	244	1840	22·3	72·8

BORON EXPERIMENT

Barley (Plumage Archer). Little Hoos, 1922.

Treatment of Plots.	Dressed Grain.						Offal Grain per Acre. lb.			Straw per Acre.						Proportion of Total Grain to 100 of Total Straw.		
	Yield per Acre. Bushels.			Weight per Bushel. lb.						Straw. lb.			Total Straw. cwt.					
	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3
Boric Acid 20 lb. per acre ...	37.9	40.8	30.8	51.1	51.8	52.0	191	138	84	2025	1875	1850	24.6	23.2	22.1	77.4	86.5	68.2
Boric Acid 8 lb. per acre ...	36.5	40.0	41.3	51.5	52.0	52.0	169	113	150	1825	1800	1850	23.4	22.8	23.0	78.1	86.0	89.2
Control ...	34.9	40.8	38.6	50.9	52.4	52.5	156	134	119	1725	1775	1850	21.4	22.5	23.4	80.5	89.9	81.7

All plots received a basal dressing of Superphosphate 3 cwt.; Sulphate of Potash 1 cwt.; Sulphate of Ammonia $1\frac{1}{2}$ cwt.

EXPERIMENTS WITH NITROGENOUS MANURES

Potatoes (Arran Chief). Sawpit Field, 1921.

Manure per Acre.	Yield per Acre.		
	1st Plot. Tons.	2nd Plot. Tons.	3rd Plot. Tons.
4 cwt. Super., 1 cwt. Sulphate Potash, 2 cwt. Sulphate Ammonia	2.27	2.24	2.43
4 cwt. Super., 1 cwt. Sulphate Potash	1.84	2.13	1.99
4 cwt. Super., 1 cwt. Sulphate Potash, 193 lb. Muriate Ammonia	2.18	2.67	2.61
Control	1.33	1.41	1.49
4 cwt. Super., 1 cwt. Sulphate Potash, 102 lb. Urea	*1.72	2.69	2.57

* The bouts on this plot were badly broken down due to extra hoeing on account of growth of Wheatbind.

Barley (Plumage Archer). Stackyard Field, 1921.

Manures per Acre.	Dressed Grain.						Offal Grain per Acre.			Straw per Acre.						Proportion of Total Grain to 100 of Total Straw.		
	Yield per Acre. Bushels.			Weight per Bushel. lb.						Straw. lb.			Total Straw. cwt.					
	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3
1½ cwt. Super., 145lb. M./Amm.	40.4	34.8	—	54.7	55.5	—	197	135	—	2000	2000	—	23.5	25.9	—	91	71	—
1½ cwt. Super....	27.2	27.1	24.1	56.0	55.5	55.0	153	103	109	1325	1475	1350	17.1	18.4	17.2	88	78	75
1½ cwt. Super., 1½cwt. S./Amm.	38.3	36.5	30.2	55.7	55.2	54.2	144	175	194	1900	2050	1825	23.6	24.9	22.1	87	79	74
1½ cwt. Super., 76¾ lb. Urea ...	38.2	34.6	29.2	55.0	54.5	54.2	150	150	169	2000	2025	1775	24.2	24.7	21.3	83	74	73
No Manure ...	27.5	24.7	—	55.0	54.5	—	103	97	—	1400	1450	—	17.3	17.9	—	83	72	—

MALTING BARLEY EXPERIMENT.

Plumage Archer. Long Hoos Field, 1922.

Manures per Acre.	Dressed Grain.		Offal Grain per Acre.	Straw per Acre		Proportion of Total Grain to 100 of Total Straw.
	Yield per Acre.	Weight per Bushel.		Straw.	Total Straw	
	Bushels	lb.	lb.	lb.	cwt.	
Super. 3 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 1 cwt.	36·0	50·8	163	1213	17·1	104
Super. 3 cwt., Sul./Pot. 1½ cwt., Mur./Amm. 93 lb.	35·7	51·0	169	1388	18·5	96
Super. 3 cwt., Sul./Pot. 1½ cwt. ...	31·0	50·8	188	1263	17·0	93
Super. 3 cwt., Sul./Amm. 1 cwt. ...	30·0	50·3	175	975	14·1	107
Super. 3 cwt., Sul./Amm. 1 cwt., Mur./Pot. 1½ cwt.*	34·8	50·0	206	not	reco	ded.
Sul./Amm. 1 cwt., Sul./Pot. 1½ cwt.	36·8	50·3	191	1438	19·9	92
No Manure	28·6	50·5	184	1125	15·5	94

*Muriate of Potash applied on April 3rd. Other Manures on March 24th.

MISCELLANEOUS EXPERIMENTS.

Clover. Hoos Field, 1921 and 1922.

(Formerly Barley after Alsike).

Plot.	Manures per Acre.	Yield per Acre.	
		1921.	1922.
		cwt.	cwt.
1	Slag 8 cwt., Lime 10 cwt.	45·3	17·4
2	Farmyard Manure 14 tons, Super. 5 cwt., Lime 10 cwt.	53·8	17·9
3	Lime 10 cwt.	35·9	17·6
4	Super. 5 cwt., Lime 10 cwt., Sulph. Potash 1½ cwt. ...	40·6	19·6
5	Super. 5 cwt., Lime 10 cwt.	45·3	13·0
6	Lime 10 cwt.	41·1	13·0
7	Farmyard Manure 14 tons, Lime 10 cwt.	54·5	16·7
8	Slag 8 cwt.	42·9	11·4
9	Farmyard Manure 14 tons, Super. 5 cwt.	50·5	17·2
10	Control	36·8	14·1
11	Super. 5 cwt., Sulph. Potash 1½ cwt.	45·1	20·3
12	Super. 5 cwt.	49·1	14·3
13	Control	36·6	9·4
14	Farmyard Manure 14 tons	46·2	10·7
15	Horse Dung 14 tons, Lime 10 cwt.	35·3	6·7
16	Control	35·5	7·1
17	Horse Dung 14 tons	54·9	11·6
18	Super. 5 cwt.	39·7	6·3
19	Cattle Dung 14 tons, Lime 10 cwt.	50·5	13·0
20	Control	33·3	3·6
21	Cattle Dung 14 tons	41·5	5·8

Manures applied and Clover sown in 1920.

Barley. Hoos Field. Leguminous Strips, 1921, 1922.

Description of Plot.	Manurial Treatment	1921.						1922.					
		Dressed Grain.			Straw per Acre.	Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.	Dressed Grain.			Straw per Acre.	Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.
		Yield per Acre.	Weight per Bushel.	lb.				lb.	lb.	Yield per Acre.			
After Lucerne ...	Sulphate Amm. 1½ cwt. ... S. Amm. 1½ cwt. Super. 3 cwt.	14·8	56·5	134	688	10·2	85·5	27·2	51·4	188	1921	22·7	62·3
		37·8	57·3	154	1310	18·0	115·4	41·2	52·0	161	1884	22·2	92·6
After Red Clover	Sulphate Amm. 1½ cwt. ... S. Amm. 1½ cwt.; Super. 3 cwt.	12·1	56·4	122	555	8·4	85·2	25·2	50·6	134	1556	18·1	69·3
		31·1	57·4	134	1037	15·7	109·0	35·4	51·5	109	1579	18·0	96·0
After Alsike ...	Sulphate Amm. 1½ cwt. ... S. Amm. 1½ cwt.; Super. 3 cwt.	11·0	56·3	100	557	8·3	77·6	25·9	50·9	125	1481	17·2	74·8
		28·7	57·8	137	871	15·2	105·4	33·6	52·0	92	1421	16·5	99·6

Leguminous crops ploughed in November, 1911.

OUTSIDE CENTRES—MALTING BARLEY EXPERIMENTS.

Yield in bushels per acre; nitrogen expressed as per cent. of dry matter; value in shillings per quarter assigned by the expert valuation committee.

Treatment.	Rothamsted.			Cawkwell.			Wellingore.			Barneyhill.			Milverton.			Dereham.		
	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.
No Manure ...	25·8	1·60	s. 36	25·2	1·56	s. 28	36·1	1·76	s. 36	78·5	1·36	s. 48	26·0	1·53	s. 45	36·7	1·62	s. 31
Complete Manure ...	32·6	1·64	31	30·9	1·57	30	39·0	1·80	36	89·0	1·47	48	—	—	—	41·0	1·67	31
No Potash ...	27·0	1·69	31	22·6	1·49	30	43·5	1·78	36	84·0	1·50	48	27·2*	1·64	48	42·0	1·62	31
No Phosphate ...	33·0	1·63	32	28·8	1·49	30	40·5	1·89	32	85·0	1·47	45	—	—	—	36·0	1·69	31
No Nitrogen ...	28·2	1·58	31	23·1	1·50	30	37·3	1·72	40	75·5	1·38	53	23·3	1·48	55	30·0	1·66	31
	Woburn.			Orwell Park.			Walcott.			Eyton.			Dunmow.			Eye.		
	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.	Bushels.	Per cent. Nitrogen.	Value.
No Manure ...	42·5	1·78	s. 27	16·2	1·43	s. 65	56·9	1·70	s. 30	36·0	1·85	s. 32	51·0	1·75	s. 42	39·5	2·11	s. 30
Complete Manure ...	44·7	2·09	27	21·6	1·52	65	60·3	1·88	30	28·0	1·90	32	49·5	1·75	42	43·5	2·17	30
No Potash ...	41·8	2·10	27	24·6	1·53	64	58·7	1·79	30	45·2	2·10	32	47·0	1·85	42	44·1	2·13	30
No Phosphate ...	39·9	1·91	27	27·9	1·55	64	61·3	1·88	30	44·5	1·87	40	49·0	1·75	42	50·9	2·23	30
No Nitrogen ...	45·0	1·87	27	18·2	1·51	60	60·8	1·72	30	48·0	1·90	40	45·0	1·78	42	59·1	2·02	30

* Superphosphate only given in this case.