


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Report for 1927-28

[Full Table of Content](#)



Barley

Rothamsted Research

Rothamsted Research (1928) *Barley* ; Report For 1927-28, pp 26 - 30 - DOI:
<https://doi.org/10.23637/ERADOC-1-85>

No Potash.	Muriate of Potash, cwt. per acre.		
	1	1½	2
9.75	10.25	10.87	12.32

Each plot also received 1 cwt. sulphate of ammonia and 4 cwt. super-phosphate per acre.

Increase per cwt. muriate of potash—20 cwt. sugar beet.

It is somewhat curious that the factory determinations of the percentage of sugar in the roots were consistently lower than ours made on samples taken direct from the field.

CEREALS.

Under British conditions the most important fertilisers for cereals are the nitrogen compounds: these act with considerable uniformity, 1 cwt. sulphate of ammonia giving increases that range about 4½ bushels of wheat, 6 bushels of barley and 8 bushels of oats. The figures vary from season to season, but their relationships to the meteorological data are not yet fully known.

Barley. The experiments on barley are made in conjunction with the Institute of Brewing to ascertain the effect of soil, season and manuring on the yield and malting quality of barley. During 1927 and 1928 they have been continued at Rothamsted, Woburn, Wellingore (Lincs.), Chisleborough (S. Somerset), Fitzhead (Vale of Taunton) and Longniddry: they were, however, discontinued at the other centres, sufficient information having already been obtained. The most striking effect again was the increase given by 1 cwt. sulphate of ammonia per acre; this was greatest in years of low yield and least in years of high yield so that the effect of the fertiliser is to even up the results: the yields have been, in bushels per acre:—

	Lowest	Highest	Range of Variation
Without nitrogenous fertiliser	19.9	47.9	28.0
With nitrogenous fertiliser ...	32.5	44.4	11.9

So far as the data go, the increments of yield seem to be affected by:—

- (1) sufficient rainfall in spring to allow of tillering proportionate to the nitrogen supply;
- (2) sufficient sunshine in July to allow of head formation proportionate to the nitrogen supply.

The values of the increments for the past eight years have been at Rothamsted:—

	1923.	1927.	1924.	1928.	1921.	1925.	1922.	1926.
Increment of yield, bushels ...	12.6	10.4	7.7	7.0	6.8	6.0	5.0	-3.5
Yield without nitrogenous manure, bushels ...	19.9	23.6	22.1	28.6	27.2	25.0	31.0	47.9
Rainfall in inches:								
March ...	2.48	2.38	1.14	2.40	1.07	1.22	2.29	0.22
April ...	1.48	1.86	3.18	0.91	1.57	1.70	3.52	2.96
May ...	1.68	1.19	4.63	1.45	1.45	2.48	1.58	1.95
July sunshine hrs.	223.8	130.4	236.6	276.3	240.0	183.6	149.5	151.1

The high increments are associated with years of 3 to 4 inches March and April rain and 200 or more hours of July sunshine, 1927 being the only exception. A higher increment might have been expected in 1924, but the abnormally wet summer and autumn greatly encouraged the growth of weeds and protracted the harvest.

THE NEW NITROGENOUS MANURES.

Four nitrogenous fertilisers have been compared in detail. The results were, at Rothamsted, for barley :—

	1927.				1928.			
	Grain, bushels per acre.		Straw, cwt. per acre.		Grain, bushels per acre.		Straw, cwt. per acre.	
	Single dose.	Double dose.	Single dose.	Double dose.	Single dose.	Double dose.	Single dose.	Double dose.
Sulphate of ammonia	34.0	37.8	20.4	22.2	35.5	34.6	32.1	34.5
Muriate of ammonia	36.2	47.0	20.0	27.0	34.6	37.5	31.3	36.2
Urea... ..	32.8	43.8	20.0	24.3	35.0	35.8	31.1	32.8
Cyanamide	36.0	35.8	20.8	20.7	33.4	37.5	28.8	33.8
No nitrogen ...	23.6		15.4		28.6		24.4	

All fertilisers markedly increase the yield, with muriate of ammonia coming out best as usual in 1927, and quite well in 1928. In both years cyanamide has done well: it was applied a few days before sowing. Urea does not come up to muriate of ammonia.

These nitrogenous manures act best when they are applied with the seed—cyanamide should go on even earlier. Used as top dressing, even ammonium nitrate (nitrochalk) is ineffective, and when given late it only raised the percentage of nitrogen in the grain.

Barley sown.		No top dressing.	Nitrochalk applied:		
			May 12.	June 4.	June 19.
Grain, bushels per acre	...	31.1	30.8	33.0	31.4
Straw, cwt. per acre	...	30.1	31.9	32.1	29.5
Nitrogen per cent. in grain	...	2.075	2.118	2.110	2.160

Superphosphate on barley. The Hoosfield barley plots afford the best demonstration in the world of the effects of phosphate, potash and nitrogen starvation on the barley plant. In British practice, phosphate starvation is rare, the barley being grown only one or two years after a root crop which has been manured with a phosphatic fertiliser. The farmer is more interested in the other problem: the effect of doses of phosphate larger than are needed to supply the bare necessities of the plant. This depends very much on the season, but also on the soil. In the outside experiments the glacial drift soils at each of the three Norfolk centres have always responded to superphosphate. On other soils, however, the response varies from season to season: e.g., at Rothamsted, a response was obtained in 1927, but hardly in 1928 :—

Rothamsted.

	Grain, bushels per acre.		Straw, cwt. per acre.	
	1927.	1928.	1927.	1928.
Superphosphate	35.0	33.6	20.7	30.3
No superphosphate	31.4	32.8	18.9	29.4
Effect of superphosphate	+3.6	+0.8	+1.8	+0.9

The figures for the straw vary in the same direction. No connection between the meteorological data and the response to phosphate has yet been traced.

Potassic fertilisers on barley. The effect of potassic fertiliser, like that of phosphate, is much less marked than that of nitrogenous fertiliser. Few soils, except perhaps the thin chalks and light sands, show signs of potash starvation, and it is not clear that excess of potash over and above a margin of safety is advantageous: indeed, in some seasons, especially the good ones, sulphate of potash appears to be harmful. During its 77 successive years under barley, Hoosfield has passed through three stages: the first, when sulphate of potash not infrequently reduced the yield; the second, when it had no effect; and the third, when it increased the yield, potash starvation having set in at the end of about 32 years. The yields of grain have been, in bushels per acre:—

	Plot.	Early years.					Mean of 40 years. 1852-91.	Mean of 8 years. 1908-15.	Mean of 12 years 1916-27.
		1st 8 yrs. 1852-59.	1863.	1864.	1865.	1866.			
Complete artificial	4A	45.4	55.4	55.4	46.5	47.0	43.5	40.4	32.0
No potash	2A	44.9	61.6	58.5	48.4	50.5	42.75	28.5	26.5
Difference ¹		+0.5	-6.2	-3.1	-1.9	-2.5	+0.75	+11.8	+5.5

(1) Sulphates of soda and of magnesia are also omitted as well as sulphate of potash, but other plots show that their effects are relatively small.

Sulphate of potash caused a marked depression in yield of malting barley at Rothamsted in 1924 and at certain of the outside centres in other years: this is not common, but it appears to be a true result. The present data suggest that:—

- (1) in years of high spring rainfall and good ripening weather, *i.e.*, years favourable to the formation of well-matured grain of low nitrogen content, sulphate of potash may decrease the yield of barley;
- (2) in years unfavourable to ripening, sulphate of potash has less depressing effect and may even raise the yield of barley.

These variations are of the same kind as for wheat and potatoes, on both of which sulphate of potash acts beneficially in unfavourable seasons, and has less effect in good seasons, the badness of the season being in each instance measured by the yield of crop receiving no potash.

Actual depression of crop, however, seems to be confined to

barley, and apparently to sulphate of potash, for it has been observed with muriate of potash only in 1924; whether the chlorine ion is beneficial and the sulphate ion harmful, is not known.

EFFECT OF FERTILISERS ON COMPOSITION AND QUALITY OF THE GRAIN.

The percentage of nitrogen in the grain of barley depends on the amount of nitrogen the plant has taken from the soil and on the amount of carbohydrate it has synthesised during its growth. A high nitrogen uptake makes possible considerable growth and sufficient carbohydrate formation to over-balance the nitrogen: the grain then has a low nitrogen content. It depends on the favourableness of the conditions whether this possibility eventuates. Late sowing, or a check in growth due to drought, or a late supply of nitrogen to the plant, may so cut down the available time that the plant cannot make the necessary carbohydrate: the nitrogen content of the grain then becomes high. On the other hand, high rainfall in the weeks after sowing, by reducing the nitrate in the soil, but otherwise favouring growth, lowers the nitrogen content of the grain, as shown by the following data, obtained at Woburn:—

Nitrogen per cent.	2.01	1.95	1.71	1.57	1.23
Year	1925	1922	1923	1926	1924
Barley sown ...	March 31	April 19	April 10	March 4	March 11
Rainfall in inches after sowing.					
March	—	—	—	0.09	0.35
April	1.59	1.93	1.34	2.59	2.97
May 1st-15th inclusive	1.18	0.35	0.79	1.43	1.35

In sufficiently favourable conditions, sulphate of ammonia may still further increase the carbohydrate production and thus further reduce the proportion of nitrogen in the grain; in less favourable seasons, however, insufficient carbohydrate is produced and the nitrogen content of the grain may be raised. As the nitrogen content is already low in favourable and high in unfavourable seasons, it follows that sulphate of ammonia tends to lower the nitrogen content of the grain in years when it is low and to raise it in years when it is high. Larger quantities (2 cwt. per acre) tend to raise it in any case. The Rothamsted results have been:—

Percentage of Nitrogen in Grain.

	1925.	1926.	1927.	1928.	
No Nitrogen... ..	1.597	1.599	1.452	1.915	
Sulphate of Ammonia ...	1.585	1.711	1.442	2.029	Double dressing.
Muriate of Ammonia ...	1.552	1.684	1.438	1.985	2.220
					2.112

As in previous years muriate of ammonia gave grain of lower nitrogen content than sulphate of ammonia. Potassic fertilisers counteract to some extent the tendency for sulphate of

ammonia to raise the percentage of nitrogen; at Woburn, in 1928, the percentages of nitrogen in the grain were :—

Effect of Sulphate of Ammonia.		Effect of Sulphate of Potash.		Sulphate of Ammonia and Sulphate of Potash. No Superphosphate.
Superphosphate and Sulphate of Potash. + Sulphate of Ammonia.	No Sulphate of Ammonia.	Superphosphate and Sulphate of Ammonia. + Sulphate of Potash.	No Sulphate of Potash.	
1.372	1.310	1.372	1.398	1.346

The nitrogen content of the barley, more than any other single factor, determines its malting value. It is closely connected with the amount of "extract" and with the diastatic power of the malt, the extract varying inversely and the diastatic power directly with the nitrogen: it has also more subtle effects.

The investigations by Mr. Bishop on the nitrogen compounds of the barley grain have reached an interesting stage. The proportions of hordein, glutelin, and salt-soluble compounds are all connected with the percentage of nitrogen in the grain: for different samples of the same variety (Plumage-Archer) the glutelin increases proportionately to the nitrogen, the hordein increases more rapidly, and the salt-soluble compounds less rapidly than the nitrogen. The relationships are the same, whether the variation results from changes in soil, season or manuring; it appears, therefore, that the ratio glutelin/nitrogen may be a varietal constant of considerable interest to the breeder of barley for quality, and this is being determined for some of the new barleys grown by the National Institute of Agricultural Botany: the barleys are also being malted by the experts of the Institute of Brewing.

The large number of analyses of British-grown barleys made in recent years at Rothamsted has shown that the grinding barleys are richer in protein than is usually supposed. The figure quoted in the standard British tables is 8.6 per cent. of protein, corresponding to 1.38 per cent. of nitrogen; our results show that the figures have been, for barleys which buyers would not take for malting :—

		Valuation 45/- and less	Valuation below 40/-
1922	...	1.72	1.76
1923	...	1.73	1.95
1925	...	1.86	2.16
1926	...	1.58	1.65
Mean	...	1.72	1.88
Protein on conventional basis	...	10.8 %	11.8 %

The results show that less protein concentrates, such as decorticated cotton seed cake or meal, or decorticated ground nut cake, than is usually recommended, need be mixed with barley meal for feeding to farm animals.

WINTER-SOWN OATS AND WHEAT.

The experiments have given further information as to the effect of time of application of the sulphate of ammonia, and we are now able to sketch out the following as the probable