


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# The Soil at Rothamsted

[Full Table of Content](#)



---

## Appendix A: Descriptions of Sampled Profiles

### Rothamsted Research

Rothamsted Research (1995) *Appendix A: Descriptions of Sampled Profiles* ; The Soil At Rothamsted, pp 19 - 36 - DOI: <https://doi.org/10.23637/ERADOC-1-143>

## Appendix A: Descriptions of Sampled Profiles

### Profile 1 (B.W. Avery, 1945)

|                 |                                                                                                                                                      |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Park Grass (TL 122130)                                                                                                                               |
| Land use:       | old grass                                                                                                                                            |
| Soil series:    | Batcombe (intergrading to Hook)                                                                                                                      |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth.<br>U.S.D.A. (1992): Aquic Paleudalf<br>F.A.O. (1990): Chromic Luvisol (or Alisol) |
| Horizons (cm):  |                                                                                                                                                      |
| 0 - 20 Ah       | Thin turf mat over dark greyish brown, very slightly flinty silty clay loam. (U.S. silt loam)                                                        |
| 20 - 45 Eb      | Brown to yellowish brown, very slightly flinty silty clay loam (U.S. silt loam)                                                                      |
| 45 - 69 2Bt(g)1 | Strong brown, slightly or very slightly flinty silty clay loam (U.S. silty clay loam) with common paler mottles                                      |
| 69 - 90 2Bt(g)2 | Strong brown, slightly flinty silty clay (U.S. silty clay loam) with common reddish, greyish and paler brown mottles.                                |

### Profile 2 (B.W. Avery, 1945)

|                 |                                                                                                                                                                  |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | High Field (TL129131)                                                                                                                                            |
| Land use:       | old grass                                                                                                                                                        |
| Soil series:    | Batcombe                                                                                                                                                         |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Aquic Paleudalf or Paleudult<br>F.A.O. (1990): Chromic Luvisol (or Alisol) |
| Horizons (cm):  |                                                                                                                                                                  |
| 0 - 18 Ah       | Dark greyish brown flinty silty clay loam (U.S. silt loam)                                                                                                       |
| 18 - 30 Eb      | Brown flinty silty clay loam (U.S. silty clay loam)                                                                                                              |
| 30 - 60+ 2Bt(g) | Strong brown slightly flinty clay (U.S. clay) with paler and redder mottles becoming more prominent with depth                                                   |

### Profile 3 (B.W. Avery, 1945)

|                 |                                                                                                                                         |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Sawyers III (TL122142)                                                                                                                  |
| Land use:       | arable                                                                                                                                  |
| Soil series:    | Hornbeam                                                                                                                                |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Typic Paleudalf<br>F.A.O. (1990): Chromic Luvisol |
| Horizons (cm):  |                                                                                                                                         |
| 0 - 20 Ap       | Dark greyish brown, very slightly flinty clay loam (U.S. loam)                                                                          |
| 20 - 38 Eb      | Brown, slightly flinty clay loam (U.S. loam)                                                                                            |
| 38 - 90 2Bt(g)  | Strong brown to yellowish red, slightly flinty clay (U.S. clay) with paler and redder mottles                                           |

#### Profile 4 (B.W. Avery, 1945)

|                 |                                                                                                                                      |
|-----------------|--------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Harwood's Piece (TL134132)                                                                                                           |
| Land use:       | arable                                                                                                                               |
| Soil series:    | Notley                                                                                                                               |
| Classification: | England and Wales (1980): colluvial brown earth<br>U.S.D.A. (1992): Typic Hapludalf or Paleudalf?<br>F.A.O. (1990): Chromic Luvisol? |
| Horizons (cm):  |                                                                                                                                      |
| 0 - 20 Ap       | Dark greyish brown, flinty, slightly calcareous clay loam to silty clay loam (U.S. clay loam)                                        |
| 20 - 51+ Bw     | Brown, flinty silty clay loam (U.S. silt loam); few chalk fragments (augering proved brighter brown silty clay or clay at 80 cm)     |

#### Profile 5 (B.W. Avery, 1951)

|                 |                                                                                                                                                            |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Knott Wood (TL117133)                                                                                                                                      |
| Land use:       | woodland (hazel with ash and brambles)                                                                                                                     |
| Soil series:    | Winchester                                                                                                                                                 |
| Classification: | England and Wales (1980): typical paleo-argillic brown earth<br>U.S.D.A. (1992): Typic (or Vertic) Hapludalf<br>F.A.O. (1990): Chromic (or Vertic) Luvisol |
| Horizons (cm):  |                                                                                                                                                            |
| 0 - 10 Ah       | Very dark greyish brown (10YR3/2) flinty silty clay loam (U.S. silty clay loam)                                                                            |
| 10 - 23 Eb      | Brown flinty clay (U.S. clay)                                                                                                                              |
| 23 - 38 2Bt     | Strong brown (7.5 YR5/6-8) flinty clay (U.S. clay) with paler brown ped faces                                                                              |
| 38 - 66 2Bt     | Yellowish red (5 YR5/8) flinty clay (U.S. clay) with brown ped faces                                                                                       |
| 66 - 76 3Bt     | Yellowish red (5YR5/6) clay (U.S. clay) with large unbroken flints and black manganiferous segregations                                                    |
| 76+ 4Cu         | Fragmentary Chalk with clayey inclusions                                                                                                                   |

#### Profile 6 (B.W. Avery, 1951)

|                 |                                                                                                                      |
|-----------------|----------------------------------------------------------------------------------------------------------------------|
| Location:       | Knott Wood (TL116133)                                                                                                |
| Land use:       | woodland; clearing in hazel coppice with field layer of grass and herbs                                              |
| Soil series:    | Wallop                                                                                                               |
| Classification: | England and Wales (1980): brown rendzina<br>U.S.D.A. (1992): Typic Udorthent<br>F.A.O. (1990): Eutric Leptosol       |
| Horizons (cm):  |                                                                                                                      |
| 0 - 10 Ah       | Very dark greyish brown (7.5-10YR3/2) calcareous silty clay (U.S. silty clay); few chalk fragments                   |
| 10 - 23 Bw      | Brown to strong brown (7.5YR4/4-5/6) flinty, slightly calcareous clay (U.S. silty clay to clay); few chalk fragments |

|             |                                                                                             |
|-------------|---------------------------------------------------------------------------------------------|
| 23 - 46 BC  | Reddish yellow, flinty, very calcareous clay containing many soft chalk fragments           |
| 46 - 60+ Cu | Columns of hard white blocky chalk alternating with more friable chalk brash and brown clay |

#### Profile 7 (B.W. Avery, 1951)

|                 |                                                                                                                                      |
|-----------------|--------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Knott Wood (TL116133)                                                                                                                |
| Land use:       | woodland; hazel coppice with field layer of grass and herbs.                                                                         |
| Soil series:    | unnamed, fine silty over clayey, over chalky drift                                                                                   |
| Classification: | England and Wales (1980): Typical argillic brown earth<br>U.S.D.A. (1992): Typic Hapludalf<br>F.A.O. (1990): Haplic Alisol           |
| Horizons (cm):  |                                                                                                                                      |
| 0 - 13 Ah       | 2 cm turfy (not sampled) over dark brown (7.5YR4/3) flinty silty clay loam (U.S. silt loam)                                          |
| 13 - 30 Eb      | Brown (7.5YR4/4) flinty silty clay loam (U.S. silty clay loam)                                                                       |
| 30 - 56 Bt1     | Brown, flinty silty clay loam (U.S. silty clay loam)                                                                                 |
| 56 - 86 Bt2     | Reddish brown, flinty silty clay (U.S. silty clay loam) resting with an irregular boundary on chalky material with clayey inclusions |

#### Profile 8 (B.W. Avery, 1951)

|                 |                                                                                                                                        |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Knott Wood (TL115135)                                                                                                                  |
| Land use:       | woodland; hazel coppice with field layer of dogs mercury and bluebells                                                                 |
| Soil series:    | Panholes                                                                                                                               |
| Classification: | England and Wales (1980): typical brown calcareous earth<br>U.S.D.A. (1992): Rendollic Eutrochrept<br>F.A.O. (1990): Calcaric Cambisol |
| Horizons (cm):  |                                                                                                                                        |
| 1 - 0           | Litter layer (not sampled)                                                                                                             |
| 0 - 10 Ah       | Dark brown (7.5YR3/2), very slightly flinty, calcareous, humose silty clay loam (U.S. silt loam)                                       |
| 10 - 36 Bw      | Brown (7.5YR4/4), slightly flinty, calcareous silty clay loam (U.S. silty clay loam)                                                   |
| 36 - 61 BC      | Brown flinty, very calcareous, silty clay loam to silty clay (U.S. silty clay loam) containing abundant chalk fragments                |
| 61 + Cu         | Hard blocky Chalk with inclusions of brown loamy material and fresh white-coated flints                                                |

#### Profile 9 (B.W. Avery, 1951)

|                 |                                                                                             |
|-----------------|---------------------------------------------------------------------------------------------|
| Location:       | Knott Wood (TL115134)                                                                       |
| Land use:       | woodland (cut-over); hawthorn scrub with elder and brambles; field layer of grass and herbs |
| Soil series:    | Wold                                                                                        |
| Classification: | England and Wales (1980): typical argillic brown earth                                      |

|                |                                                                                                                         |
|----------------|-------------------------------------------------------------------------------------------------------------------------|
|                | U.S.D.A. (1992): Typic Hapludalf<br>F.A.O. (1990): Haplic Luvisol                                                       |
| Horizons (cm): |                                                                                                                         |
| 0 - 10 Ah      | 2 cm turfy over very dark greyish brown (7.5-10YR3/1) flinty silty clay loam (U.S. silt loam)                           |
| 10 - 25 Eb     | Brown (7.5YR4/3) flinty silty clay loam (U.S. silty clay loam)                                                          |
| 25 - 51 Bt     | Brown to strong brown (7.5YR4/4-5/6) flinty silty clay (U.S. silty clay)                                                |
| 51 - 55/76 BC  | Brown to pale reddish yellow, flinty, very calcareous silty clay loam to silty clay containing abundant chalk fragments |
| 55/76 + Cu     | Hard white blocky Chalk with loamy inclusions                                                                           |

#### Profile 10 (D.W. King, 1953)

|                 |                                                                                                                                                     |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Broadbalk Wilderness (TL121136)                                                                                                                     |
| Land use:       | woodland; self-sown sycamore, hazel and hawthorn; field layer predominately ivy.                                                                    |
| Soil series:    | Batcombe                                                                                                                                            |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Aquic (or Typic) Paleudalf<br>F.A.O. (1990): Chromic Luvisol. |
| Horizons (cm):  |                                                                                                                                                     |
| 0 - 15 Ah       | Very dark greyish brown (10YR3/2), flinty, slightly calcareous clay loam (U.S. loam)                                                                |
| 15 - 30 Eb/Bw   | Brown (10YR4/3), flinty, calcareous silty clay loam (U.S. clay loam to silty clay loam); few small chalk fragments                                  |
| 30 - 41 Eb      | Brown (7.5YR4/4), flinty, very slightly calcareous silty clay loam (U.S. silt loam)                                                                 |
| 41 - 91 2Bt(g)  | Reddish brown to yellowish red (5YR4/4-5/6) flinty clay (U.S. clay) with greyish and reddish mottles becoming more prominent with depth             |

#### Profile 11 (D.W. King, 1953)

|                 |                                                                                                                                             |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Broadbalk Wilderness (TL121136)                                                                                                             |
| Land use:       | grassland regularly mown, with hogweed and nettles.                                                                                         |
| Soil series:    | Batcombe                                                                                                                                    |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Aquic Paleudalf<br>F.A.O. (1990): Chromic Luvisol     |
| Horizons:       |                                                                                                                                             |
| 0 - 15 Ah       | Dark greyish brown (10YR4/2), slightly flinty, slightly calcareous silty clay loam (U.S. silt loam to silty clay loam); few chalk fragments |
| 15 - 25 Eb      | Brown (9YR4/4) slightly flinty, slightly calcareous silty clay loam (U.S. silt loam to silty clay loam); chalk fragments more common        |
| 25 - 51 2Bt(g)  | Brown to strong brown (7.5YR4/4-5/6), very slightly flinty, very                                                                            |

slightly calcareous silty clay (U.S. silty clay) with redder mottles and paler ped faces  
 51 - 76 + 2Bt(g)2 Strong brown (7.5YR5/6), very slightly flinty clay (U.S. clay) with common redder and greyer mottles

**Profile 12** (B.W. Avery & A.J. Thomasson, 1956)

Location: Meadow Field, Scout Farm (TL109130)  
 Land use: old grass  
 Soil series: Charity (intergrading to Hamble)  
 Classification: England and Wales (1980): typical argillic brown earth (possibly disturbed)  
 U.S.D.A. (1992): Typic Hapludalf  
 F.A.O. (1990): Haplic (or Chromic) Luvisol.

Horizons (cm):  
 0 - 13 Ah Thin turfy mat over dark greyish brown (10YR4/2) slightly flinty silt loam (U.S. silt loam)  
 13 - 76 Eb/Bw Brown (7.5YR4-5/4) very slightly flinty silt loam  
 76 - 117 Bt1 Brown very slightly flinty silty clay loam (U.S. silt loam) with faint paler and redder mottles and clay skins on ped surfaces  
 117 - 160 Bt2 Strong brown (7.5YR5/6) very slightly flinty silty clay loam (U.S. silty clay loam) with faint paler and redder mottles and clay skins on ped surfaces  
 160 + Becoming gravelly

**Profile 13** (B.W. Avery and D. Mackney, 1963)

Location: High Field (TL124129)  
 Land use: old grass  
 Soil series: Batcombe  
 Classification: England and Wales (1980): stagnogleyic paleo-argillic brown earth  
 U.S.D.A. (1992): Aquic Paleudalf  
 F.A.O. (1990): Chromic Luvisol

Horizons (cm):  
 0 - 6 Ah Dark greyish brown (10YR4/2) stoneless silty clay loam (U.S. silt loam); added chalk  
 6 - 20 Ah/Eb Dark greyish brown and dark yellowish brown (10YR4/4) very slightly flinty silty clay loam (U.S. silt loam)  
 20 - 35 Eb Yellowish brown (10YR5/4) slightly to moderately flinty silty clay loam (U.S. silt loam) with paler ped faces  
 35 - 65 2Bt(g)1 Brown (10YR5/4) to strong brown (7.5YR5/6-8) slightly flinty silty clay (U.S. silty clay) with paler and redder mottles  
 65 - 125 2Bt(g)2 Strong brown (7.5YR5/6) to yellowish red (5YR5/6-8) very slightly flinty clay (U.S. clay) with pale brown and greyish mottles becoming more prominent with depth

#### Profile 14 (R.G. Sturdy, 1966)

Location: Meadow Field, Scout Farm (TL107128)  
 Land use: old grass  
 Soil series: Rowton? (needs deeper sampling for positive identification)  
 Classification: England and Wales (1980): typical argillic brown earth or typical brown earth?  
 U.S.D.A. (1992): Typic Hapludalf or Glossudalf?  
 F.A.O. (1990): Haplic (or Chromic) Luvisol?

Horizons (cm):  
 0 - 11 Ah Dark greyish brown (10YR4/2), very slightly flinty silty clay loam (U.S. silt loam)  
 11 - 43 Bw/Eb Brown (7.5YR4/4) slightly flinty silt loam to silty clay loam (U.S. silt loam)  
 43 - 56 Bw/Eb Brown (7.5-10YR4/4) moderately flinty silt loam (U.S. silt loam)  
 56 - 81 2Bw/Eb Brown (7.5YR5/4), very gravelly sandy silt loam (U.S. silt loam); stones mainly rounded flint pebbles with some larger angular flints

#### Profile 15 (B.W. Avery, 1966)

Location: Scout Field, Scout Farm (TL108134)  
 Land use: old grass  
 Soil series: Hamble intergrading to Charity  
 Classification: England and Wales (1980): typical argillic brown earth  
 U.S.D.A. (1992): Typic Hapludalf  
 F.A.O. (1990): Haplic Luvisol

Horizons (cm):  
 0 - 18 Ah Dark greyish brown (10YR4/2-3) very slightly flinty silty clay loam (U.S. silt loam)  
 18 - 38 Eb Brown (10-7.5YR4/3) silty clay loam (U.S. silt loam) with occasional flints < 5 cm  
 38 - 61 Bt1 Brown (7.5YR4-5/4) silty clay loam (U.S. silt loam to silty clay loam) with occasional small flints as above  
 61 - 89 Bt2(g) Brown (7.5YR5/4-6) silty clay loam to silty clay (U.S. silty clay loam) with faint mottling in shades of brown; occasional flints as above

#### Profile 16 (B.W. Avery and R.G. Sturdy, 1966)

Location: Osier Field, Scout Farm (TL109132)  
 Land use: old grass  
 Soil series: Maplestead  
 Classification: England and Wales (1980): typical argillic brown earth  
 U.S.D.A. (1992): Typic Paleudalf (or Glossudalf)  
 F.A.O. (1990): Chromic Luvisol (or Eutric Podzoluvisol).

Horizons:  
 0 - 10 Ah Very dark greyish brown (10YR3/2) slightly flinty sandy silt loam to clay loam (U.S. silt loam)

|                           |                                                                                                                                                                                                            |
|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 10 - 25 Ah/Eb             | Dark greyish brown to brown (10YR4/2-3) moderately flinty (c.20%) sandy silt loam (U.S. silt loam); added chalk                                                                                            |
| 25 - 48 2Eb               | Brown (7.5YR4-5/4) very gravelly (mainly small angular flints) sandy silt loam (U.S. loam); wavy lower boundary                                                                                            |
| 48 - 58/66<br>2 Eb and Bt | Strong brown to yellowish brown (7.5-10YR5/4-6) very flinty (as above) sandy silt loam (U.S. silt loam) with light yellowish brown (10YR6/4) patches; irregular lower boundary with more clayey inclusions |
| 58/66 - 91 3Bt            | Reddish yellow to strong brown (7.5YR6/8-5/6), moderately to slightly flinty silty clay loam to silty clay (U.S. silty clay loam) with redder (5YR5/6) patches and thick clay skins                        |

**Profile 17 (B.W. Avery and R.G. Sturdy, 1966)**

|                 |                                                                                                                                                                                                             |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Flint Field, Scout Farm (TL112132)                                                                                                                                                                          |
| Land use:       | old grass with rushes, buttercups and tufted hair grass                                                                                                                                                     |
| Soil series:    | unnamed; fine silty over non-calcareous gravel                                                                                                                                                              |
| Classification: | England and Wales (1980): argillic humic gley soil<br>U.S.D.A. (1992): Typic Argiaquoll<br>F.A.O. (1990): Luvic (or Gleyic) Phaeozem                                                                        |
| Horizons (cm):  |                                                                                                                                                                                                             |
| 0 - 9 Ah1       | Very dark brown (10YR2/2) very slightly flinty (mainly < 1 cm) humose silty clay loam (U.S. silt loam)                                                                                                      |
| 9 - 25 Ah2      | Very dark grey (10YR3/1) slightly to moderately flinty (gravelly) humose silty clay loam (U.S. silt loam) with fine reddish brown (rusty) mottles, ochreous stains on flints, and paler grey lenses at base |
| 25 - 41 bAh1    | Very dark greyish brown (10YR3/2) slightly to moderately flinty (gravelly) silty clay loam (U.S. silt loam) with common fine rusty mottles                                                                  |
| 41 - 46 2bAh2   | Loose flint gravel with dark grey loamy matrix and common ochreous stains on flints                                                                                                                         |
| 46 - 71 2Eg     | Loose flint gravel with yellowish red mottles and ochreous deposits on flints, concentrated at 61-66 cm                                                                                                     |
| 71 - 90 3Btg    | Light grey to light greyish brown (10YR-2.5Y6/1) flinty sandy clay loam, sticky and plastic                                                                                                                 |

**Profile 18 (R.G. Sturdy, 1966)**

|                 |                                                                                                                                      |
|-----------------|--------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Webb's Field, Scout Farm (TL113128)                                                                                                  |
| Land use:       | arable                                                                                                                               |
| Soil series:    | Coombe                                                                                                                               |
| Classification: | England and Wales (1980): typical brown calcareous earth<br>U.S.D.A. (1992): Rendollic Eutrochrept<br>F.A.O. (1990): Eutric Cambisol |
| Horizons (cm):  |                                                                                                                                      |
| 0 - 25 Ap       | Dark greyish brown (10YR4/2) slightly flinty, slightly calcareous silty clay loam (U.S. silty clay loam)                             |

- 25 - 53/58 B Brown to dark brown (7.5YR4/4) slightly flinty, very slightly calcareous silty clay loam (U.S. silty clay loam) becoming slightly more clayey towards base; wavy lower boundary
- 53/58 - 86 Cu Light yellowish brown (10YR6/4) and very pale brown (10YR8/3), compact, extremely calcareous silty clay loam (U.S. silty clay loam) containing common sub-rounded chalk fragments and brown to strong brown loamy inclusions

**Profile 19** (B.W. Avery, 1968; Profile 1 in Avery & Bullock, 1969)

- Location: Broadbalk: intersection of Sections IB/2, plots 17/18 (TL122135)
- Land use: arable
- Soil series: Batcombe
- Classification: England and Wales (1980): stagnogleyic paleo-argillic brown earth  
U.S.D.A. (1992): Aquic Paleudalf  
F.A.O. (1990): Chromic (or Vertic) Luvisol
- Horizons (cm):
- 0 - 20 Ap Dark brown (10YR4/3-2) slightly flinty, slightly calcareous silty clay loam (U.S. silty clay loam); added chalk
- 20 - 51 2Bt(g)1 Yellowish red (5YR5/6-8) and strong brown moderately flinty clay (U.S. clay) with common red and lighter brown to greyish brown mottles
- 51 - 94 2Bt(g)2 Yellowish red, strong brown and red slightly flinty clay (U.S. clay) with common brown to light brownish grey (10YR6/2) mottles, mainly as vertical streaks

**Profile 20** (B.W. Avery, 1968; Profile 2 in Avery & Bullock, 1969)

- Location: Broadbalk: intersection of Sections 3/4, plots 3/5 (TL123136)
- Land use: arable
- Soil series: Batcombe
- Classification: England and Wales (1980): stagnogleyic paleo-argillic brown earth  
U.S.D.A. (1992): Aquic Paleudalf  
F.A.O. (1990): Chromic Luvisol
- Horizons (cm):
- 0 - 20 Ap Dark brown (10YR4/3) slightly flinty, slightly calcareous silty clay loam (U.S. silt loam); added chalk
- 20 - 51 2Bt(g)1 Yellowish red to strong brown (5 - 7.5YR5/6-8) very slightly flinty clay (U.S. clay) with few to common red mottles and paler brown ped faces
- 51 - 96 2Bt(g)2 Yellowish red and strong brown slightly flinty clay (U.S. clay) with red and brown (10YR5/3) to light grey (10YR6/1) mottles
- 96 - 142 2BC(g) Similar mottled clay containing common large, little broken nodular flints and fewer mottles

**Profile 21** (B.W. Avery, 1968; Profile 3 in Avery & Bullock, 1969)

- Location: Broadbalk: intersection of Sections 3/4 and plots 8/9 (TL123136)

|                   |                                                                                                                                                                     |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Land Use:         | arable                                                                                                                                                              |
| Soil Series:      | Batcombe                                                                                                                                                            |
| Classification:   | England and Wales (1980): stagnogleyic paleo-argillic brown earth.<br>U.S.D.A. (1992): Aquic Paleudalf (or Hapludalf)<br>F.A.O. (1990): Chromic Luvisol             |
| Horizons (cm):    |                                                                                                                                                                     |
| 0 - 18 Ap         | Dark greyish brown to brown (10YR4/2-3) slightly flinty, very slightly calcareous silty clay loam (U.S. silt loam)                                                  |
| 18 - 48 Bt        | Brown (7.5YR4/4) slightly flinty silty clay (U.S. silty clay)                                                                                                       |
| 48 - 71 2Bt(g)1   | Strong brown to reddish yellow (7.5YR5/6-8) very slightly flinty clay (U.S. clay) with more sand than above, many red and few to common light grey (10Y7/1) mottles |
| 71 - 102 2Bt(g)2  | Strong brown to yellowish red (7.5-5YR5/8) very slightly flinty clay (U.S. clay) with appreciable sand and common red and light grey mottles                        |
| 102 - 152 2BCt(g) | Yellowish red to strong brown stoneless clay loam to clay (U.S. clay loam) with fewer red and grey mottles, platy structure                                         |

**Profile 22** (B.W. Avery, 1968; Profile 4 in Avery & Bullock, 1969)

|                  |                                                                                                                                                        |
|------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:        | Broadbalk: intersection of Sections 2/3 & plots 7/8 (TL122136)                                                                                         |
| Land use:        | arable                                                                                                                                                 |
| Soil series:     | Batcombe, intergrading to Hook                                                                                                                         |
| Classification:  | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Aquic Paleudalf (or Hapludalf)<br>F.A.O. (1990): Chromic Luvisol |
| Horizons (cm):   |                                                                                                                                                        |
| 0 - 23 Ap        | Dark greyish brown to brown (10YR4/2-3) slightly flinty, very slightly calcareous silty clay loam (U.S. silt loam)                                     |
| 23 - 41 Eb/Bt(g) | Brown (7.5-10YR) very slightly flinty, very slightly calcareous silty clay loam (U.S. silt loam) with faint greyer and brighter brown mottles          |
| 41 - 66 Bt(g)1   | Brown to strong brown (7.5YR4/4-5/6) very slightly flinty silty clay (U.S. silty clay loam) with paler brown to light brownish grey (10YR6/2) mottles  |
| 66 - 94 Bt(g)2   | Strong brown (7.5YR5/6) very slightly flinty silty clay (U.S. silty clay) with mottles as above more common                                            |
| 94 - 117 2Bt(g)3 | Yellowish red (5YR5/8) and strong brown moderately flinty silty clay with common light brownish grey to light grey (10YR6/1) mottles                   |

**Profile 23** (B.W. Avery, 1968; Profile 5 in Avery & Bullock, 1969)

|                 |                                                                                              |
|-----------------|----------------------------------------------------------------------------------------------|
| Location:       | Broadbalk: south corner, headland beside plot 19 (TL124135)                                  |
| Land use:       | arable                                                                                       |
| Soil series:    | Notley                                                                                       |
| Classification: | England and Wales (1980): colluvial brown earth intergrading to gleyic argillic brown earth. |

|                   |                                                                                                                                                                                          |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                   | U.S.D.A. (1992): Aquic Paleudalf<br>F.A.O. (1990): Chromic Luvisol                                                                                                                       |
| Horizons (cm):    |                                                                                                                                                                                          |
| 0 - 25 Ap         | Dark greyish brown (10YR3/2) slightly flinty silty clay loam (U.S. silt loam)                                                                                                            |
| 25 - 41 Bw/Eb1    | Brown (7.5YR4/4) slightly flinty silty clay loam (U.S. silt loam)                                                                                                                        |
| 41 - 53 2Bw/Eb(g) | Brown (7.5YR4-5/4) moderately flinty silty clay loam (U.S. silt loam) with common, faint, paler brown (6/3) mottles                                                                      |
| 53 - 71 3Bt(g)1   | Brown (7.5YR5/4) very slightly flinty (< 2%) silty clay loam (U.S. silty clay loam) with many paler brown to light grey (10YR6-7/1) mottles and black manganiferous segregations         |
| 71 - 91 3Bt(g)2   | Strong brown (7.5YR5/6-8), brown, and light brownish grey (10YR6/2) prominently mottled, very slightly flinty silty clay (U.S. silty clay loam) with manganiferous segregations as above |

**Profile 24** (B.W. Avery, 1971; Profile 5 in Avery *et al.*, 1972)

|                 |                                                                                                                                                                                                                  |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Barnfield: section 6, between plots 4 and 5 (TL132132)                                                                                                                                                           |
| Land use:       | arable                                                                                                                                                                                                           |
| Soil series:    | Batcombe                                                                                                                                                                                                         |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Aquic Paleudalf<br>F.A.O. (1990): Chromic (or Vertic) Luvisol                                                              |
| Horizons (cm):  |                                                                                                                                                                                                                  |
| 0 - 18 Ap       | Dark brown (10YR4/3) very slightly flinty, very slightly calcareous silty clay loam (U.S. silty clay loam)                                                                                                       |
| 18 - 35 2Bt(g)1 | Strong brown (7.5YR5/6) to yellowish red, stoneless clay (U.S. clay) with fine redder mottles and paler brown ped faces                                                                                          |
| 35 - 55 2Bt(g)2 | Strong brown very slightly flinty clay (U.S. clay) with red and paler brown mottles and paler brown ped faces                                                                                                    |
| 55 - 90 2Bt(g)3 | Strong brown to reddish yellow (7.5YR5/8) very slightly flinty clay (U.S. clay) with many red and common light brownish grey (10YR6/2) to grey (6/1) mottles, particularly around stones and on structural faces |

**Profile 25** (B.W. Avery, 1971; Profile 7 in Avery *et al.*, 1972)

|                 |                                                                                                                                    |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Barnfield: section 3, between plots 4 and 5 (TL132134)                                                                             |
| Land use:       | arable                                                                                                                             |
| Soil series:    | Carstens (intergrading to Notley)                                                                                                  |
| Classification: | England and Wales (1980): typical paleo-argillic brown earth<br>U.S.D.A. (1992): Typic Paleudalf<br>F.A.O. (1990): Chromic Luvisol |
| Horizons (cm):  |                                                                                                                                    |
| 0 - 15 Ap1      | Dark brown (10YR4/3) slightly flinty, slightly calcareous silty clay loam (U.S. silt loam)                                         |
| 15 - 30 Ap2     | Dark brown (10YR4/3) slightly flinty (more than above), very slightly calcareous silty clay loam (U.S. silt loam)                  |
| 30 - 41 Bt      | Brown (10-7.5YR4/3-4) moderately flinty silty clay loam (U.S. silty                                                                |

|              |                                                                                        |
|--------------|----------------------------------------------------------------------------------------|
| 41 - 70 2Bt1 | clay loam)<br>Brown to yellowish red (7.5-5YR4/4-5/6) slightly flinty clay (U.S. clay) |
| 70 - 90 2Bt2 | Yellowish red (5YR5/6-8) slightly flinty clay (U.S. clay) with few paler brown mottles |

**Profile 26** (B.W. Avery, 1971; Profile 8 in Avery *et al.*, 1972)

|                 |                                                                                                                                                                 |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Barnfield, section 1, between plots 4 & 5 (TL132135)                                                                                                            |
| Land Use:       | arable                                                                                                                                                          |
| Soil series:    | Batcombe (intergrading to Givendale)                                                                                                                            |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Aquic (or Vertic) Paleudalf<br>F.A.O. (1990): Chromic (or Vertic) Luvisol |
| Horizons (cm):  |                                                                                                                                                                 |
| 0 - 19/21 Ap    | Dark brown (10YR4/3) slightly flinty, very slightly calcareous, silty clay loam to silty clay (U.S. silty clay loam)                                            |
| 19/21 - 35 2Bt1 | Strong brown to yellowish red (7.5-5YR4-5/6) very slightly flinty clay (U.S. clay) with faint brownish mottles                                                  |
| 35 - 55 2Bt2(g) | Strong brown to yellowish red, very slightly flinty clay (U.S. clay) with paler brown (7.5YR5/4) structural faces                                               |
| 55 - 90 2Bt3(g) | Brown and yellowish red, slightly flinty clay (U.S. clay) with common to many red (2.5YR4/6) and few to common light brownish grey to light grey mottles        |

**Profile 27** (B.W. Avery, 1971; Profile 11 in Avery *et al.*, 1972)

|                 |                                                                                                                                                                                                      |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Barnfield; section 3, between plots 7 & 8 (TL133134)                                                                                                                                                 |
| Land use:       | arable                                                                                                                                                                                               |
| Soil series:    | Notley                                                                                                                                                                                               |
| Classification: | England and Wales (1980): colluvial brown earth<br>U.S.D.A. (1992): Typic Hapludalf (or Paleudalf)?<br>F.A.O. (1990): Chromic Luvisol?                                                               |
| Horizons (cm):  |                                                                                                                                                                                                      |
| 0 - 18 Ap1      | Dark greyish brown (10YR4/2-3) slightly to moderately flinty, slightly calcareous silty clay loam (U.S. silt loam)                                                                                   |
| 18 - 28 Ap2     | Dark brown (10YR4/3) slightly to moderately flinty, slightly calcareous silty clay loam (U.S. silt loam), more compact than above                                                                    |
| 28 - 50 Bw      | Brown (7.5-10YR4/4) slightly flinty silty clay loam (U.S. silt loam)                                                                                                                                 |
| 50 - 70 2Eb/Bt  | Paler yellowish brown (10YR5/4) very flinty sandy silt loam (U.S. loam to silt loam) becoming redder and finer in texture in places near lower boundary; stones mainly less than 5 cm (loamy gravel) |
| 70 - 90 3Bt(g)  | Strong brown to reddish yellow (7.5YR5-6/6) stoneless silty clay loam to silty clay (U.S. silty clay loam) with pale brown to light yellowish brown (10YR6/3-4) mottles                              |

### Profile 28 (B.W. Avery, 1971)

|                 |                                                                                                                             |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------|
| Location:       | Stubbings Field, Scout Farm (TL114131)                                                                                      |
| Land use:       | arable                                                                                                                      |
| Soil series:    | Charity (intergrading to Notley)                                                                                            |
| Classification: | England and Wales (1980): typical argillic brown earth<br>U.S.D.A. (1992): Typic Hapludalf<br>F.A.O. (1990): Haplic Luvisol |
| Horizons (cm):  |                                                                                                                             |
| 0 - 25 Ap       | Dark greyish brown (10YR4/2) slightly flinty silty clay loam (U.S. silt loam)                                               |
| 25 - 52 Eb      | Brown (7.5-10YR4/4) slightly flinty silty clay loam (U.S. silt loam)                                                        |
| 52 - 90 Bt      | Brown (7.5YR4/4) moderately flinty silty clay loam (U.S. silt loam to silty clay loam)                                      |

### Profile 29 (B.W. Avery, 1971)

|                 |                                                                                                                                                    |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Stubbings Field, Scout Farm (TL114133)                                                                                                             |
| Land use:       | arable                                                                                                                                             |
| Soil series:    | unnamed; fine silty over clayey, on chalky drift                                                                                                   |
| Classification: | England and Wales (1980): typical argillic brown earth<br>U.S.D.A. (1992): Typic Hapludalf<br>F.A.O. (1990): Haplic Luvisol                        |
| Horizons (cm):  |                                                                                                                                                    |
| 0 - 25 Ap       | Dark brown (10YR3/3, 4/3 rubbed); slightly flinty, slightly calcareous silty clay loam (U.S. silty clay loam)                                      |
| 25 - 42/55 Bt   | Brown (7.5YR4/4) slightly flinty silty clay (U.S. silty clay loam); few rotting chalk fragments; fine earth mainly non-calcareous                  |
| 42/55 - 90 Cu   | Brown and pale brown, very slightly flinty, very calcareous silty clay loam (U.S. silty clay loam) containing many soft, weathered chalk fragments |

### Profile 30 (B.W. Avery, 1971)

|                 |                                                                                                                                                            |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Great Knott (TL117136)                                                                                                                                     |
| Land use:       | arable                                                                                                                                                     |
| Soil series:    | Winchester                                                                                                                                                 |
| Classification: | England and Wales (1980): typical paleo-argillic brown earth<br>U.S.D.A. (1992): Typic (or Vertic) Hapludalf<br>F.A.O. (1990): Chromic (or Vertic) Luvisol |
| Horizons (cm):  |                                                                                                                                                            |
| 0 - 20 Ap       | Dark brown (10YR4/3) slightly to moderately flinty, slightly calcareous clay loam to clay (U.S. silty clay loam); added chalk                              |
| 20 - 43 2Bt1    | Yellowish red (5 YR4-5/8) flinty clay (U.S. clay) with some brown (7.5-5YR4/4) faces around flints                                                         |
| 43 - 67 2Bt2    | Yellowish red (5YR4-5/8) flinty clay (U.S. clay) with common slickensided structural faces; flints include large little-broken nodules                     |

67 - 90 Cu Chalk fragments in a matrix of finely divided chalk with few flints and inclusions of yellowish red clay near upper boundary

**Profile 31 (B.W. Avery, 1971)**

Location: Whittlocks (TL118143)  
 Land use: arable  
 Soil series: Hamble  
 Classification: England and Wales (1980): typical argillic brown earth  
 U.S.D.A. (1992): Typic Hapludalf  
 F.A.O. (1990): Haplic Luvisol

Horizons (cm):  
 0 - 18 Ap Dark greyish brown to brown (10YR4/2-3) very slightly flinty silty clay loam (U.S. silt loam)  
 18 - 38 Bt1 Brown (7.5YR4/4) very slightly flinty silty clay loam (U.S. silt loam)  
 38 - 68 Bt2 Brown (7.5YR4/4) stoneless silty clay loam (U.S. silty clay loam) with few faint paler brown mottles  
 68 - 90 Bc(g) Brown stoneless silty clay loam (U.S. silt loam) with paler brown and dark greyish brown mottles

**Profile 32 (B.W. Avery, 1973)**

Location: Little Knott (TL121135)  
 Land use: arable  
 Soil series: Hook  
 Classification: England and Wales (1980): gleyic argillic brown earth  
 U.S.D.A. (1992): Typic Paleudalf  
 F.A.O. (1990): Chromic Luvisol

Horizons (cm):  
 0 - 20 Ap Dark greyish brown to dark brown (10YR4/2-3) very slightly flinty, slightly calcareous silty clay loam (U.S. silt loam)  
 20 - 40 Eb Brown (7.5-10YR4/4) very slightly flinty silty clay loam (U.S. silt loam)  
 40 - 65 Bt(g)1 Brown (7.5YR4-5/4) stoneless silty clay loam (U.S. silty clay loam) with common fine strong brown (7.5YR5/6) mottles  
 65 - 91 Bt(g)2 Strong brown (7.5YR5/6) and brown (7.5YR4-5/4), mottled, stoneless silty clay loam to silty clay (U.S. silty clay loam) with some paler brown (5/2-3) structural faces  
 91 - 136 Bct(g)1 As above, with few light grey (10YR6/2) faces  
 136 - 182 Bct(g)2 Strong brown stoneless silty clay (U.S. silty clay loam) with many paler brown to light grey mottles

**Profile 33 (B.W. Avery, 1973)**

Location: Little Knott (TL121135)  
 Land use: arable  
 Soil series: Hook (intergrading to Batcombe)  
 Classification: England and Wales (1980): gleyic argillic brown earth

|                   |                                                                                                                                                                                                 |
|-------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                   | U.S.D.A. (1992): Typic (or Aquic) Paleudalf<br>F.A.O. (1990): Chromic (or Haplic) Luvisol                                                                                                       |
| Horizons (cm):    |                                                                                                                                                                                                 |
| 0 - 18 Ap         | Dark greyish brown to dark brown (10YR4/2-3) very slightly flinty, slightly calcareous silty clay loam (U.S. silt loam)                                                                         |
| 18 - 37 Eb        | Brown (7.5-10YR4/4) very slightly flinty silty clay loam (U.S. silt loam)                                                                                                                       |
| 37 - 55/60 Bt(g)1 | Brown (7.5YR5/4) very slightly flinty silty clay loam (U.S. silt loam) with few faint paler and stronger brown mottles                                                                          |
| 55/60 - 90 Bt(g)2 | Brown and strong brown (7.5YR5/6-8), stoneless to very slightly flinty silty clay loam to silty clay (U.S. silty clay loam) with few light grey mottles mainly associated with structural faces |
| 90 - 135 BCt(g)   | Strong brown (7.5YR5/6) and brown (7.5YR5/4) stoneless silty clay loam (U.S. silty clay loam) with light grey mottles as above                                                                  |
| 135 - 300 BC(g)   | As above, mainly strong brown with few grey to light grey sub-vertical veins and mottles                                                                                                        |

**Profile 34 (R.G. Sturdy, 1976)**

|                 |                                                                                                                                                      |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Blackhorse Field, Scout Farm (TL105130)                                                                                                              |
| Land use:       | arable                                                                                                                                               |
| Soil series:    | Batcombe                                                                                                                                             |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic brown earth<br>U.S.D.A. (1992): Typic (or Glossic) Paleudalf<br>F.A.O. (1990): Chromic Luvisol |

|                  |                                                                                                                                                                                                                    |
|------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Horizons (cm):   |                                                                                                                                                                                                                    |
| 0 - 27 Ap        | Dark greyish brown (10YR4/2), slightly flinty, very slightly calcareous, clay loam to silty clay loam (U.S. silt loam to loam)                                                                                     |
| 27 - 38 Eb       | Brown (7.5YR5/4) and dark brown (7.5YR4/3) moderately flinty (including common flint pebbles) silty clay loam (U.S. silt loam)                                                                                     |
| 38 - 70 Bt(g)    | Brown (7.5YR4-5/4) moderately flinty (as above) silty clay loam (U.S. silt loam to silty clay loam) with reddish yellow (7.5YR6/6) mottles and pale brown (10YR6/3) inclusions extending locally to greater depths |
| 70 - 110 2Bt(g)  | Red (2.5YR3/6) moderately flinty clay (U.S. clay) with many yellowish brown (10-7.5YR5/4) mottles and brown (7.5YR5/4) ped faces                                                                                   |
| 110 - 130 2BC(g) | Red (2.5YR3/6) moderately flinty clay (U.S. clay) with common yellowish brown to pale brown (7.5YR5/2) mottles                                                                                                     |

**Profile 35 (B.W. Avery, 1976)**

|                 |                                                                                                                            |
|-----------------|----------------------------------------------------------------------------------------------------------------------------|
| Location:       | Meadow Field, Scout Farm (TL107127)                                                                                        |
| Land use:       | arable                                                                                                                     |
| Soil series:    | Carstens (intergrading to Batcombe)                                                                                        |
| Classification: | England and Wales (1980): typical (grading to stagnogleyic) paleo-argillic brown earth<br>U.S.D.A. (1992): Typic Paleudalf |

|                    |                                                                                                                                                                                                    |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                    | F.A.O. (1990): Chromic Luvisol                                                                                                                                                                     |
| Horizons (cm):     |                                                                                                                                                                                                    |
| 0 - 25 Ap          | Dark greyish brown (10YR4-3/2) slightly flinty, very slightly calcareous clay loam (U.S. silt loam)                                                                                                |
| 25 - 35/50 Eb      | Brown (7.5YR4/4) moderately (varying from slightly to very) flinty silty clay loam (U.S. silt loam) with wavy lower boundary and paler brown (7.5YR5-6/4) inclusions in lower part                 |
| 35/50 - 65/92 Bt1  | Reddish brown to yellowish red (5YR4/4-5/6) moderately flinty silty clay loam (U.S. silty clay loam) with brown (7.5YR4-5/4) inclusions and similar colours on some ped faces; wavy lower boundary |
| 65/92 - 120 Bt2(g) | Yellowish red (5YR5/8) to reddish yellow (7.5YR6/6-8) slightly to moderately flinty clay (U.S. clay to clay loam) with paler brown (7.5YR5/4-3) ped faces and few redder mottles                   |

#### Profile 36 (R.G. Sturdy, 1976)

|                 |                                                                                                                                                       |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Osier Field, Scout Farm (TL110128)                                                                                                                    |
| Land use:       | old grass                                                                                                                                             |
| Soil series:    | Charity                                                                                                                                               |
| Classification: | England and Wales (1980): typical argillic brown earth.<br>U.S.D.A. (1992): Typic Paleudalf<br>F.A.O. (1990): Chromic (or Haplic) Luvisol             |
| Horizons (cm):  |                                                                                                                                                       |
| 0 - 10 Ah       | Dark greyish brown (10YR4/2) very slightly flinty silty clay loam (U.S. silt loam)                                                                    |
| 10 - 25 AE      | Dark greyish brown to dark brown (10YR4/2-3) very slightly flinty silty clay loam (U.S. silt loam)                                                    |
| 25 - 40 Eb      | Brown (7.5YR4/4) very slightly flinty silty clay loam (U.S. silt loam)                                                                                |
| 40 - 70/84 Bt1  | Brown (7.5YR5/4) slightly flinty silty clay loam (U.S. silty clay loam), with darker brown (7.5YR4/2-3) inclusions and ped faces; wavy lower boundary |
| 70/84 - 105 Bt2 | Strong brown (7.5YR5/6) slightly flinty silty clay loam (U.S. silty clay loam) with reddish brown (5YR4/4) clay coats on ped faces                    |
| 105 - 170 BC(t) | Brown to strong brown (7.5YR5/5) very slightly flinty silty clay (U.S. silty clay loam) with few paler brown (7.5YR5/2) veins and ped faces           |

#### Profile 37 (R.G. Sturdy, 1976)

|                 |                                                                                                                            |
|-----------------|----------------------------------------------------------------------------------------------------------------------------|
| Location:       | Ver Field, Scout Farm (TL111128)                                                                                           |
| Land use:       | old grass (made ground in 1993)                                                                                            |
| Soil series:    | unnamed; fine silty over non-calcareous gravel                                                                             |
| Classification: | England and Wales (1980): argillic humic gley soil<br>U.S.D.A. (1992): Mollic Endoaqualf<br>F.A.O. (1990): Gleyic Luvisol? |
| Horizons (cm):  |                                                                                                                            |
| 0 - 18 Ahg1     | Very dark grey (10YR3/1) very slightly flinty, humose sandy silt                                                           |

|                |                                                                                                                                                  |
|----------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
|                | loam (U.S. loam) with common fine strong brown (7.5YR4/6) mottles along root channels                                                            |
| 18 - 36 Ahg2   | Dark greyish brown to dark grey (10YR4/2-1) slightly flinty sandy silt loam to silty clay loam (U.S. silt loam) with mottles as above            |
| 36 - 50 Eg     | Dark greyish brown (10YR4/2) moderately flinty silty clay loam (U.S. silt loam) with few mottles                                                 |
| 50 - 58 Btg    | Dark grey (10YR4/1) slightly flinty silty clay loam (U.S. silt loam) with many dark yellowish brown (10YR4/4) to strong brown (7.5YR5/6) mottles |
| 58 - 95 2Btg   | Dark grey (10YR4/1) very to extremely flinty sandy clay loam (loamy gravel) with many strong brown mottles                                       |
| 95 - 110 2BCgf | Strong brown (7.5YR5/8) very flinty clay loam (U.S. loam) with many pale brown (7.5YR4/2) mottles                                                |

**Profile 38 (B.W. Avery, 1976)**

|                 |                                                                                                                                                  |
|-----------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Ver Field, Scout Farm (TL112128)                                                                                                                 |
| Land use:       | old grass (made ground in 1993)                                                                                                                  |
| Soil series:    | Tidmarsh                                                                                                                                         |
| Classification: | England and Wales (1980): gleyic argillic brown earth<br>U.S.D.A. (1992): Mollic Hapludalf (or Glossudalf)<br>F.A.O. (1990): Chromic Luvisol?    |
| Horizons (cm):  |                                                                                                                                                  |
| 0 - 15 Ah1      | Very dark greyish brown (10YR3/2) very slightly flinty, humose clay loam (U.S. loam)                                                             |
| 15 - 32 Ah2     | Dark greyish brown (10YR4/2) moderately to very flinty clay loam (U.S. loam)                                                                     |
| 32 - 50 Bw?     | Brown very flinty sandy silt loam or sandy loam (loamy gravel of subangular flints with few flint pebbles)                                       |
| 50 - 85 2Eb     | Brown to light yellowish brown (10YR5/4) very to extremely flinty sandy loam with paler mottles (loamy gravel with inclusions of 'clean' gravel) |
| 85 - 130 2Btgf  | Prominently mottled, light yellowish brown and strong brown to yellowish red, very flinty sandy clay loam with clayey coats and inclusions       |

**Profile 39 (B.W. Avery, 1976)**

|                 |                                                                                                                               |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Ver Field, Scout Farm (TL112129)                                                                                              |
| Land use:       | old grass (made ground in 1993)                                                                                               |
| Soil series:    | Binstead (phase with superficial silty calcareous alluvium)                                                                   |
| Classification: | England and Wales (1980): typical argillic gley soil<br>U.S.D.A. (1992): Aeric Endoaqualf<br>F.A.O. (1990): Gleyic Luvisol?   |
| Horizons (cm):  |                                                                                                                               |
| 0 - 12 Ah1      | Very dark grey to greyish brown (10YR3/1-2) stoneless, calcareous, humose silty clay loam (U.S. silt loam to silty clay loam) |
| 12 - 25 Ah2(g)  | Dark greyish brown, very slightly flinty, calcareous silty clay loam                                                          |

- 25 - 48 2Eg (U.S. silt loam) with common strong brown to yellowish red mottles  
Brown to greyish brown very flinty sandy silt loam to sandy loam  
(loamy gravel) with fine strong brown mottles
- 48 - 150 2Eg/Btgf Pale brown, very to extremely flinty sandy loam (loamy gravel) with  
yellowish red to dark reddish brown ferruginous bands; matrix  
becoming sandy clay loam or finer below 90 cm; wet at base

**Profile 40** (L. De Klerk and C. Van Beek, 1977)

- Location: Knott Wood (TL117135)  
Land use: deciduous woodland  
Soil series: Carstens  
Classification: England and Wales (1980): typical paleo-argillic brown earth  
U.S.D.A. (1992): Typic Hapludalf  
F.A.O. (1990): Chromic Luvisol
- Horizons (cm):
- 0 - 15 Ah Dark brown (10YR3/3) slightly flinty silty clay loam (U.S. silt loam)
- 15 - 30 Eb Brown to dark brown (7.5YR4/4) moderately flinty silty clay loam  
(U.S. silt loam)
- 30 - 50 Bt Strong brown (7.5YR5/6) slightly flinty silty clay loam (U.S. silty  
clay loam)
- 50 - 90 2Bt Reddish brown to yellowish red (5YR4/5) very slightly flinty clay  
(U.S. silty clay to clay) with common fine red (2.5YR5/6) mottles
- 90 - 100 Cu Pinkish grey (7.5YR7/2) extremely calcareous silty clay (U.S. silty  
clay loam) containing many chalk fragments

**Profile 41** (L. De Klerk and C. Van Beek, 1977)

- Location: Little Knott (TL120134)  
Land use: temporary grass  
Soil series: Hook (intergrading to Batcombe)  
Classification: England and Wales (1980): gleyic argillic brown earth  
U.S.D.A. (1992): Typic Paleudalf  
F.A.O. (1990): Haplic Luvisol
- Horizons (cm):
- 0 - 24 Ap Dark brown (10YR4/3) very slightly flinty, very slightly calcareous  
silty clay loam (U.S. silt loam)
- 24 - 40 Bt1 Dark yellowish brown (10YR4/4) very slightly flinty silty clay loam  
(U.S. silty clay loam)
- 40 - 65 Bt2(g) Brown (7.5YR5/4) stoneless silty clay loam (U.S. silty clay loam)  
with common strong brown (7.5YR5/7) mottles
- 65 - 105 2Bt3(g) Brown (7.5YR5/4) stoneless to very slightly flinty silty clay (U.S.  
silty clay loam) with common strong brown mottles
- 105 - 110 2BCg Strong brown (7.5YR5/7) stoneless to very slightly stony silty clay  
to clay (U.S. silty clay) with common yellowish red (5YR5/6)  
mottles

**Profile 42 (L. De Klerk and C. Van Beek, 1977)**

|                 |                                                                                                                                         |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| Location:       | Bones Close                                                                                                                             |
| Land use:       | arable                                                                                                                                  |
| Soil series:    | Batcombe                                                                                                                                |
| Classification: | England and Wales (1980): stagnogleyic paleo-argillic grown earth<br>U.S.D.A. (1992): Aquic Paleudalf<br>F.A.O. (1990): Chromic Luvisol |
| Horizons (cm):  |                                                                                                                                         |
| 0 - 20 Ap       | Dark brown (10YR4/3) slightly flinty, very slightly calcareous silty clay loam (U.S. silt loam)                                         |
| 20 - 40 Bt1     | Strong brown (7.5YR5/6) and dark yellowish brown (10YR4/4), slightly flinty faintly mottled silty clay loam (U.S. silty clay loam)      |
| 40 - 120 2Bt(g) | Strong brown (7.5YR5/8) very slightly flinty clay (U.S. clay) with common pinkish grey (7.5YR7/2) mottles                               |