


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Memoranda of the Field Experiments at Rothamsted, May 1872



[Full Table of Content](#)

Experiments on the Economy of Nitrogenous Manures

Rothamsted Research

Rothamsted Research (1873) *Experiments on the Economy of Nitrogenous Manures* ; Memoranda Of The Field Experiments At Rothamsted, May 1872, pp 11 - 11 - DOI:
<https://doi.org/10.23637/ERADOC-1-236>

(11)

EXPERIMENTS WITH A VIEW TO ECONOMY IN THE USE OF EXPENSIVE NITROGENOUS MANURES.

It is found that generally less than half the nitrogen supplied in such manures as guano, ammonia-salts, or nitrate of soda, is recovered in the increase of the crop for which they are used; that a considerable quantity may remain in the soil in a comparatively inactive state, yielding increase very slowly; and that a considerable quantity may be carried away by drainage, and lost. It seemed desirable, therefore, to

commence a series of experiments to determine whether any saving can be effected by applying comparatively small quantities near to the seed, instead of larger amounts in the usual mode of broadcast sowing and harrowing-in.

It is also intended to make experiments with a view to ascertain the best periods of the year for the application of such manures to different crops.

FIRST SEASON, 1871.

Experiments upon Wheat. Little Hoos Field. Plots $\frac{1}{2}$ acre each.

PLOT No.	MANURES PER ACRE, &c.	PRODUCE PER ACRE.		
		Dressed Corn.		Total Straw.
		Quantity.	Weight per Bushel.	
1	Unmanured. Seed 1 bushel, dibbled 6 inches apart in the rows	Bushels, 23 $\frac{3}{4}$	lbs. 59.3	cwts. 24 $\frac{1}{2}$
2	{ 146 lbs. (1) Sulphate Ammonia. Seed 1 bushel; } { Holes dibbled 6 inches apart in the rows; manure (mixed with Ashes) put in, and seed above }	31 $\frac{1}{2}$	59.1	36 $\frac{1}{2}$
3	{ 292 lbs. Sulphate Ammonia. Seed 1 bushel; } { Manure (mixed with Ashes) sown broadcast, seed dibbled 6 inches apart in the rows }	28 $\frac{3}{4}$	58.3	35 $\frac{1}{2}$

(1) Containing Nitrogen equal to that in 15 bushels of grain, with its average proportion of Straw.

Experiments upon Barley. Thirty-acres Field. Plots $\frac{1}{2}$ acre each.

PLOT No.	MANURES PER ACRE, &c.	PRODUCE PER ACRE.		
		Dressed Corn.		Total Straw.
		Quantity.	Weight per Bushel.	
1	Unmanured. Seed 3 bushels; drilled	Bushels, 40 $\frac{1}{2}$	lbs. 53.9	cwts. 24 $\frac{1}{2}$
2	{ 1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels; } { Manures mixed with Ashes and sown broadcast; seed drilled .. }	49 $\frac{1}{2}$	53.3	30 $\frac{1}{2}$
3	{ 1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels; .. } { Manures mixed with Ashes and drilled; seed drilled above .. }	49 $\frac{1}{2}$	53.4	28 $\frac{1}{2}$
4	{ 1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels; } { Manures, Ashes, and Seed mixed, and drilled together .. }	51	53.0	30 $\frac{1}{2}$
5	{ 1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 1 $\frac{1}{2}$ bushel; } { Holes dibbled, 6 inches apart in the rows; Manures (mixed with Ashes) put in, and Seed above }	51 $\frac{1}{2}$	53.3	28 $\frac{1}{2}$
6	{ 2 cwt. Superphosphate, 2 cwt. Nitrate Soda. Seed 3 bushels; } { Manures mixed with Ashes and sown broadcast; seed drilled }	56 $\frac{1}{2}$	51.6	32 $\frac{1}{2}$

SECOND SEASON, 1872.

Experiments upon Barley. Thirty-acres Field. Plots $\frac{1}{2}$ acre each.

PLOT No.	MANURES PER ACRE, &c.	PRODUCE PER ACRE.		
		Dressed Corn.		Total Straw.
		Quantity.	Weight per Bushel.	
1	Unmanured. Seed 2 $\frac{1}{2}$ bushels, drilled	Bushels.	lbs.	cwts.
2	{ 3 cwt. Superphosphate, 2 cwt. Nitrate Soda. Seed 2 $\frac{1}{2}$ bushels; } { Manures made up to 15 bushels per acre with Ashes, and sown broadcast; seed drilled .. }			
3	{ 3 cwt. Superphosphate, 2 cwt. Nitrate Soda. Seed 2 $\frac{1}{2}$ bushels; } { The Superphosphate mixed with 40 lbs. slaked Lime to neutralize the acid, the Nitrate added, and the whole made up to 15 bushels per acre with Ashes, and sown broadcast; Seed drilled }			
4	{ 1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 2 $\frac{1}{2}$ bushels; } { Manures and Seed made up to 15 bushels per acre with Ashes, and the whole (Manure, Seed, and Ashes) drilled together }			
5	{ 1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 2 $\frac{1}{2}$ bushels; } { Manures and Seed made up to 15 bushels per acre with a mixture of half Lime and half Ashes, and the whole (Manure, Seed, Lime, and Ashes) drilled together }			