


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1878

[Full Table of Content](#)



Memoranda of the Field Experiments at Rothamsted May 1878

Rothamsted Research


Rothamsted Research (1879) *Memoranda of the Field Experiments at Rothamsted May 1878* ; Yields Of The Field Experiments 1878, pp 1 - 23 - DOI: <https://doi.org/10.23637/ERADOC-1-242>

MEMORANDA
OF THE
ORIGIN, PLAN, AND RESULTS
OF THE
FIELD EXPERIMENTS
CONDUCTED ON THE
FARM OF JOHN BENNET LAWES, Esq.,
AT
ROTHAMSTED, HERTS;
ALSO A STATEMENT OF THE
PRESENT AND PREVIOUS CROPPING, ETC.,
OF THE
ARABLE LAND NOT UNDER EXPERIMENT.

MAY, 1878.

CONTENTS.

	PAGE
Origin, Scope, and Plan of the Rothamsted Experiments	3-5
Lists of Papers published	6-7
Experiments on Permanent Meadow Land; The Park	8
Experiments on Barley; Hoos Field	9
Experiments on Wheat; Broadbalk Field	10
Experiments on Oats; Geescroft Field	11
Experiments on Beans; Geescroft Field	12
Experiments on Clover; Hoos Field	13-14
Experiments on Turnips; Barn Field	15
Experiments on Sugar-beet; Barn Field	16-17
Experiments on Mangold-Wurzel; Barn Field	18
Experiments on Potatoes; Hoos Field	19
Experiments on Rotation; Agdell Field	20
Experiments with different descriptions of Wheat; Foster's Field	21
Cropping, &c., of the Arable Land not under Experiment	22-23



ORIGIN, SCOPE, AND PLAN, OF THE ROTHAMSTED EXPERIMENTS.

THE following statement of the origin, scope, and plan, of the Rothamsted Investigations, was drawn up in answer to a circular letter issued by a Committee appointed to arrange for the commemoration of the twenty-fifth anniversary of the establishment of the First Experimental Station in Germany (Möckern), which was held in Leipzig in September 1877. The precise form of the statement depended on the order and form of the questions to which it is an answer. It has already been published in German, almost in full, with the series of reports of other Experimental Stations, which was issued at the time of the Jubilee Meeting.¹ It has been thought that it would be of some interest as an introduction to the *Memoranda of the Plan and Results of Field Experiments* annually issued at Rothamsted, and which here follow it. To the general statement, which, with a few slight alterations correcting it up to date, is given in the form in which it was originally drawn up, are appended lists of the titles of all the papers already published, with full reference to the Journals in which they appeared.

Mr. Lawes was the founder of the Rothamsted Experimental Station. He commenced experiments with different manuring substances, first with plants in pots, and afterwards in the field, soon after entering into possession of his hereditary property at Rothamsted² in 1834. The researches of De Saussure on vegetation were the chief subjects of his study to this end. Of all the experiments so made, those in which the neutral phosphate of lime, in bones, bone-ash, and apatite, was rendered soluble by means of sulphuric acid, and the mixture applied for root-crops, gave the most striking results. The results obtained on a small scale in 1837, 1838, and 1839, were such as to lead to more extensive trials in the field in 1840 and 1841, and subsequently.

In 1843 more systematic field experiments were commenced, and a barn, which had previously been partially applied to laboratory purposes, became almost exclusively devoted to agricultural investigations. The foundation of the Rothamsted Experimental Station may be said to date from that time (1843).

The Rothamsted station has up to the present time been entirely disconnected from any external organization, and has been maintained entirely by Mr. Lawes. He has further set apart a sum of £100,000, and certain areas of land, for the continuance of the investigations after his death.

In 1854-5 a new laboratory was built, by public subscription of agriculturists, and presented to Mr. Lawes, in July 1855, from which date the old barn-laboratory was abandoned, and the new one has been occupied.

From June 1843, up to the present time, Dr. J. H. Gilbert has been associated with Mr. Lawes, and has had the direction of the laboratory.

The number of assistants and other helps has increased from time to time. At first only one laboratory man was employed; but very soon a chemical assistant was necessary, and next a computer and record-keeper.

During the last twenty-five years the staff has consisted of—
One or two, and sometimes three, chemists.

Two or three general assistants. One of these is usually employed in routine chemical work, but sometimes in more general work. The chief occupation of the general assistants

is to superintend the field experiments—that is, the making of the manures, the measurement of the plots, the application of the manures, and the harvesting of the crops; also the taking of samples, the preparation of them for preservation or analysis, and the determinations of dry matter, ash, &c. These assistants also superintend any experiments made with animals. There are now more than 25,000 bottles of samples of experimentally-grown vegetable produce, of animal products, of ashes, or of soils, stored in the laboratory.

A botanical assistant is also occasionally employed, with from three to six boys under him, and with him is generally associated one of the permanent general assistants, who at other times undertakes the botanical work.

Two or three (for some time past three) computers and record-keepers have been occupied in calculating and tabulating field, feeding, and laboratory results, copying, &c.

One, and sometimes two, laboratory men are employed.

Besides the permanent laboratory staff, chemical assistance is frequently engaged in London, or elsewhere; and in this way, for some years past, Mr. R. Richter, of Berlin, has been almost constantly occupied with analytical work sent from Rothamsted.

The field experiments, and occasionally feeding experiments, also employ a considerable but a very variable number of agricultural labourers.

Nothing has been done at Rothamsted in the way of manure-, feeding-stuff-, or seed-control.

The investigations may be classed under two heads:—

I.—FIELD EXPERIMENTS, EXPERIMENTS ON VEGETATION, &c.

The general scope and plan of the field experiments has been:—

To grow some of the most important crops of rotation, each separately, year after year, for many years in succession on the same land, without manure, with farmyard-manure, and with a great variety of chemical manures; the same description of manure being, as a rule, applied year after year on the same plot. Experiments on an actual course of rotation, with different manures, have also been made. In this way field experiments have been conducted as follows:—

(¹) Die landwirtschaftlichen Versuchs-Stationen Organ. Band xxii. 1877.

(²) Rothamsted is in Hertfordshire, twenty-five miles from London, on the Midland Railway; station, Harpenden.

On Wheat, thirty-five years in succession; 18 acres, 35 plots, many of which are duplicates of others.

On Barley, twenty-seven years in succession; $4\frac{1}{2}$ acres, 23 (or 29) plots.

On Oats, ten years (including one year fallow); $\frac{3}{4}$ acre, 6 plots.

On Wheat, alternated with fallow, twenty-seven years; 1 acre, 2 plots.

On different descriptions of Wheat, eleven years; 4-8 acres (each year in a different field), now more than 20 plots.

On Beans, thirty-two years (including one year Wheat and five years fallow); $1\frac{1}{4}$ acre, 10 plots. Also twenty-seven years; 5 plots, 1 acre.

On Beans, alternated with Wheat, twenty-eight years; 1 acre 10 plots.

On Clover, with fallow or a corn-crop intervening, twenty-eight years; 3 acres, 18 plots.

On Turnips, twenty-eight years (including three years' barley); about 8 acres, 40 plots.

On Sugar Beet, five years; about 8 acres, 40 plots.

On Mangel Wurzel, two years (in progress); about 8 acres, 40 plots.

On Potatoes, two years (in progress); 2 acres, 10 plots.

On rotation, thirty-one years; about $2\frac{1}{2}$ acres, 12 plots.

On permanent grass-land, twenty-three years; about 7 acres, 20 plots.

Comparative experiments with different manures have also been made on other descriptions of soil, in other localities.

Samples of all the experimental crops are taken, and brought to the laboratory. Weighed portions of each are partially dried, and preserved for future reference or analysis. Duplicate weighed portions of each are dried at 100°C , the dry matter determined, and then burnt to ash on platinum sheets in cast-iron muffles. The quantities of ash are determined and recorded, and the ashes themselves are preserved for reference, or analysis.

In a large proportion of the samples the nitrogen is determined.

In selected cases, illustrating the influence of season, manures, exhaustion, &c., complete ash analyses have been made, numbering in all more than 500.

Also in selected cases, illustrating the influence of season and manuring, quantities of the experimentally-grown Wheat grain have been sent to the mill, and the proportion and composition of the different mill-products determined.

In the case of Sugar Beet the sugar, by polariscope, has in most cases been determined.

In the case of the experiments on the mixed herbage of permanent grass land, besides the samples taken for the determination of chemical composition (dry matter, ash, nitrogen, woody fibre, fatty matter, and composition of ash), carefully averaged samples have frequently been taken for the determination of the botanical composition. In this way, on four occasions, at intervals of five years—viz., in 1862, 1867, 1872, and 1877—a sample of the produce of each plot was taken, and submitted to careful botanical separation, and the percentage by weight of each species in the mixed herbage determined. Partial separations have also been made in other years.

INVESTIGATION OF SOILS.

Samples of the soils of most of the experimental plots have been taken from time to time, generally to the depth of 9, 18, and 27 inches, but sometimes to twice this depth. In this way nearly 600 samples have been taken, submitted to partial mechanical separation, and portions of the mould have been carefully prepared and preserved for analysis. In a large proportion of the samples the loss on drying at different temperatures,

and at ignition, has been determined. In most the nitrogen determinable by burning with soda-lime has been estimated. In some the carbon, and in some the nitrogen as nitric acid, have been determined. Some experiments have also been made on the comparative absorptive capacity (for water and ammonia) of different soils and subsoils. The systematic investigation of the amount and condition of the nitrogen, and of some of the more important mineral constituents, of the soils of the different plots, and from different depths, is now in progress or contemplated.

RAINFALL AND DRAINAGE.

Almost from the commencement of the experiments the rainfall has been measured—for twenty-four years in a gauge of one-thousandth of an acre area, as well as in an ordinary small funnel-gauge of 5 inches diameter. From time to time the nitrogen, as ammonia and as nitric acid, has been determined in the rain waters.

Three "drain gauges," also of one-thousandth of an acre each, for the determination of the quantity and composition of the water percolating respectively through 20 inches, 40 inches, and 60 inches depth of soil (with its subsoil in natural state of consolidation) have also been constructed. A more numerous series of smaller "drain gauges," arranged for the investigation of the influence of different crops and of different manures, are in course of construction. Each of the differently manured plots of the permanent experimental Wheat-field having a separate pipe-drain, the drainage-waters have frequently been collected and analysed.

Professor Frankland has determined the nitrogen, as ammonia, as nitric acid, and as organic nitrogen, and also some other constituents, in many samples both of the rain and of the various drainage waters collected at Rothamsted; and Dr. Voelcker has determined the combined nitrogen, and also the incombustible constituents, in many of the drainage waters.

AMOUNT OF WATER TRANSPIRED BY PLANTS.

For several years in succession, experiments were made to determine the amount of water given off by plants during their growth. In this way various plants, including representatives of the gramineous, the leguminous, and other families, have been experimented upon. Similar experiments have also been made with various trees.

BOTANICAL CHARACTERISTICS, &c.

Having regard to the difference in the character and amount of the constituents assimilated by plants of different botanical relationships, under equal external conditions, or by the same description of plants, under varying conditions, observations have been made on the character and range of the roots of different plants, and on their relative development of stem, leaf, &c. In the case of various crops, but more especially with Wheat and Beans, samples have been taken at different stages of growth, and the composition determined, in more or less detail, sometimes of the entire plant, and sometimes of the separated parts. In a few cases the amounts of dry matter, ash, nitrogen, &c., in the above-ground growth of a given area, at different stages of development, have been determined. The amounts of stubble of different crops have also occasionally been estimated.

ASSIMILATION OF FREE NITROGEN.

Experiments were made for several years in succession to determine whether plants assimilate free or uncombined nitrogen, and also various collateral points. Plants of the gramineous, the leguminous, and of other families were operated upon. The late Dr. Pugh took a prominent part in this inquiry.

II.—EXPERIMENTS ON ANIMALS, &c.

Experiments with the animals of the farm were commenced early in 1847, and have been continued, at intervals, up to the present time.

The following points have been investigated :—

1. The amount of food and of its several constituents consumed in relation to a given live weight of animal within a given time.
2. The amount of food and of its several constituents consumed to produce a given amount of increase in live weight.
3. The proportion and relative development of the different organs or parts of different animals.
4. The proximate and ultimate composition of the animals in different conditions as to age and fatness, and the probable composition of their increase in live weight during the fattening process.
5. The composition of the solid and liquid excreta (the manure) in relation to that of the food consumed.
6. The loss or expenditure of constituents by respiration and the cutaneous exhalations—that is, in the mere sustenance of living meat-and-manure-making machine.

The general plan of experimenting was as follows :—

To provide data as to the amount of food or its several constituents consumed in relation a given live weight of animal within a given time, and to produce a given amount of increase in live weight, several hundred animals—oxen, sheep, and pigs—have been experimented upon. Selected lots of animals were supplied, for many weeks, or for months consecutively, with weighed quantities of foods, selected and allotted according to the special point under inquiry. The composition of the foods was determined by analysis. The weights of the animals were taken at the commencement, at intervals during the progress, and at the conclusion of the experiment.

The amount and relative development of the different organs and parts were determined in two calves, two heifers, fourteen bullocks, one lamb, 249 sheep, and fifty-nine pigs.

The percentage of water, mineral matter, fat, and nitrogenous substance, were determined in certain separated parts, and in the entire bodies, of ten animals—namely, one calf, two oxen one lamb, four sheep, and two pigs. Complete analyses of the ashes, respectively, of the entire carcasses, of the mixed internal and other offal parts, and of the entire bodies, of each of these ten animals have also been made.

From the data provided, as just described, as to the chemical composition of the different descriptions of animal, in different conditions as to age and fatness, the composition of the increase whilst fattening, and the relation of the constituents stored up in increase to those consumed in food, have been estimated.

To ascertain the composition of the manure in relation to that of the food consumed, oxen, sheep, and pigs have been experimented upon.

In the case of oxen, the food and litter (sometimes with an acid absorbent), were weighed, sampled, and analysed; the animals were fed in boxes, for periods of from five to nine weeks, and the total dung produced was well mixed, weighed, sampled, and analysed. The constituents determined in the food and litter on the one hand, and in the dung on the other, were dry matter, ash, and nitrogen.

In the case of sheep no litter was used; the animals were kept in lots of five, on rafters, through which (but with some little loss) the solid and liquid excreta passed on to a sheet-zinc flooring at such an incline that the liquid drained off at once into carboys containing acid, and the solid matter was removed two or three times daily, and also mixed with acid. The constituents determined in the food and manure were dry matter, mineral matter, sometimes woody fibre, and nitrogen.

In the case of pigs, individual male animals were experimented upon, each for periods of three, five, or ten days only. Each animal was kept in a frame, preventing it from turning round, and having a zinc bottom, with an outlet for the liquid to run into a bottle, and it was watched night and day, and the voidings carefully collected as soon as passed, which could easily be done, as the animal never passed either fæces or urine without getting up, and in getting up he rang a bell, and so attracted the notice of the attendant. The constituents determined were, in the food and fæces, dry matter, ash and nitrogen, and in the urine, dry matter, ash, nitrogen, and urea.

The loss or expenditure of constituents, by respiration and the cutaneous exhalations, has not been determined directly, that is by means of a respiration-apparatus, but only by difference, that is, by calculation, founded on the amounts of dry matter, ash, and nitrogen, in the food, and in the (increase) fæces, and urine.

Independently of the points of inquiry above enumerated, the results obtained have supplied data for the consideration of the following questions :—

1. The characteristic demands of the animal body (for nitrogenous or non-nitrogenous constituents of food) in the exercise of muscular power.
2. The sources in the food of the fat produced in the animal body.
3. The comparative characters of animal and vegetable food in human dietaries.

SUPPLEMENTARY INVESTIGATIONS.

In conjunction with Professor Way, an extensive investigation was undertaken on the application of town sewage to different crops, but especially to grass. The amount, and the composition, of both the sewage and the produce grown were determined, and in selected cases the composition of the land drainage-water was also determined. Comparative experiments were also made on the feeding qualities of the differently grown produce, the amount of increase yielded by oxen, and the amount and composition of the milk yielded by cows, being determined. In this inquiry part of the analytical work was performed at Rothamsted, but most of it by Professor Way in London.

The chemistry of the malting process, the loss of food constituents during its progress, and the comparative feeding value of barley and malt, have been investigated.

Although many of the results of the investigations above enumerated have already been published, a large proportion as yet remains unpublished.

The following lists give the titles of the papers already published, arranged in two series, and within each series arranged in chronological order; and they show in what Journal each paper appeared.

SERIES I.—REPORTS OF FIELD EXPERIMENTS, EXPERIMENTS ON VEGETATION, &c. PUBLISHED 1847—1878 INCLUSIVE.

1. Agricultural Chemistry (Jour. Roy. Ag. Soc. Eng., vol. viii., p. 226) 1847
2. Agricultural Chemistry, Turnip Culture (Jour. Roy. Ag. Soc. Eng., vol. viii., p. 494) 1847
3. Experimental Investigation into the Amount of Water Given Off by Plants during their Growth, especially in relation to the Fixation and Source of their various Constituents (Jour. Hort. Soc. Lond., vol. v., p. 38) 1850
4. Report of some Experiments undertaken at the suggestion of Professor Lindley, to ascertain the Comparative Evaporating Properties of Evergreen and Deciduous Trees (Jour. Hort. Soc. Lond., vol. vi., p. 227) 1851
5. Agricultural Chemistry, especially in relation to the Mineral Theory of Baron Liebig (Jour. Roy. Ag. Soc. Eng., vol. xii., p. 1) 1851
6. On the Amounts of, and Methods of Estimating, Ammonia and Nitric Acid in Rain-water (Report of the British Association for the Advancement of Science for 1854) 1854
7. Report of the Right Hon. the Earl of Leicester, on the Experiments, conducted by Mr. Keary, on the Growth of Wheat upon the same land for four successive years, at Holkham Park Farm (Jour. Roy. Ag. Soc. Eng., vol. xvi., p. 207) 1855
8. On some points connected with Agricultural Chemistry; being a reply to Baron Liebig's "Principles of Agricultural Chemistry" (Jour. Roy. Ag. Soc. Eng., vol. xvi., p. 411) 1855
9. On the Growth of Wheat by the Lois Weedon System, on the Rothamsted Soil; and on the Combined Nitrogen in Soils (Jour. Roy. Ag. Soc. Eng., vol. xvii., p. 582) 1856
10. On some points in the Composition of Wheat Grain, its Products in the Mill, and Bread (Journal of the Chemical Society of London, vol. x., p. 1) 1857
11. On the Growth of Barley by Different Manures continuously on the Same Land; and on the Position of the Crop in Rotation (Jour. Roy. Ag. Soc. Eng., vol. xviii., p. 454) 1857
12. Report of Experiments with different Manures on Permanent Meadow Land, with Tabular Appendix (Jour. Roy. Ag. Soc. Eng., vols. xix., p. 552, and xx., pp. 228 and 398) 1858-9
13. Report of Experiments on the Growth of Red Clover by different Manures (Jour. Roy. Ag. Soc. Eng., vol. xxi., p. 178) 1860
14. On the Sources of the Nitrogen of Vegetation; with special reference to the question whether Plants Assimilate Free or Uncombined Nitrogen.—Abstract (Proceedings of the Royal Society of London, vol. x., p. 544) 1860
15. On the Application of Different Manures to Different Crops, and on their Proper Distribution on the Farm 1861
16. On some Points in connection with the Exhaustion of Soils.—Abstract (Report of the British Association for the Advancement of Science for 1861) 1861
17. On the Sources of the Nitrogen of Vegetation, with special reference to the question whether Plants Assimilate Free or Uncombined Nitrogen (Philosophical Transactions, part 2, 1861) 1861
18. Report of Experiments made at Rodmersham, Kent, on the Growth of Wheat by different Descriptions of Manure for several years in succession on the same land (Jour. Roy. Ag. Soc. Eng., vol. xxiii., p. 31) .. 1862
19. The Effects of Different Manures on the Mixed Herbage of Grass Land (Jour. Roy. Ag. Soc. Eng., vol. xxiv., p. 131) 1863
20. On the Sources of the Nitrogen of Vegetation, with special reference to the question whether Plants assimilate Free or Uncombined Nitrogen (Jour. Chem. Soc., new series, vol. i.; entire series, vol. xvi.) 1863
21. Liebig and the "Mineral Theory" (note, extracted from a paper by Messrs. Lawes and Gilbert, Jour. Roy. Ag. Soc. Eng., vol. xxiv., part 2) 1863
22. Further Report of Experiments with Different Manures on Permanent Meadow Land (Jour. Roy. Ag. Soc. Eng., vol. xxiv., part 2) 1863
23. Report of Experiments on the Growth of Wheat for Twenty Years in Succession on the same land (Jour. Roy. Ag. Soc. Eng., vol. xxv., parts 1 and 2) 1864
24. On the Selection of Artificial Manures for the Sugar-cane 1864
25. Preliminary Notice of Results on the Composition of Wheat grown for twenty years in succession on the same land (Report of the British Association for the Advancement of Science for 1867) 1867
26. On the Home Produce, Imports, and Consumption of Wheat (Jour. Roy. Ag. Soc. Eng., vol. vi., s.s., part 2) 1868
27. Exhaustion of the Soil in relation to Landlords' Covenants, and the Valuation of Unexhausted Improvements (read before the London Farmers' Club, April 4, 1870) 1870
28. Scientific Agriculture with a view to Profit (read before the Maidstone Farmers' Club, Dec. 15, 1870) 1870
29. Reports of Experiments on the Influence of various Manures on different Species of Plants (Proceedings of the Royal Horticultural Society) 1870
30. Effects of the Drought of 1870 on some of the Experimental Crops at Rothamsted (Jour. Roy. Ag. Soc. Eng., vol. vii., s.s., part 1) 1871
31. Notes on Clover Sickness (Jour. Roy. Hort. Soc., vol. iii.) 1871
32. Report of Experiments on the Growth of Barley for Twenty Years in Succession on the same land (Jour. Roy. Ag. Soc. Eng., vol. ix., s.s., parts 1 and 2) 1873
33. Unexhausted Tillages and Manures, with reference to the Landlord and Tenant (Ireland) Act, 1870 .. 1874
34. On the more frequent Growth of Barley on Heavy Land (read before the London Farmers' Club, Feb. 1, 1875) 1875
35. On the Valuation of Unexhausted Manures (Jour. Roy. Ag. Soc. Eng., vol. xi., s.s., part 1) 1875
36. Note on the Occurrence of "Fairy Rings" (Jour. Linn. Soc., Botany, vol. xv., p. 17) 1875
37. On some points in connection with Vegetation (Address delivered at South Kensington in the Chemical Section of the Science Conferences) 1876
38. On Rainfall, Evaporation, and Percolation (Proceedings of the Inst. of Civil Engineers, vol. xiv., part 3) 1876
39. Freedom in the Growth and Sale of the Crops of the Farm, considered in relation to the interests of the Landowner and the Tenant Farmer (Jour. Soc. Arts, December 14, 1877) 1877
40. On Nitrification; a Report of Experiments made in the Rothamsted Laboratory (Jour. Chem. Soc., January, 1878) 1878

SERIES II.—REPORTS OF EXPERIMENTS ON THE FEEDING OF ANIMALS, SEWAGE UTILISATION, &c.
PUBLISHED 1849—1877, INCLUSIVE.

1. Agricultural Chemistry: Sheep Feeding and Manure, Part I. (With Tabular Appendix in 1856.) (Jour. Roy. Ag. Soc. Eng., vol. x., p. 276) 1849
2. Report of Experiments on the Comparative Fattening Qualities of Different Breeds of Sheep; Hampshire and Sussex Downs (Jour. Roy. Ag. Soc. Eng., vol. xii., p. 414) 1851
3. Report of Experiments on the Comparative Fattening Qualities of Different Breeds of Sheep—Cotswolds (Jour. Roy. Ag. Soc. Eng., vol. xiii., p. 179) 1852
4. On the Composition of Foods in relation to Respiration and the Feeding of Animals (Report of the British Association for the Advancement of Science for 1852) 1852
5. Agricultural Chemistry: Pig Feeding (Jour. Roy. Ag. Soc. Eng., vol. xiv., p. 459) 1853
6. On the Equivalency of Starch and Sugar in Food (Report of the British Association for the Advancement of Science for 1854) 1854
7. Experiments on the Comparative Fattening Qualities of Different Breeds of Sheep—Leicesters and Cross-breeds (Jour. Roy. Ag. Soc. Eng., vol. xvi., p. 45) 1855
8. On the Sewage of London (Journal of the Society of Arts, March 7, 1855) 1855
9. Letter on the Utilisation of Town Sewage (from the Report ordered by the House of Commons to be printed, Aug. 3, 1857. Appendix xii., p. 477) 1857
10. Experimental Inquiry into the Composition of some of the Animals Fed and Slaughtered as Human Food. Abstract (Proceedings of the Royal Society of London, vol. ix., p. 348) 1858
11. Observations on the recently-introduced Manufactured Foods for Agricultural Stock (Jour. Roy. Ag. Soc. Eng., vol. xix., p. 199) 1858
12. Experimental Inquiry into the Composition of some of the Animals Fed and Slaughtered as Human Food (Philosophical Transactions, Part 2, 1859) 1859
13. On the Composition of Oxen, Sheep, and Pigs, and of their Increase while Fattening (Jour. Roy. Ag. Soc. Eng., vol. xxi., p. 433) 1860
14. On the Composition of the Animal Portion of our Food, and on its relations to Bread—Abstract (Jour. Chem. Soc., vol. xii., p. 54) 1860
15. Fifth Report of Experiments on the Feeding of Sheep (Jour. Roy. Ag. Soc. Eng., vol. xxii., p. 189) 1861
16. Report of Experiments on the Fattening of Oxen at Woburn Park Farm (Jour. Roy. Ag. Soc. Eng., vol. xxii., p. 200) 1861
17. Experiments on the Question whether the Use of Condiments increases the Assimilation of Food by Fattening Animals, or adds to the Profits of the Feeder (Edinburgh Veterinary Review and Annals of Comparative Pathology, July, 1862) 1862
18. Supplementary Report of Experiments on the Feeding of Sheep (Jour. Roy. Ag. Soc. Eng., vol. xxiii., p. 191) 1862
19. The Utilisation of Town Sewage (Jour. Roy. Ag. Soc. Eng., vol. xxiv., p. 65) 1863
20. On the Chemistry of the Feeding of Animals for the Production of Meat and Manure (read before the Royal Dublin Society, March 31, 1864) 1864
21. On the Sewage of Towns (Third Report and Appendices 1, 2, and 3, of the Royal Commission. Presented to Parliament) 1865
22. Report (presented to Parliament) of Experiments undertaken by Order of the Board of Trade to Determine the Relative Values of Unmalted and Malted Barley as Food for Stock 1866
23. On the Composition, Value, and Utilisation of Town Sewage (Jour. Chem. Soc., New Series, vol. iv.; Entire Series, vol. xix.) 1866
24. Food, in its Relations to the various Exigencies of the Animal Body (Phil. Mag., July, 1866) 1866
25. On the Sources of the Fat of the Animal Body (Phil. Mag., December, 1866) 1866
26. Note—On Sewage Utilisation (Proceedings of the Institution of Civil Engineers, vol. xiv., Part 3) 1876
27. On some Points in connection with Animal Nutrition (Address delivered at South Kensington in the Biological Section of the Science Conferences) 1876
28. On the Formation of Fat in the Animal Body (Journal of Anatomy and Physiology, vol. xi., Part 4) 1877

THE PARK.

EXPERIMENTS WITH DIFFERENT MANURES ON PERMANENT MEADOW LAND.

The Land has probably been laid down with Grass for some centuries. No fresh seed has been artificially sown within the last 40 years certainly; nor is there record of any having been sown since the Grass was first laid down. The experiments commenced in 1856, at which time the character of the herbage appeared uniform over all the Plots. Excepting as explained in the Table, and in the foot-notes, the same description of Manure has been applied year after year to the same Plot.

During the first 19 years of the experiments, 1856-1874, the first crop only, each year, was mown, made into hay, removed from the land, and weighed. As a rule, the second crop was fed-off by sheep having no other food, the object being not to disturb the condition of the manuring. A given number was allotted to each Plot, according to the amount of produce, penned upon a portion of it, and the area extended, day by day, until the whole was eaten down. Frequently, however, the animals suffered considerably; and in 1866, 1870, 1873, and 1874, the second crops (and third, if any) were cut, and spread on the respective Plots. In the twentieth season, 1875, the second crops being unusually heavy, and the weather favourable, they were, for the first time, cut, weighed as hay, and removed. In 1876 they were cut and spread on the Plots. In 1877 the second crops were again made into hay, weighed, and removed; and it is intended, in future, to adopt this plan, whenever the weather will permit.

(Area under experiment, about 7 acres.)

PLOTS.		PRODUCE PER ACRE, WEIGHED AS HAY.										PLOTS.		
		Average per Annum.					Twenty-first Season, 1876.		Twenty-second Season, 1877.					
		10 Years, 1866-75.		20 Years, 1846-75.		Cwts.	First Crop.	Second Crop.	Total.	Cwts.	First Crop.		Second Crop.	Total.
		Cwts.	Cwts.	Cwts.	Cwts.									
	1 acre.. .. = (about) 0-40 Hectare or 1-39 Prussian Morgen. 1 lb. (pound avoird.) .. = (about) 5-10 Kilogrammes or 0-91 Zollverein Pfund. 1 cwt. (hundredweight) .. = (about) 50-56 Kilogrammes or 20-33 Cantars. 1 ton = (about) 1016-0 Kilogrammes or 20-33 Cantars. 1 lb. per acre = (about) 1-12 Kilogrammes per Hectare or 0-57 Zollv. Pfd. per Pr. Morgen. 1 cwt. per acre = (about) 125-5 Kilogrammes per Hectare or 0-84 Cantar per Pr. Morgen. 1 ton per acre = (about) 2510-0 Kilogrammes per Hectare or 12-82 Cantar per Pr. Morgen.	Manures, per acre, per Annum.												
	(1856-63, 8 years, 14 tons Farmyard Manure, and 200 lbs. Ammonia-salts (a); average produce 49½ cwt. { 1864 and since, 200 lbs. Ammonia-salts alone; average produce (12 years, 1864-75) 38½ cwt. }	Cwts. 48½	Cwts. 37½	Cwts. 43	Cwts. 29½	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	1
	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwt. { 1864 and since, unmanured; average produce (12 years, 1864-75) 32½ cwt. }	41½	32	36½	20½	..	..	..	..	..	..	..	..	2
	Unmanured, continuously	22½	20	21½	12½	..	..	..	..	..	..	..	..	3
	3½ cwt. Superphosphate of Lime (a)	23½	21½	22½	16½	..	..	..	..	..	..	..	..	4
	3½ cwt. Superphosphate of Lime, and 400 lbs. Ammonia-salts	33½	30½	32½	18½	..	..	..	..	..	..	..	..	4 1/2
	400 lbs. Ammonia-salts	30½	22	26½	17½	..	..	..	..	..	..	..	..	5
	(1856-68, 13 years, 400 lbs. Ammonia-salts; average produce 30½ cwt. { 1869 and since, 300 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwt. Superphosphate; av. prod. (7 yrs., 1869-75) 31½ cwt. }	31½	30½	30½	32	..	..	..	..	..	..	..	..	6
	300 lbs. Sulphate Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulphate Magnesia, and 3½ cwt. Superphosphate	33½	36½	35½	34½	..	..	..	..	..	..	..	..	7
	(1856-61, 6 years, 300 lbs. Sulph. Potash, 200 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwt. Superphosphate; average produce 36 cwt. { 1862 and since, 250 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulphate Magnesia, and 3½ cwt. Superphosphate; average produce (14 years, 1862-75) 27½ cwt. }	33½	26½	30½	24½	..	..	..	..	..	..	..	..	8
	300 lbs. Sulphate Potash, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, 3½ cwt. Superphosphate, and 400 lbs. Ammonia-salts	53½	48½	51	50	..	..	..	..	..	..	..	..	9
	(1856-61, 6 yrs. 300 lbs. Sulph. Potash, 200 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwt. Superphosphate, 400 lbs. Ammonia-salts; av. prod. 55½ cwt. { 1862 and since, 250 lbs. Sulph. Potash, 100 lbs. Sulph. Magnesia, 3½ cwt. Superphosphate, 400 lbs. Ammonia-salts; av. prod. (14 yrs., 1862-75) 42½ cwt. }	52½	39½	46½	40	..	..	..	..	..	..	..	..	10
	300 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwt. Superphosphate, 800 lbs. Ammonia-salts	61½	53½	57½	57½	..	..	..	..	..	..	..	..	11
	300 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwt. Superphosphate, 800 lbs. Ammonia-salts	63½	61½	62½	64½	..	..	..	..	..	..	..	..	11 1/2
	Unmanured continuously	25	22½	24	14½	..	..	..	..	..	..	..	..	12
	300 lbs. Sulphate Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwt. Superphosphate, 400 lbs. Ammonia-salts, and 400 lbs. Cut Wheat-straw	55½	59½	57	64½	..	..	..	..	..	..	..	..	13
	550 lbs. Nitrate of Soda (a), 300 lbs. Sulphate Potash, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwt. Superphosphate	53½	60½	57	64½	..	..	..	..	..	..	..	..	14
	1858-75, 18 years, 550 lbs. Nitrate Soda	36½	35	33½	30½	..	..	..	..	..	..	..	..	15
	1876, and since, 300 lbs. Sulphate Potash, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwt. Superphosphate	45½	47½	46½	41½	..	..	..	..	..	..	..	..	16
	275 lbs. Nitrate of Soda, 300 lbs. Sulphate Potash, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwt. Superphosphate	34½	33½	33½	25½	..	..	..	..	..	..	..	..	17
	275 lbs. Nitrate of Soda	21	33½	32½	37	..	..	..	..	..	..	..	..	18
	Mixture supplying the quantity of Potash, Soda, Lime, Magnesia, Phosphoric acid, Silica, and Nitrogen, contained in 1 ton of Hay (commencing 1865)	..	..	..	..	..	..	..	..	..	..	..	..	19
	275 lbs. Nitrate of Soda, 290 lbs. Sulphate of Potash, and 3½ cwt. Superphosphate (commencing 1872)	..	..	..	..	..	..	..	..	..	..	..	..	19
	275 lbs. Nitrate of Potash, and 3½ cwt. Superphosphate (commencing 1872)	..	..	..	..	..	..	..	..	..	..	..	..	20

(a) "Ammonia-salts"—in all cases equal parts Sulphate and Muriate of Ammonia of Commerce.
 (b) The "Superphosphate of Lime" is, in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid Sp. gr. 1.7 (and water).
 (c) Plots 6, 8, and 10, had, besides the Manures specified, 2000 lbs. Sawdust per acre per annum for the first 7 years, 1856-1862, but without effect.
 (d) 200 lbs. 1856-63 inclusive.
 (e) 500 lbs. in 1862 and 1863.
 (f) Only 400 lbs. in 1869-70-71.
 (g) The application of Silicates did not commence until 1862; 9 years (1862-1870), 200 lbs. Silicate years the first crop only was weighed and removed.

HOOS FIELD.

EXPERIMENTS ON THE GROWTH OF BARLEY YEAR AFTER YEAR ON THE SAME LAND, WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

Previous Cropping—1847, Swedish Turnips, with Dung and Superphosphate of Lime, the Roots carted off; 1848, Barley; 1849, Clover; 1850, Wheat; 1851, Barley manured with Ammonia-salts. First Experimental Barley Crop in 1852. Barley every year since; and, unless stated to the contrary in the Table, or in the foot-notes, the same Manure has been applied year after year to the same Plot.

(Area under experiment, about 44 acres.)

Plots.	PRODUCE PER ACRE.												Plots.
	Average per Annum.												
	Dressed Corn.												
	Total Straw.												
	Dressed Corn.												
Manures, per acre, per annum.													
Unmanured continuously													
1 O.	1 acre	0-40 Hectare	..	..	..	or 1-59 Prussian Morgen.	12 Years, 1862-63.	24 Years, 1862-76.	12 Years, 1862-63.	24 Years, 1862-76.	12 Years, 1862-63.	24 Years, 1862-76.	
2 O.	1 bushel	0-36 Hectolitre	..	..	..	or 0-66 Prussian Scheffel.	Bushels.	Bushels.	Cwts.	Cwts.	Bushels.	Cwts.	
3 O.	1 lb. (pound avoird.)	0-45 Kilogramme	..	..	..	or 0-91 Zollverein Pfund.	21½	52½	15½	38½	52½	79½	
4 O.	1 cwt. (hundredweight)	51-0 Kilogrammes	..	..	..	or 1-02 Centner.	208	531	144	107	231	591	
1 A.	1 bushel per acre	0-9 Hectolitre per Hectare	..	..	..	or 0-42 Pr. Scheffel per Pr. Morgen.	17½	52½	18½	9½	201	52½	
2 A.	1 lb. per acre	1-12 Kilogrammes per Hectare	..	..	..	or 0-57 Zollv. Pfd. per Pr. Morgen.	30	52½	15½	10½	231	53½	
3 A.	1 cwt. per acre	125-5 Kilogrammes per Hectare	..	..	..	or 0-84 Centner per Pr. Morgen.	21½	52½	15½	10½	231	53½	
1 AA.	200 lbs. Ammonia-salts ^(a)	..	..	..	..	..	34½	53½	19½	15½	35½	15½	
2 AA.	200 lbs. Ammonia-salts, and 3½ cwt. Superphosphate	..	..	..	..	..	47½	53½	24½	20½	54½	19½	
3 AA.	200 lbs. Ammonia-salts, 200 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia	..	..	..	..	..	36½	53½	22½	17½	53½	19½	
4 A.	200 lbs. Ammonia-salts, 200 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag., 3½ cwt. Superphos.	..	..	..	..	..	47½	54½	28½	26½	56½	24½	
1 AAS.	275 lbs. Nitrate Soda	..	..	..	..	..	39½	53½	18½	21½	37½	19½	
2 AAS.	275 lbs. Nitrate Soda, and 3½ cwt. Superphosphate	..	..	..	..	..	50½	53½	24½	23½	46½	21½	
3 AAS.	275 lbs. Nitrate Soda, 200 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag., 3½ cwt. Superphos.	..	..	..	..	..	39½	53½	20½	23½	46½	21½	
4 AAS.	275 lbs. Nitrate Soda, 200 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag., 3½ cwt. Superphos.	..	..	..	..	..	50½	53½	28½	31½	56½	30	
1 C.	1000 lbs. Rape-cake	..	..	..	..	..	47	53½	22½	26½	44½	20½	
2 C.	1000 lbs. Rape-cake, and 3½ cwt. Superphosphate	..	..	..	..	..	48½	53½	24½	27½	44½	20½	
3 C.	1000 lbs. Rape-cake, 200 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	..	..	..	..	..	44½	53½	24½	27½	44½	20½	
4 C.	1000 lbs. Rape-cake, 200 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	..	..	..	..	..	44½	53½	24½	27½	44½	20½	
1 N.	275 lbs. Nitrate of Soda	..	..	..	..	..	38½	53½	20½	23½	44½	20½	
2 N.	275 lbs. Nitrate of Soda	..	..	..	..	..	43½	53½	24½	27½	44½	20½	
3 N.	275 lbs. Nitrate of Potash, 3½ cwt. Superphosphate ^(a)	..	..	..	..	..	24½	53½	24½	27½	44½	20½	
4 N.	275 lbs. Nitrate of Potash, 3½ cwt. Superphosphate ^(a)	..	..	..	..	..	24½	53½	24½	27½	44½	20½	
5 O.	200 lbs. Sulphate of Potash, 3½ cwt. Superphosphate ^(a)	..	..	..	..	..	24½	53½	24½	27½	44½	20½	
5 A.	200 lbs. Sulphate of Potash, 3½ cwt. Superphosphate ^(a)	..	..	..	..	..	24½	53½	24½	27½	44½	20½	
5 M.	200 lbs. Sulphate of Potash, 3½ cwt. Superphosphate ^(a)	..	..	..	..	..	24½	53½	24½	27½	44½	20½	
6 ⁽¹⁾	Unmanured continuously	..	..	..	..	..	24½	53½	24½	27½	44½	20½	
7 ⁽¹⁾	Ashea (burnt soil and turf)	..	..	..	..	..	24	53½	24	27½	44½	20½	
7 ⁽²⁾	Farmyard Manure 14 tons, 20 yrs., 1852-71, av. prod. 48½ bush.; unmanured since, av. prod., 6 yrs., 1872-7, 38½ bush.	..	..	..	..	..	49½	54½	28½	27½	44½	20½	
	Farmyard Manure 14 tons, every year; av. produce, 20 years, 1852-71, 48½ bush.; 6 years, 1872-7, 50 bush.	..	..	..	..	..	46½	54½	29½	29½	44½	20½	

(¹) The "Superphosphate of Lime" is, in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid sp. gr. 1.7 (and water).

(²) 300 lbs. per annum for the first six years, 1852-7.

(³) 300 lbs. per annum for the first six years, 1852-7.

(⁴) The "Ammonia-salts"—in all cases equal parts Sulphate and Murate of Ammonia of Commerce.

(⁵) First 6 years, 1852-7, instead of Nitrate of Soda, 400 lbs. Ammonia-salts per annum; next 10 years, 1858-67, 200 lbs. Ammonia-salts per annum; 1868, and since, 275 lbs. Nitrate of Soda per annum, 275 lbs. Nitrate of Soda is reckoned to contain the same amount of Nitrogen as 200 lbs. "Ammonia-salts."

(⁶) The application of Silicates did not commence until 1864; in 1864-5-6 and 7, 200 lbs. Silicate of Soda and 200 lbs. Silicate of Lime were applied per acre, but in 1868, and since, 400 lbs. Silicate of Soda, and no Silicate of Lime. These plots ("AAS") comprise, respectively, one half of the original "AA" plots, and,

excepting the addition of the Silicates, have been, and are, in other respects, manured in the same way as the

AA plots.

(⁷) 300 lbs. Rape-cake per annum for the first six years, and 1000 lbs. only, each year since.

(⁸) 300 lbs. Silicate of Potash, and 3½ cwt. Superphosphate of Lime, without Nitrate of Soda, the first year (1852); Nitrate of Soda for 1853-4-5-6, and 7; and 275 lbs. only, each year since.

(⁹) 550 lbs. Nitrate of Soda for 1853-4-5-6, and 7; and 275 lbs. only, each year since.

(¹⁰) Ammonia-salts also the first year, but not since.

(¹¹) Averages of 11 years, 12 years, and 23 years.

(¹²) Averages of 6 years, 12 years, and 18 years.

(¹³) Averages of 20 years (with dung), 4 years (unmanured), and 24 years.

BROADBALK FIELD.

EXPERIMENTS ON THE GROWTH OF WHEAT YEAR AFTER YEAR ON THE SAME LAND; WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

Previous Cropping—1839, Turnips, with Farmyard Manure; 1840, Barley; 1841, Peas; 1842, Wheat; 1843, Oats; the last four crops Unmanured. First Experimental Wheat Crop in 1844. Wheat every year since; and, with some exceptions, nearly the same description of Manure on the same Plots each year—especially during the last 27 years (1852 and since). From the commencement of the experiments in 1843-4 up to 1876-7 inclusive, the mineral manures, the ammonia-salts, and rape-cake, &c., if any, were sown in the autumn, before the seed; excepting, in 1845, when, owing to the wet autumn and winter, all the manures were spring-sown; and for the crops of 1873, 4, 5, 6, and 7, the ammonia-salts applied to Plot 15 were top-dressed in the spring. Nitrate of soda has, however, always been sown in the spring. But, in consequence of the ascertained great loss of the nitrogen of the manures by drainage, especially in wet winters, it has been decided to apply only the mineral manures (and Farmyard-manure) in the autumn, and the ammonia-salts, as well as the nitrate, in the spring; excepting on Plot 15, where, for comparison, the ammonia-salts are to be sown in the autumn; and on Plot 19, where the ammonia-salts are mixed in, when the superphosphate is made. This plan is adopted for the first time for the present crop, 1877-8.

(Area under experiment, about 13 acres.)

Plots.	Manures, per acre, per annum.	PRODUCE PER ACRE.												Thirty-Fourth Season, 1877.		Plots.	
		Average per Annum.						Total Straw.						Dressed Corn.	Weight of Straw.		
		Dressed Corn.			Weight per Bushel.			Total Straw.			Weight per Bushel.						
		12 Years, 1862-75.	24 Years, 1864-76.	Quantity.	12 Years, 1862-75.	24 Years, 1864-76.	Quantity.	12 Years, 1862-75.	24 Years, 1864-76.	Quantity.	12 Years, 1862-75.	24 Years, 1864-76.	Quantity.	12 Years, 1862-75.	24 Years, 1864-76.		Quantity.
0	Superphosphate of Lime (three times as much as on No. 5 and succeeding Plots)	Bushels. 18½	Bushels. 17½	Bushels. 12½	Bushels. 57½	Bushels. 58½	Bushels. 16½	Bushels. 13	Bushels. 12	Bushels. 12	Bushels. 12	Bushels. 12	Bushels. 12	Bushels. 12	Cwts. 8½		
1	Sulphates of Potash, Soda, and Magnesia (twice as much as on No. 5 and succeeding Plots)	Bushels. 16½	Bushels. 14½	Bushels. 12½	Bushels. 57½	Bushels. 58½	Bushels. 16½	Bushels. 13	Bushels. 12	Bushels. 12	Bushels. 12	Bushels. 12	Bushels. 12	Bushels. 12	Cwts. 8½		
2	Farmyard Manure (14 tons every year)	Bushels. 35½	Bushels. 36½	Bushels. 14	Bushels. 59½	Bushels. 60	Bushels. 34½	Bushels. 33½	Bushels. 12½	Bushels. 12½	Bushels. 12½	Bushels. 12½	Bushels. 12½	Bushels. 12½	Cwts. 8½		
3	Unmanured continuously	Bushels. 15½	Bushels. 17	Bushels. 13½	Bushels. 56½	Bushels. 59½	Bushels. 15½	Bushels. 13½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Cwts. 8½		
4	Unmanured for Crop of 1852, and since; previously Superphosphate (made with Muratic Acid), and Sulphate Ammonia	Bushels. 17	Bushels. 18½	Bushels. 13½	Bushels. 57½	Bushels. 59½	Bushels. 16½	Bushels. 14½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Cwts. 8½		
5 (a and b)	200 lbs. 0 Sulphate Potash, 100 lbs. 0 Sulphate Soda, 100 lbs. 0 Sulphate Magnesia, 3½ cwt. Superphos., 200 lbs. Ammonia-salts 0	Bushels. 28½	Bushels. 23½	Bushels. 25½	Bushels. 58½	Bushels. 60½	Bushels. 26½	Bushels. 20½	Bushels. 14½	Bushels. 14½	Bushels. 14½	Bushels. 14½	Bushels. 14½	Bushels. 14½	Cwts. 8½		
6 (a and b)	200 lbs. 0 Sulphate Potash, 100 lbs. 0 Sulphate Soda, 100 lbs. 0 Sulphate Mag., 3½ cwt. Superphos., 400 lbs. Ammonia-salts 0	Bushels. 36½	Bushels. 32½	Bushels. 34½	Bushels. 58½	Bushels. 60½	Bushels. 34½	Bushels. 31½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Cwts. 8½		
7 (a and b)	200 lbs. 0 Sulphate Potash, 100 lbs. 0 Sulphate Soda, 100 lbs. 0 Sulphate Mag., 3½ cwt. Superphos., 600 lbs. Ammonia-salts 0	Bushels. 38½	Bushels. 37½	Bushels. 37½	Bushels. 57½	Bushels. 60½	Bushels. 40½	Bushels. 41½	Bushels. 24½	Bushels. 24½	Bushels. 24½	Bushels. 24½	Bushels. 24½	Bushels. 24½	Cwts. 8½		
8 (a and b)	200 lbs. 0 Sulphate Potash, 100 lbs. 0 Sulphate Soda, 100 lbs. 0 Sulphate Mag., 3½ cwt. Superphos., 550 lbs. Nitrate Soda 0	Bushels. 34½	Bushels. 32½	Bushels. 34½	Bushels. 57½	Bushels. 59½	Bushels. 33½	Bushels. 34½	Bushels. 40½	Bushels. 40½	Bushels. 40½	Bushels. 40½	Bushels. 40½	Bushels. 40½	Cwts. 8½		
9 {a	550 lbs. Nitrate of Soda 0. (The Nitrate for both 9a and 9b always sown in the Spring)	Bushels. 23½	Bushels. 21½	Bushels. 23½	Bushels. 55½	Bushels. 58½	Bushels. 22½	Bushels. 18½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Cwts. 8½		
10 {b	400 lbs. Ammonia-salts alone, for 1845, and each year since; Mineral Manure in 1844	Bushels. 27	Bushels. 23	Bushels. 25	Bushels. 57½	Bushels. 58½	Bushels. 25½	Bushels. 20½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Cwts. 8½		
11 (a and b)	400 lbs. Ammonia-salts, 3½ cwt. Superphosphate	Bushels. 29½	Bushels. 25½	Bushels. 27½	Bushels. 56½	Bushels. 59½	Bushels. 28½	Bushels. 23½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Bushels. 17½	Cwts. 8½		
12 (a and b)	400 lbs. Ammonia-salts, 3½ cwt. Superphosphate, and 366½ lbs. 0 Sulphate of Soda	Bushels. 35½	Bushels. 31½	Bushels. 33½	Bushels. 58½	Bushels. 60½	Bushels. 33½	Bushels. 30½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Cwts. 8½		
13 (a and b)	400 lbs. Ammonia-salts, 3½ cwt. Superphosphate, and 200 lbs. 0 Sulphate of Potash	Bushels. 34½	Bushels. 32½	Bushels. 33½	Bushels. 58½	Bushels. 60½	Bushels. 33½	Bushels. 30½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Cwts. 8½		
14 (a and b)	400 lbs. Ammonia-salts, 3½ cwt. Superphosphate, and 280 lbs. 0 Sulphate of Magnesia	Bushels. 35½	Bushels. 31½	Bushels. 33½	Bushels. 58½	Bushels. 60½	Bushels. 33½	Bushels. 30½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Cwts. 8½		
15 {a	400 lbs. 0 Sulph. Pot., 100 lbs. 0 Sulph. Sod., 100 lbs. 0 Sulph. Mag., 3½ cwt. Superphos., 0; 400 lbs. Amm.-salts, in Autm. 0	Bushels. 33½	Bushels. 31½	Bushels. 32½	Bushels. 58½	Bushels. 60½	Bushels. 32½	Bushels. 31½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Cwts. 8½		
15 {b	200 lbs. 0 Sulph. Pot., 100 lbs. 0 Sulph. Sod., 100 lbs. 0 Sulph. Mag., 3½ cwt. Superphos., 0; 400 lbs. Amm.-salts, in Autm. 0	Bushels. 34½	Bushels. 32½	Bushels. 33½	Bushels. 58½	Bushels. 60½	Bushels. 33½	Bushels. 31½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Bushels. 18½	Cwts. 8½		
16 (a and b)	1852-64, 13 years, 200 lbs. Sulph. Potash, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag., 3½ cwt. Superphos., and 800 lbs. Ammonia-salts; average produce 39½ bush. Corn, 4½ cwt. Straw	Bushels. 38½	Bushels. 19½	Bushels. 29	Bushels. 57½	Bushels. 60½	Bushels. 46	Bushels. 31½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Bushels. 9½	Cwts. 8½		
17 (a and b)	1865 and since, unmanured; average produce (12 years, 1865-76) 16½ bushels Corn, 1½ cwt. Straw	Bushels. 32½	Bushels. 28½	Bushels. 30½	Bushels. 58½	Bushels. 58½	Bushels. 33½	Bushels. 27½	Bushels. 10	Bushels. 10	Bushels. 10	Bushels. 10	Bushels. 10	Bushels. 10	Cwts. 8½		
18 (a and b)	400 lbs. Ammonia-salts, 3½ cwt. Superphosphate, and 280 lbs. 0 Sulphate of Magnesia, and 3½ cwt. Superphosphate	Bushels. 31½	Bushels. 28½	Bushels. 30½	Bushels. 58½	Bushels. 60½	Bushels. 31½	Bushels. 28½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Bushels. 19½	Cwts. 8½		
19	3½ cwt. Superphosphate of Lime 0, 300 lbs. Sulphate of Ammonia, and 500 lbs. Rape-cake, in Autm.	Bushels. 15½	Bushels. 13½	Bushels. 13½	Bushels. 57½	Bushels. 59½	Bushels. 15½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Cwts. 8½		
20	Unmanured continuously	Bushels. 22½	Bushels. 19½	Bushels. 20½	Bushels. 57½	Bushels. 59½	Bushels. 20½	Bushels. 16½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Cwts. 8½		
21	200 lbs. 0 Sulph. Potash, 100 lbs. 0 Sulph. Soda, 100 lbs. Sulph. Mag., 3½ cwt. Superphos., 100 lbs. Muratic Ammonia	Bushels. 21½	Bushels. 19½	Bushels. 20½	Bushels. 57½	Bushels. 59½	Bushels. 20½	Bushels. 16½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Cwts. 8½		
22	200 lbs. 0 Sulph. Potash, 100 lbs. 0 Sulph. Soda, 100 lbs. Sulph. Mag., 3½ cwt. Superphos., 100 lbs. Sulphate Ammonia	Bushels. 21½	Bushels. 19½	Bushels. 20½	Bushels. 57½	Bushels. 59½	Bushels. 20½	Bushels. 16½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Bushels. 11½	Cwts. 8½		

- (1) 800 lbs. per annum for Crop of 1858, and previously.
 (2) 200 lbs. per annum for Crop of 1858, and previously.
 (3) a Superphosphate of Lime—in all cases, excepting for Plot 19, made from 200 lbs. Bone-ash, 150 lbs. Sulphate of lime sp. 17 (and water).
 (4) The Ammonia-salts, in all cases, equal parts Sulphate and Muric Acid of Ammonia of Commerce.
 (5) 34 1/2 lbs. Nitrate of Soda in 1852, 27 1/2 lbs. in 1853, and 1854, 550 lbs. each year since. In 1852, 550 lbs. Potash, or Magnesia, or Superphosphate, in 1852, 1853, or 1854, 96, 475 lbs. Nitrate in 1852, 550 lbs. each year since. 550 lbs. Nitrate is reckoned to contain the same amount of Nitrogen as 400 lbs. "Ammonia-salts."
 (6) For 1858, and previously—1 1/2 times as much.
 (7) For 1872 and previously, made with Muric Acid instead of Sulphuric Acid.
 (8) For 1872 and previously, 400 lbs. Sulphate Ammonia, sown in the Autumn; for 1873, 4, 5, 6, and 7, 400 lbs. Ammonia-salts, sown in the Spring; for 1878, 400 lbs. Ammonia-salts, sown in the Autumn.
 (9) For 1872 and previously, 300 lbs. Sulphate Ammonia and 500 lbs. Rape-cake, sown in the Autumn; for 1873, 4, 5, 6, and 7, 400 lbs. Ammonia-salts, sown in the Spring; for 1878, 400 lbs. Ammonia-salts, sown in the Autumn.
 (10) For 1872 and previously, 300 lbs. Sulphate Ammonia and 500 lbs. Rape-cake, sown in the Autumn; for 1873, 4, 5, 6, and 7, 400 lbs. Ammonia-salts, sown in the Spring; for 1878, 400 lbs. Ammonia-salts, sown in the Autumn.

(11)

GEESCROFT FIELD.

EXPERIMENTS ON THE GROWTH OF OATS YEAR AFTER YEAR ON THE SAME LAND; WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

Previous Cropping—1847 and 1848, Clover, Experimental Manures; 1849—1865, Beans, Experimental Manures; 1860, Fallow; 1861 and 1862, Wheat, Unmanured; 1863, Fallow; 1864, Beans, Dunged; 1865, Wheat, Unmanured; 1866, Beans, Unmanured; 1867 and 1868, Wheat, Unmanured.

First Experimental Oat Crop in 1869.

(Area under Experiment, $\frac{1}{2}$ acre.)

PLOTS.	MANURES, PER ACRE, PER ANNUM.	PRODUCE PER ACRE.												AVERAGE PER ANNUM 5 YEARS, 1869-1873.		
		1ST SEASON, 1869.			2ND SEASON, 1870.			3RD SEASON, 1871.			4TH SEASON, 1872.			5TH SEASON, 1873.		
		Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.
		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.	
1	Unmanured	Bushels. 36 $\frac{1}{2}$	lbs. 36 $\frac{1}{2}$	cwts. 19 $\frac{1}{2}$	Bushels. 16 $\frac{1}{2}$	lbs. 35	cwts. 9 $\frac{1}{2}$	Bushels. 20 $\frac{1}{2}$	lbs. 33 $\frac{1}{2}$	cwts. 11 $\frac{1}{2}$	Bushels. 15	lbs. 36 $\frac{1}{2}$	cwts. 7 $\frac{1}{2}$	Bushels. 19 $\frac{1}{2}$	lbs. 33 $\frac{1}{2}$	cwts. 10 $\frac{1}{2}$
2	{ 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate of Lime (1) }	45	38 $\frac{1}{2}$	24 $\frac{1}{2}$	19 $\frac{1}{2}$	35 $\frac{1}{2}$	9 $\frac{1}{2}$	22	35 $\frac{1}{2}$	13 $\frac{1}{2}$	19 $\frac{1}{2}$	37 $\frac{1}{2}$	10 $\frac{1}{2}$	24 $\frac{1}{2}$	35	13 $\frac{1}{2}$
3	400 lbs. Ammonia-salts (2)	56 $\frac{1}{2}$	37 $\frac{1}{2}$	36 $\frac{1}{2}$	30	34 $\frac{1}{2}$	17 $\frac{1}{2}$	57 $\frac{1}{2}$	36 $\frac{1}{2}$	40 $\frac{1}{2}$	55 $\frac{1}{2}$	37 $\frac{1}{2}$	30 $\frac{1}{2}$	47	35 $\frac{1}{2}$	28 $\frac{1}{2}$
4	{ 400 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate .. }	75 $\frac{1}{2}$	39 $\frac{1}{2}$	54	50 $\frac{1}{2}$	36	28 $\frac{1}{2}$	58 $\frac{1}{2}$	35 $\frac{1}{2}$	50	62 $\frac{1}{2}$	39 $\frac{1}{2}$	45 $\frac{1}{2}$	48 $\frac{1}{2}$	37	41 $\frac{1}{2}$
5	550 lbs. Nitrate of Soda (3)	62 $\frac{1}{2}$	38 $\frac{1}{2}$	42 $\frac{1}{2}$	36 $\frac{1}{2}$	35 $\frac{1}{2}$	23	55	36 $\frac{1}{2}$	34 $\frac{1}{2}$	42 $\frac{1}{2}$	36 $\frac{1}{2}$	20 $\frac{1}{2}$	39 $\frac{1}{2}$	35 $\frac{1}{2}$	27 $\frac{1}{2}$
6	{ 550 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate .. }	69 $\frac{1}{2}$	38 $\frac{1}{2}$	49 $\frac{1}{2}$	50	35 $\frac{1}{2}$	25 $\frac{1}{2}$	60 $\frac{1}{2}$	33 $\frac{1}{2}$	48 $\frac{1}{2}$	44 $\frac{1}{2}$	37 $\frac{1}{2}$	24	57 $\frac{1}{2}$	35 $\frac{1}{2}$	35

SECOND 5 YEARS; MINERAL MANURES AS BEFORE, AMMONIA-SALTS AND NITRATE OF SODA ONLY HALF AS MUCH AS PREVIOUSLY.																	
6TH SEASON, 1874.			7TH SEASON, 1875.			8TH SEASON, 1876 (4).			9TH SEASON, 1877 (5).			10TH SEASON, 1878.			AVERAGE PER ANNUM 5 YEARS, 1874-1878.		
Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.	Dressed Corn.		Total Straw.
Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.		Quantity.	Weight per Bushel.	
1	Unmanured	Bushels. 12	lbs. 31 $\frac{1}{2}$	cwts. 7	Bushels. 12 $\frac{1}{2}$	lbs. 29 $\frac{1}{2}$	cwts. 5 $\frac{1}{2}$	Bushels. 8 $\frac{1}{2}$	lbs. 32	cwts. 2 $\frac{1}{2}$	Bushels. ..	lbs. ..	cwts. ..	Bushels. ..	lbs. ..	cwts. ..	Bushels. ..
2	{ 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate of Lime (1) }	13 $\frac{1}{2}$	31 $\frac{1}{2}$	6 $\frac{1}{2}$	13 $\frac{1}{2}$	29 $\frac{1}{2}$	6 $\frac{1}{2}$	7 $\frac{1}{2}$	30	2 $\frac{1}{2}$	..	..	..	..	..	..	..
3	200 lbs. Ammonia-salts (2)	37 $\frac{1}{2}$	33 $\frac{1}{2}$	22 $\frac{1}{2}$	30 $\frac{1}{2}$	32 $\frac{1}{2}$	15 $\frac{1}{2}$	17 $\frac{1}{2}$	34 $\frac{1}{2}$	6	..	..	..	..	..	..	..
4	{ 200 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate .. }	46 $\frac{1}{2}$	34 $\frac{1}{2}$	24 $\frac{1}{2}$	30 $\frac{1}{2}$	34 $\frac{1}{2}$	20 $\frac{1}{2}$	29 $\frac{1}{2}$	35 $\frac{1}{2}$	12 $\frac{1}{2}$	..	..	..	..	..	..	..
5	275 lbs. Nitrate of Soda (3)	35 $\frac{1}{2}$	30 (4)	16 $\frac{1}{2}$ (4)	23 $\frac{1}{2}$ (4)	31 $\frac{1}{2}$ (4)	11 $\frac{1}{2}$ (4)	12 $\frac{1}{2}$	30 $\frac{1}{2}$	8 $\frac{1}{2}$	..	..	..	..	..	..	..
6	{ 275 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate .. }	28 $\frac{1}{2}$ (4)	33 $\frac{1}{2}$ (4)	16 $\frac{1}{2}$ (4)	28 $\frac{1}{2}$ (4)	33 $\frac{1}{2}$ (4)	14 $\frac{1}{2}$ (4)	19 $\frac{1}{2}$	33 $\frac{1}{2}$	8	..	..	..	..	..	..	..

(1) "Superphosphate of Lime"—in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid sp. gr. 1.7 (and water).

(2) "Ammonia-salts"—in each case, equal parts Sulphate and Murate of Ammonia of Commerce.

(3) 550 lbs. Nitrate of Soda is reckoned to contain the same amount of Nitrogen as 400 lbs. "Ammonia-salts."

(4) On these plots, where large quantities of Nitrate of Soda had been applied year after year, the land, though more worked, was so wet that it could not be got into favourable condition for sowing, and the plant was very irregular.

(5) Owing to the extremely wet condition of the land, especially on the Nitrate plots, it was not sown until April 6, and then with a very unfavourable seed bed; and there being a heavy fall of snow a week later, the plant came up very irregularly, and much of it perished from standing surface-water.

(6) Owing to the very wet winter, 1876-7, the land could not be worked in time for sowing, and was therefore left fallow in 1877.

B 6

EXPERIMENTS ON THE GROWTH OF LEGUMINOUS CROPS.

I.—BEANS, PEAS, AND TARES—GEESCROFT FIELD.

EXPERIMENTS on the growth of Leguminous corn-crops (beans, peas, and tares), with different descriptions of manure, were commenced in 1847, about nine acres being devoted to the purpose.

Experiments with BEANS were continued for thirteen consecutive seasons, to 1859 inclusive; but, during the later years, the crop fell off very much, and the land became very foul.

In 1860 the land was fallowed.

In 1861 a crop of wheat, without manure, was taken.

In 1862 beans were again sown, but with some variation in the manuring.

In 1863 the land was fallowed.

In 1864, 5, 6, 7, 8, and 9, beans were grown, with much the same manures on the same plots, each year, as in 1862.

In the winter of 1869-70, 5000 lbs. of fresh burnt lime were applied per acre, over all the plots.

In 1870 beans were grown with the same manures on the respective plots as in 1864-69.

In October 1870 winter beans were sown (without manure), but the plants were to so great an extent destroyed by the severe weather which followed, that, in April 1871, the crop was ploughed up, and the land left fallow.

During the winter and early spring of 1871-2, the land was so wet that it could not be prepared in time for sowing. It was therefore left fallow for 1872, at the end of May subsoiled to a depth of about 12 inches, and re-ploughed in July. The winter and early spring of 1872-3 were also so extremely wet, that it was again impossible to prepare the land in time for sowing; it was, however, ploughed up towards the end of March, again left fallow, and re-ploughed in July and October (1873). On February 2, 1874, the land was again set with Beans, but without manure. In 1875 Beans were re-sown, with the same manures on the respective plots as in 1864-1870; but owing to the wetness of the land in the first instance, and the subsequent hindrance by other spring sowing, they were not put in until April 1 and 2. The wetness of the winter 1875-6, again prevented the preparation of the land in due time; and, though the manures were sown, and the land ploughed, it was left fallow during the summer of 1876. Winter Beans were put in (drilled), without further manuring, early in October, 1876. For the present crop (1878), the usual manures were sown, and beans were drilled on February 26.

The general result of the experiments with BEANS has been that mineral constituents used as manure (more particularly potass), increased the produce very much during the early years; and, to a certain extent, afterwards, whenever the season was favourable for the crop. Ammonia-salis, on the other hand, produced very little effect; notwithstanding that a Leguminous crop contains two, three, or more times as much nitrogen as a Gramineous one grown under similar conditions as to soil, &c. Nitrate of soda has, however, produced marked effects. But Leguminous crops grown too frequently on the same land seem to be peculiarly subject to disease, which no conditions of manuring that we have hitherto tried seem to obviate.

Experiments with PEAS were soon abandoned, owing to the difficulty of keeping the land free from weeds, and an alternation of BEANS and WHEAT was substituted; the beans being manured much as in the experiments with the same crop grown continuously as above described. But the wetness of the winter of 1871-72 prevented the sowing of the Beans for the season of 1872; and again the wetness of the autumn and winter of 1872-3 prevented the sowing of the wheat until April 4, 1873, when Nursery wheat was put in, which, however, did not come to maturity, but was cut in the middle of September, yielding about 27 cwts. of gross produce per acre, containing too little corn to be worth thrashing. The land was ploughed in October 1873, and sown with beans February 3, 1874. On October 23, 1874, wheat was sown without manure. Beans should have been sown in 1876; indeed, the manures were sown, but, for the reason stated above, the land was left fallow; and wheat was put in October 24 (1876). In 1878 Beans were drilled, on February 26, with the usual manures.

In alternating WHEAT with BEANS, the remarkable result had been obtained, that nearly as much wheat, and nearly as much nitrogen, were yielded in eight crops of wheat in alternation with the highly nitrogenous beans, as in sixteen crops of wheat grown consecutively without manure in another field, and also nearly as much as were obtained in a third field in eight crops alternated with bare fallow.

Experiments with TARES, like those with Peas, were soon abandoned, and for the same reasons. Beans were at first substituted, with some variation in the description of the manures employed; but this experiment has likewise been abandoned for some years.

EXPERIMENTS ON THE GROWTH OF LEGUMINOUS CROPS—continued.

II.—RED CLOVER (*Trifolium pratense*)—HOOS FIELD.

EXPERIMENTS on the growth of Clover, with many different descriptions of manure, were commenced in 1849, and, with the occasional interposition of a corn-crop, or fallow, have been continued up to the present time.

As with other *Leguminous* crops, the result was, that mineral constituents applied as manure (particularly potass) considerably increased the early crops; whereas ammonia-salts had little or no beneficial effect, and were sometimes injurious. It may be added that, even up to the present, the beneficial effects of long previous applications of potass are apparent when there is any growth at all. To go a little more into detail:—

In the first year, 1849, the crops were throughout very heavy; especially with mineral, and without nitrogenous manure.

In autumn 1849 wheat was sown, and in spring 1850 Red Clover. In 1851 small cuttings were taken; and in 1852, though the crops were not heavy, there was by no means a failure. Since that time, however, all attempts to grow clover year after year on the same land have failed to give anything like a full crop, or a plant which would stand the usual time on the ground. Small cuttings were obtained in the autumns of 1855 and 1859 from seed sown in the spring of those years, and small but rather heavier cuttings in June and August 1865, from seed sown in 1864.

On two occasions (1851 and 1854) heavy dressings of Farmyard dung were applied to some of the plots; and in 1854 some received a dressing of 20 tons of dung, and 5000 lbs. of lime, per acre.

On some portions of the land Clover was sown 10 times during the 23 years, 1848–1870, and more frequently alone than with a corn-crop; but in 7 out of the last 8 trials the plant died off in the winter and spring succeeding the sowing the seed.

In view of these failures in the field, it is a fact of much interest, that in 1854 Red Clover was sown in a garden, only a few hundred yards distant from the experimental field, on soil which has been under ordinary garden cultivation for probably two or three centuries, and it has every year since shown very luxuriant growth. Seed was re-sown in 1860, 1865, 1868, and 1871. A small cutting was taken in the autumn of 1871, two cuttings in 1872, and two in 1873. Notwithstanding some injury from dodder in 1873, there still remained too much plant to break up; and, accordingly, fresh seed was sown between the rows on May 4, and this failing, again on July 7, 1874. Small cuttings were taken June 11, July 22, and September 30, 1874. A small cutting was again taken on June 22, 1875. On July 13 the old plants were dug in, and seed again sown, and this failing, seed was re-sown September 22. In spring 1876 there was luxuriant growth, but deficient plant; from which two small cuttings were taken, on June 26, and August 7. On September 1, the beds were dug up, and resown with seed, which came up fairly, but the plant suffered during the winter, and in May 1877 it was dug up and resown. On September 5 (1877) a cutting was taken, and at the present time (May 1878) a vigorous plant remains. This (1878) is, therefore, the 25th season of the growth of Clover, year after year, on this plot of garden ground.

In reference to the field experiments, it may be added that, in 1864, a portion of the land was trenched 2 feet deep, and one-third of the manure was mixed with the layer from 24 to 16 inches, one-third from 16 to 8 inches, and the remainder from 8 inches upwards. Owing to the characters of the season, the mechanical condition of the land was at first very unfavourable after this treatment; but, although many years have now elapsed, and the excess of constituents supplied was in some cases considerable, the plant has died off as completely on these plots as elsewhere.

Again, in the winter of 1867–8 small portions of the experimental land were dug, some to the depth of 9 inches, some to the depth of 18, some to the depth of 27, and some to the depth of 36 inches, and sown to the respective depths with different mixtures; supplying in some cases very large amounts of potass, soda, lime, magnesia, phosphoric acid, sulphuric acid, nitrate of soda, &c. From other similar sized plots, the soil was removed to the depths of 9, 18, and 27 inches respectively, and replaced by soil taken at the same depths from the garden border, on a portion of which clover had been grown successfully since 1854, as above referred to. In April 1868 clover was sown over the whole of these small plots, and on some other portions of the land not so treated; but the plant for the most part died off during the following winter.

In April 1869 the same portions were re-sown, small quantities of clover were cut in September of that year, but the plant again died off in the winter.

In April 1870 Clover was sown over the whole of the experimental land, this time in conjunction with Barley; but on those portions which had also been sown in 1868 and 1869 the plant again died off during the winter and early spring; whilst from those which had not been sown in 1868 and 1869 two small cuttings were taken in 1871. In the spring of 1872, the plant being then almost entirely gone, the land was ploughed up. It was again ploughed in July 1872, and in March 1873; the intention being to sow some other *Leguminous* crop; but owing to the wetness and lateness of the season this was not done; the land was again left fallow, and re-ploughed in the beginning of June and the end of July (1873). On May 4, 1874, the land was again ploughed, and sown with Red Clover seed, May 5, without manure. The plant came up well, and was very forward in September, when the flowering stems were cut down, but left on the land. During the winter and early spring the plant on those portions from which cuttings had been taken in 1871 almost entirely failed, and the land was ploughed up in May, and again in August (1873); whilst on those from which none had been taken since 1869 a fair plant remained, and two small cuttings were obtained, namely on June 23, and on August 9 and 12 (1875). On September 22, this portion of the land was ploughed up. In May (1876) the whole was re-ploughed, again in July and September, and left fallow. In May 1877, Barley and Clover were sown over the whole of the experimental Land, without further manuring, but the clover plant completely died off during the winter. At the present time (1878) the land is devoted to experiments with various *Leguminous* plants.

In the spring of 1871 the small plots in the field were again re-sown, and those of the garden-soil were entirely enclosed, both around and above, by galvanised wire netting. Small

EXPERIMENTS ON THE GROWTH OF LEGUMINOUS CROPS—*continued*.

cuttings were taken from these small beds in July 1872, and (excepting from the garden-soil plots, which had yielded considerably more than the others in 1872) larger cuttings were taken in July 1873. The produce was the largest where potass and nitrate of soda were employed, and where they were applied in the largest quantity, and at the greatest depths. In April 1874 there was still some healthy plant on all the plots, but it was considered to be too irregular to preserve. It was, therefore, dug in. The artificially-manured plots were remanured as before, but only to the depth of 9 inches, and seed was sown on May 4th, July 6th, and October 22nd; each time the plant coming up well, but subsequently dying off. On the Garden soil plots, the plant from the first sowing (May 4), for the most part stood; requiring only to be made good here and there on July 6; and in September small cuttings were taken. In May, 1875, the plant was entirely gone on the artificially-manured plots, which were then dug up, and prepared for resowing. On the garden soil plots, though the rows were imperfect, some healthy plants still remained, and gave a small cutting on June 22. On July 24 these plots were dug up; and they, as well as the artificially manured ones just referred to, were re-sown with seed. All came up well, but in May (1876), the plants on the garden soil plots were entirely gone, and those on the artificially manured ones nearly so, but they yielded small cuttings on July 17. More small plots were arranged in the spring of 1874; on which the manures were dug in, at the various depths, on May 11th to 14th, and the seed sown on May 16th. One series received sulphate of potass only, another nitrate of soda only, and a third the two together. The plants came up fairly well, but there were some blanks in the rows, which were re-sown on October 22 (1874). A cutting was taken on June 22 and 23 (1875); the blanks in the rows were re-sown on July 24; a second cutting taken on August 17; and the blanks again re-sown on September 22 (1875). The plant was the most even on the plots with sulphate of potass, less so on those with nitrate of soda, and less still on those with both together. The amount of produce was also greater with each of the manures used separately, than with the mixture of the two. The plants on these new artificially manured plots, like those on the older ones, showed failure in the spring of 1876; but also, like them, gave small cuttings in July. All the small beds were dug up in August; the artificially manured ones remanured as in 1874, the manures dug in to a depth of 9 inches, and seed was sown on September 1, which came up, but the plants died off on all the plots in the winter of 1876-7. In May 1877, all the small beds were dug up, and sown with Barley and Clover. To try the effects of shelter, the Barley stubble was left unusually high, but the young clover plants completely died off during the winter (1877-8). At the present time (May 1878) the beds are dug up, cleaned, and will be resown with Clover, without further manuring.

The general result of the experiments in the field has been—that neither organic matter rich in carbon as well as other constituents, nor ammonia-salts, nor nitrate of soda, nor mineral constituents, nor a complex mixture, supplied as manure, availed to restore the clover-yielding capabilities of the land; though, where some of these were applied in large quantity, and at considerable depths, the result was better than when they were used in only moderate quantities and applied only on the surface.

On the other hand, it is clear that the garden-soil has supplied the conditions under which clover can be grown year after year on the same land for many years in succession.

The results obtained on the garden-soil seem to show that what is called "clover-sickness," cannot be due to the injurious influence of excreted matters upon the immediately succeeding crop.

That Clover frequently fails coincidently with injury from parasitic plants, or insects, cannot be disputed; but it may be doubted whether such injury should be reckoned as the cause, or merely the concomitant and an aggravation, of the failing condition.

The results of the experiments seem, therefore, to exclude the supposition that the primary cause of failure is either destruction by parasitic plants or insects, injury from excreted matters, or the shade of a corn-crop, and to indicate that it must be looked for in exhaustion of the soil. Still there remain several open questions. Is it exhaustion of certain organic matters rich in carbon, of nitrogenous food, or of mineral constituents? Again: is there an absolute deficiency in the soil of some of the substances in question, or only an unfavourable condition of combination, or, so to speak, of *soil-digestion* of them, for the requirements of Leguminous plants? Or, is there only an unfavourable distribution of them within the soil, considered in relation to the extent and character of the root-range of the crop?

These various suggestions cannot be further considered within the limits of this brief notice, which may be concluded by the following quotation from Rothamsted papers on the subject ('Journal of the Royal Agricultural Society of England,' vol. xxi. Part I. p. 178; and 'Journal of the Royal Horticultural Society of London,' vol. iii. p. 86, 1872).

"When land is not what is called 'clover-sick,' the crop of clover may frequently be increased by top-dressings of manure containing potass and superphosphate of lime; but the high price of salts of potass, and the uncertainty of the action of manures upon the crop, render the application of artificial manures for clover a practice of doubtful economy.

"When the land is what is called 'clover-sick,' none of the ordinary manures, whether 'artificial' or natural, can be relied upon to secure a crop.

"So far as our present knowledge goes, the only means of insuring a good crop of Red Clover is to allow some years to elapse before repeating the crop upon the same land."

EXPERIMENTS ON ROOT-CROPS.—BARN FIELD.

Experiments with Turnips were commenced in 1843. Eight acres, divided into numerous Plots, were set apart for the purpose, and the crop was grown for ten consecutive years on the same land; "Norfolk Whites" 1843-1848, and "Swedes" 1849-1852; on some Plots without manure, and on others with different descriptions of manure. Barley was then grown for three consecutive seasons, 1853-1855, without manure, in order to test the comparative corn-growing condition of the different Plots, and also to equalise their condition, as far as possible, by the exhaustion of some of the most active and immediately available constituents supplied by the previous manuring. A new series of experiments with Swedes was arranged in 1856, having regard to the character of the manures previously applied on the different Plots, and to the results previously obtained. This second series was continued for fifteen years, namely, from 1856 to 1870 inclusive. The results obtained with Norfolk Whites in the first three years, 1843, 1844, and 1845, were published in the 'Journal of the Royal Agricultural Society of England,' vol. viii. Part II., 1847; and an abstract of the results obtained from 1845 to 1870 inclusive, is given in the Table below. During the five years, 1871-1875, the land was devoted to experiments with Sugar-Beet, for particulars of which see pp. 10 and 11. In 1876 experiments with Mangold-wurzel were substituted, and are still in progress (see p. 12).

(Area under experiment, about 8 acres; quantities, average, per acre, per annum.)

NORFOLK WHITE TURNIPS; FOUR SEASONS, 1845-1848; Roots and Leaves carted off the Land.																	
SERIES 1. Manures as under; no Cross-dressing.				Each Plot as Series 1, and Cross-dressed as under—													
				SERIES 2. No Cross-dressing.		SERIES 3. 160 lbs. Sulphate Amm. nia. 75 lbs. Muriate Ammonia.		SERIES 4. 160 lbs. Sulphate Ammonia. 75 lbs. Muriate Ammonia. 1840 lbs. Rape-cake.		SERIES 5. 1840 lbs. Rape-cake.							
Average Produce, per Acre, per Annum.																	
Plots.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.					
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.						
3	Gypsum 1845; without Manure 1846 and since (average 1846, 7, 8)	1	4	0	17	1	7	1	0	5	10	6	11	3	3		
4	Superphosphate, each year; Potass, Soda, and Magnesia, 1847-8 ..	8	1	2	15	9	15	4	3	10	5	6	1	11	2	4	12
5	Superphosphate, each year;	8	16	2	19	9	18	4	8	10	1	6	3	10	18	4	15
6	Superphosphate, each year; and Potass 1847-8	8	0	2	19	9	16	5	4	10	7	6	17	10	17	5	7
7																	
SWEDISH TURNIPS; FOUR SEASONS, 1849-1852; Roots and Leaves carted off the Land (excepting 1849, when the Leaves were too small to weigh or remove).																	
SERIES 1. Manures as under; no Cross-dressing.				Each Plot as Series 1, and Cross-dressed, as under, in 1849 and 1850. No Cross-dressing in 1851 and 1852.													
				SERIES 2. No Cross-dressing.		SERIES 3. 200 lbs. Ammonia-salts.		SERIES 4. 200 lbs. Ammonia-salts. 2000 lbs. Rape-cake.		SERIES 5. 2000 lbs. Rape-cake.							
Plots.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.	
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.		
3	Without Manure, 1846 and since	2	6	0	6	3	17	0	6	7	0	0	17	7	14	0	13
4	Superphosphate, Sulphates Potass and Magnesia, and Soda-ash ..	7	17	0	10	9	9	0	11	13	1	0	18	12	7	0	15
5	Superphosphate	7	9	0	11	8	14	0	13	11	4	1	1	10	10	0	17
6	Superphosphate, and Sulphate Potass	6	16	0	9	8	14	0	10	12	8	0	17	11	14	0	14
7																	
BARLEY, without Manure (after Roots manured as above); THREE SEASONS, 1853-1855.																	
SERIES 1.				SERIES 2.		SERIES 3.		SERIES 4.		SERIES 5.							
						Dressed Corn.	Straw.	Dressed Corn.	Straw.	Dressed Corn.	Straw.						
Plots.		Dressed Corn.		Straw.		Dressed Corn.		Straw.		Dressed Corn.		Straw.					
		Bushels.	Cwts.	Bushels.	Cwts.	Bushels.	Cwts.	Bushels.	Cwts.	Bushels.	Cwts.						
3		18½	12½	20½	12½	20½	12½	25½	15½	25½	15½	25½	15½				
4		20½	12½	22½	13	22½	13	25	14½	25½	14½	25½	14½				
5		21	11½	23	12½	23	12½	26½	15	27	15½	27	15½				
6		18½	10½	20½	11½	20½	11½	25	14½	25	14½	25	14½				
7																	
SWEDISH TURNIPS; FIFTEEN SEASONS, 1856-1870. (1) Roots and Leaves carted off the Land.																	
SERIES 1. Manures as under; no Cross-dressing.				Each Plot as Series 1, and Cross-dressed as under—													
				SERIES 2. 5 years, 1856-1860. 3/00 lbs. Saw-dust. 328 lbs. Nitric Acid.		SERIES 3. 5 years, 1856-1860. 200 lbs. Ammonia-salts.		SERIES 4. 5 years, 1856-1860. 200 lbs. Ammonia-salts. 3000 lbs. Sawdust.		SERIES 5. 5 years, 1856-1860. 3600 lbs. Sawdust.							
Plots.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.	
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.		
1	Farmyard Manure, 14 tons	6	4	0	17	7	9	1	2	8	8	1	9	8	0	1	4
2	Farmyard Manure, 14 tons, and Superphosphate	6	7	0	16	7	13	1	3	8	5	1	5	8	14	1	9
3	Without Manure, 1846, and since	0	11	0	3	0	19	0	4	0	13	0	3	3	6	0	14
4	Superphosph., each year; Sulph. Potass, Soda, and Magnesia, 1856-60	2	16	0	8	5	2	0	16	4	12	0	14	6	12	1	6
5	Superphosphate, each year	2	12	0	9	4	13	0	18	3	16	0	15	5	16	1	7
6	Superphosphate, each year; Sulphate Potass, 1856-1860	2	7	0	7	4	11	0	14	4	5	0	13	6	6	1	2
7	Superphosph., each year; Sulph. Potass, and 36½ Amm.-salts, 1856-60	2	12	0	7	4	13	0	14	4	12	0	14	6	15	1	4
8	Unman. 1853, and since; previously part Unman.; part Superphosph.	1	3	0	4	1	13	0	5	1	2	0	5	3	19	0	18

NORF.—"Sulphate of Ammonia" is estimated to contain 23 per cent. Ammonia and "Muriate of Ammonia" 27 per cent. "Ammonia-salts," in each case, equal parts Sulphate and Muriate of Ammonia of commerce; and the mixture is estimated to contain 25 per cent. Ammonia. The 328 lbs. Nitric Acid (Sp. gr. 1.35), mixed with sawdust, and used as a cross-dressing on the Plots of Series 2, from 1856-1860, were estimated to contain Nitrogen = 50 lbs. Ammonia.

(1) The crops of 1859 and 1860 failed, and were ploughed in; but, as the manures were applied, and there would be accumulation within the soil for the succeeding crops, the average produce is calculated as for 15 years, that is the produce of the 15 years is, in each case, divided by 15.

(16)

EXPERIMENTS ON SUGAR BEET (VILMORIN'S GREEN-TOP WHITE SILESIAK)—BARN FIELD.

GROWN YEAR AFTER YEAR ON THE SAME LAND, WITHOUT MANURE, AND WITH DIFFERENT DESCRIPTIONS OF MANURE, COMMENCING 1871.

Previous Cropping:—1843-48 (6 Seasons), experiments on Norfolk White Turnips, with different descriptions of Manure.

1849-52 (4 Seasons), experiments on Swede Turnips, with different descriptions of Manure.

1853-55 (3 Seasons), Barley without Manure (with a view as far as possible to equalise the condition of the Plots).

1856-70 (15 Seasons), experiments on Swede Turnips, with different descriptions of Manure, in which the arrangement of the

Plots was the same, and that of the Manures very similar—in fact, exactly the same during the last 10 years—as in the first year of Sugar Beet, excepting that, during those 10 years, the Alkalies were omitted for the Swedes. For the second and subsequent years of Sugar Beet slight alterations in the Mineral Manures were made, and in the fourth and fifth years the Farmyard Manure, Nitrate of Soda, Ammonia-salts, and Rape-cake were omitted, as will be seen below. Seed dibbled on the flat; in rows 22 inches apart, and 11 inches apart in the rows; plants moulded up afterwards. Roots all carted off, Leaves weighed, spread on the respective Plots, and ploughed in.

Area under experiment about 8 acres. The experiments are arranged as under, in 5 Series, each of which comprises 8 Plots.

		Manures, per Acre, per Annum.									
PLOTS.		SERIES 1.		SERIES 2.		SERIES 3.		SERIES 4.		SERIES 5.	
		Each Plot as Series 1.		Each Plot as Series 1, and Cross-dressed with 550 lbs. Nitrate Soda.		Each Plot as Series 1, and Cross-dressed with 400 lbs. "Ammonia-salts."		Each Plot as Series 1, and Cross-dressed with 2000 lbs. Rape-cake, and 400 lbs. "Ammonia-salts."		Each Plot as Series 1, and Cross-dressed with 2000 lbs. Rape-cake.	
FIRST SEASON, 1871.											
		PRODUCE PER ACRE (Roots trimmed as for feeding, not as for Sugar-making).									
		Roots.		Leaves.		Roots.		Leaves.		Roots.	
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)	18 3	3 5	27 13	6 19	22 1	5 6	26 4	6 14	28 18	5 14
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	14 13	2 14	25 16	5 15	21 15	4 6	25 2	6 7	25 4	5 5
3	Without Manure (1846, and since)	7 11	2 0	22 3	5 12	15 6	4 16	19 18	7 0	20 16	4 12
4	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass, 200 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia	7 11	1 5	22 15	4 8	17 10	3 5	22 15	6 3	21 7	3 19
5	3½ cwt. Superphosphate	5 12	1 8	20 19	3 14	15 4	3 19	19 18	7 12	18 19	4 5
6	3½ cwt. Superphos., 300 lbs. Sulph. Potass	5 1	1 4	21 5	3 13	17 4	3 4	23 11	6 11	21 0	3 11
7	3½ cwt. Superphos., 300 lbs. Sulph. Pot., 36½ lbs. Amm.-salts (?) ..	5 18	1 5	20 19	3 13	18 8	4 3	21 0	5 0	21 7	3 17
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	7 10	1 14	21 13	3 16	16 2	4 15	17 19	7 11	20 7	4 9
SECOND SEASON, 1872.											
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)	15 13	4 2	23 9	7 19	22 14	9 0	26 8	9 11	22 5	6 1
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	16 0	3 18	24 6	8 16	22 0	7 16	25 9	9 14	20 15	5 11
3	Without Manure (1846, and since)	7 17	1 13	21 7	6 6	15 3	4 13	20 8	10 1	16 3	3 11
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride Sodium (common salt), 200 lbs. Sulphate Magnesia	6 14	1 10	20 2	5 19	15 10	3 7	23 8	7 13	17 18	3 15
5	3½ cwt. Superphosphate	6 17	1 8	19 6	6 4	14 5	4 13	18 11	10 4	15 18	3 16
6	3½ cwt. Superphos., 500 lbs. Sulph. Potass	6 6	1 5	16 16	5 14	14 7	3 19	22 16	9 9	15 17	3 14
7	3½ cwt. Superphos., 500 lbs. Sulph. Potass, 36½ lbs. Amm.-salts (?) ..	6 15	1 8	17 0	6 1	15 9	3 19	23 9	9 10	15 10	3 15
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	5 4	1 5	15 6	5 19	13 10	4 1	19 12	9 17	15 0	4 6
THIRD SEASON, 1873.											
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)	15 2	5 12	20 5	10 9	22 2	9 18	22 15	12 10	23 10	7 8
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	14 6	5 2	21 10	11 0	19 4	8 9	23 7	13 6	21 18	6 18
3	Without Manure (1846, and since)	5 1	1 11	14 5	6 11	9 3	3 16	15 12	9 11	14 13	4 1
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride Sodium (common salt), 200 lbs. Sulphate Magnesia	5 2	1 13	16 9	6 11	12 10	3 10	20 3	8 0	16 1	3 8
5	3½ cwt. Superphosphate	5 5	1 11	18 8	5 13	10 19	5 0	14 15	9 8	13 19	4 9
6	3½ cwt. Superphos., 500 lbs. Sulph. Potass	4 12	1 5	15 17	4 4	12 18	3 12	20 2	9 5	14 14	3 11
7	3½ cwt. Superphos., 500 lbs. Sulph. Potass, 36½ lbs. Amm.-salts (?) ..	5 19	1 12	16 14	5 3	13 0	4 15	19 16	9 0	15 17	4 4
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	4 11	1 7	12 9	5 18	8 8	2 19	15 2	9 8	12 2	3 16
FOURTH SEASON, 1874 (?). Mineral Manures as in 1872 and 1873; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake.											
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Without Manure, 1874 and 1875 (Farmyard Manure in '71, '72, '73) ..	10 16	5 6	11 14	8 9	11 7	8 3	13 7	9 17	14 10	7 8
2	3½ cwt. Superphosphate (with Farmyard Manure, '71, '72, '73) ..	13 3	5 9	7 9	4 16	9 5	5 17	12 5	7 7	13 1	6 4
3	Without Manure (1846, and since)	5 2	1 5	3 2	2 6	3 7	2 2	2 11	2 10	3 19	2 9
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride Sodium (common salt), 200 lbs. Sulphate Magnesia	6 10	1 8	8 16	3 6	7 10	2 0	10 12	4 16	8 2	3 11
5	3½ cwt. Superphosphate	5 19	1 7	7 10	3 6	7 6	2 8	7 15	5 4	5 17	3 6
6	3½ cwt. Superphos., 500 lbs. Sulph. Potass	5 11	1 5	8 1	2 14	8 1	1 18	9 10	4 13	7 13	3 2
7	3½ cwt. Superphos., 500 lbs. Sulph. Pot., and Amm.-salts, '71, '72, '73 ..	6 14	1 3	9 5	2 11	8 15	1 14	11 14	4 11	8 4	3 9
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	5 0	1 2	7 13	2 16	6 10	2 0	7 6	4 7	3 12	2 1
FIFTH SEASON, 1875. Mineral Manures as in 1872, 1873, and 1874; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake.											
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Without Manure, 1874 and 1875 (Farmyard Manure in '71, '72, '73) ..	17 5	2 11	19 18	2 14	21 0	3 6	22 7	3 12	19 13	2 11
2	3½ cwt. Superphosphate (with Farmyard Manure, '71, '72, '73) ..	15 11	2 2	19 18	2 18	18 17	2 18	20 9	3 5	18 10	2 1
3	Without Manure (1846, and since)	5 9	1 1	9 5	1 12	8 0	1 3	14 1	2 13	11 17	1 10
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride Sodium (common salt), 200 lbs. Sulphate Magnesia	5 9	1 0	9 8	1 7	7 16	1 1	12 14	1 14	10 3	1 7
5	3½ cwt. Superphosphate	5 11	1 2	9 19	1 10	7 16	1 4	13 17	2 8	11 2	1 14
6	3½ cwt. Superphos., 500 lbs. Sulph. Potass	5 4	1 0	8 4	1 4	7 1	1 2	12 8	2 3	10 2	1 9
7	3½ cwt. Superphos., 500 lbs. Sulph. Pot. and Amm.-salts '71, '72, '73 ..	5 11	1 1	8 2	1 6	7 6	1 1	11 17	1 17	10 6	1 11
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	4 15	1 0	7 4	1 2	6 1	1 4	12 2	2 11	11 12	2 13

(1) "Superphosphate of Lime"—in all cases made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid sp. gr. 1.7 (and water).

(2) "Ammonia-salts"—in each case equal parts Sulphate and Muriate of Ammonia of Commerce.

(3) Owing to the deficiency of rain for some time after sowing a large proportion of the plants failed. Some were transplanted on plots 1, but not on the other plots; and eventually the plant was (excepting on plots 1) upon the whole very deficient and irregular, the remaining plants being larger than usual.

EXPERIMENTS ON SUGAR BEET—BARN FIELD—*continued*.

As it will be some time before we shall be able to report fully the results obtained illustrating the influence of different manures, and different seasons, on the composition of Sugar-beet, an abstract of the analytical results obtained is given below. In interpreting the figures it must be borne in mind that with forty different experiments each year, and in each year 4 or 5 or more times as much produce on some plots as on others, it would be impossible to sample each at its best, and all in the same condition of ripeness. Each year the seed was sown on all the Plots at the same time; and the samples (each consisting of the vertical fourths of 10 or 15 roots) were taken from all within a period of about a week, beginning with the ripest. It is obvious, however, that the smaller crops would be much riper than the larger ones. It need only further be observed that although, in comparable cases, the larger crops generally give a juice containing a lower percentage of sugar and higher percentages of mineral matter and of nitrogen, yet, the larger crops yielded very much more sugar over a given area of land.

MEAN PER CENT. SUGAR, MINERAL MATTER (CRUDE ASH), AND NITROGEN, IN JUICE, in Selected cases, each year; 5 years, 1871-5;

and

AVERAGE PRODUCE AND COMPOSITION OF THE ROOTS; FIRST THREE SEASONS, 1871, 1872, and 1873.

FOR MANURES, see page 10.		CROSS-DRESSED MANURES PER ACRE PER ANNUM.														
		SERIES 1. No Cross-dressing.			SERIES 2. As Series 1, and Cross-dressed with 550 lbs. Nitrate Soda.			SERIES 3. As Series 1, and Cross-dressed with 400 lbs. "Ammonia-salts."			SERIES 4. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake, and 400 lbs. "Ammonia-salts."			SERIES 5. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake.		
		MEAN PER CENT. SUGAR, MINERAL MATTER (CRUDE ASH), AND NITROGEN, IN JUICE.														
FIRST SEASON, 1871.																
		Sugar.	Ash.	Nitrogen.	Sugar.	Ash.	Nitrogen.	Sugar.	Ash.	Nitrogen.	Sugar.	Ash.	Nitrogen.	Sugar.	Ash.	Nitrogen.
		Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.
Plot 1		12.39	0.697	..	10.27	0.897	..	11.63	0.776	..	9.85	0.936	..	10.79	0.776	..
" 4		13.68	0.528	0.096	11.38	0.707	0.166	12.49	0.668	0.141	10.42	0.764	0.224	12.31	0.670	0.133
" 5		13.92	0.553		11.65	0.640		12.04	0.662		9.76	0.730		12.47	0.582	
" 6		13.68	0.597		11.02	0.742		12.12	0.742		10.22	0.772		12.71	0.668	
Means of Plots 4, 5, and 6 ..		13.76	0.559	0.096	11.35	0.696	0.166	12.21	0.691	0.141	10.13	0.755	0.224	12.49	0.640	0.133
SECOND SEASON, 1872.																
Plot 1		13.65	0.742	..	12.67	0.877	..	12.58	0.820	..	12.70	0.844	..	13.00	0.818	..
" 4		14.90	0.647	0.099	12.83	0.810	0.146	14.02	0.698	0.123	13.33	0.816	0.186	14.08	0.717	0.143
" 5		14.65	0.537	0.091	11.75	0.824	0.176	13.71	0.584	0.148	10.95	0.844	0.236	13.92	0.576	0.146
" 6		14.54	0.581	..	12.51	0.760	..	14.17	0.728	..	12.79	0.780	..	13.86	0.661	..
Means of Plots 4 and 5 ..		14.78	0.592	0.095	12.29	0.817	0.161	13.87	0.641	0.136	12.14	0.830	0.211	14.00	0.647	0.145
THIRD SEASON, 1873.																
Plot 1		13.40	0.756	..	11.79	0.905	..	11.93	0.845	..	10.75	0.948	..	12.25	0.540	..
" 4		14.54	0.619	0.132	12.69	0.831	0.174	13.80	0.774	0.158	11.80	0.842	0.176	13.87	0.700	0.147
" 5		15.02	0.499	0.110	12.11	0.895	0.179	13.86	0.555	0.183	12.26	0.632	0.212	14.19	0.561	0.169
" 6		15.11	0.603	0.114	13.15	0.689	0.156	13.91	0.726	0.126	12.52	0.781	0.198	13.66	0.698	0.148
Means of Plots 4, 5, and 6 ..		14.89	0.574	0.119	12.65	0.785	0.169	13.86	0.685	0.156	12.19	0.752	0.195	13.91	0.653	0.155
FOURTH SEASON, 1874 (?). Mineral Manures as in 1872 and 1873; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake.																
Plot 1		11.74	0.972	0.260	10.69	1.144	..	10.30	1.121	..	10.78	1.129	..	11.42	0.935	..
" 4		13.79	0.528	0.103	10.24	0.756	0.135	13.06	0.762	0.157	12.23	0.865	0.211	13.21	0.772	0.162
" 5		13.69	0.474	0.109	10.29	0.794	0.187	13.07	0.662	0.182	12.16	0.650	0.207	11.89	0.724	0.237
" 6		13.67	0.496	0.103	11.05	0.714	0.184	14.41	0.697	0.143	12.68	0.781	0.208	11.62	0.816	0.189
Means of Plots 4, 5, and 6 ..		13.72	0.499	0.105	10.53	0.755	0.169	13.51	0.707	0.161	12.36	0.765	0.209	12.07	0.771	0.199
FIFTH SEASON, 1875. Mineral Manures as in 1872, 1873, and 1874; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake.																
Plot 1		12.33	0.626	0.136	12.47	0.637	..	12.12	0.675	..	12.65	0.718	..	12.18	0.668	..
" 4		12.75	0.607	0.094	12.69	0.606	0.106	12.97	0.652	0.116	12.52	0.674	0.115	12.30	0.695	0.115
" 5		13.67	0.536	0.104	12.73	0.582	0.114	12.72	0.573	0.113	11.79	0.580	0.137	12.43	0.513	0.106
" 6		13.33	0.541	0.107	13.13	0.637	..	12.85	0.663	0.110	12.19	0.669	0.150	12.73	0.656	0.118
Means of Plots 4, 5, and 6 ..		13.25	0.561	0.102	12.71	0.594	0.110	12.85	0.629	0.113	12.17	0.641	0.134	12.49	0.621	0.113

AVERAGE PRODUCE AND COMPOSITION, FIRST THREE SEASONS, 1871, 1872, and 1873.

PLOT 1 (SERIES I.), Farmyard Manure (14 Tons).

Average produce per acre:—	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.
Roots	326	476	446	502	498
Leaves	86	169	161	192	128
Total	412	645	607	694	626
Average Composition of the Roots:—	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.
Dry Matter	17.49	16.11	16.56	16.23	16.66
Mineral Matter (ash) in Dry Matter ..	5.00	6.11	5.83	6.55	5.61
Nitrogen in Dry Matter (?)	0.83	1.24	1.53	1.52	1.24
Sugar in Juice	13.14	11.58	12.05	11.10	12.01
Sugar in Roots, if 95, P.C. Juice ..	12.48	11.00	11.45	10.55	11.41

MEANS OF PLOTS 4, 5, and 6 (SERIES I.), Superphosphate, with or without other Mineral Manures, every year.

Average produce per Acre:—	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.
Roots	118	382	290	413	346
Leaves	28	102	76	165	76
Total	146	484	366	578	422
Average Composition of the Roots ..	Per Cent.	Per Cent.	Per Cent.	Per Cent.	Per Cent.
Dry Matter	18.53	15.93	17.43	15.93	17.66
Mineral Matter (ash) in Dry Matter ..	4.30	5.73	4.81	5.98	4.50
Nitrogen in Dry Matter (?)	0.54	1.20	0.87	1.52	0.83
Sugar in Juice	14.45	12.12	13.35	11.56	13.45
Sugar in Roots, if 95, P.C. Juice ..	13.73	11.51	12.68	10.98	12.78

(1) Owing to the deficiency of Rain for some time after sowing a large proportion of the plants failed. Some were transplanted on plots 1, but not on the other plots; and eventually the plant was (excepting on plots 1) upon the whole very deficient and irregular, the remaining plants being larger than usual.

(2) The percentages of Nitrogen in the roots relate to the first year only; but the percentages of Nitrogen determined in the Juice, in selected cases, each year, confirm the indications of the nitrogen in the roots in the first year.

(18)

EXPERIMENTS ON MANGOLD WURZEL.—BARN FIELD (after SUGAR-BEET); commencing 1876.

The arrangement of the Plots is precisely the same as previously for Sugar-beet, excepting that Plot 9, which was unmanured for Sugar-beet, and also previously for Swedes, is now added as a manured Plot. With this exception, the manures are also substantially the same as previously for Sugar-beet; in fact, precisely the same as for the Sugar-beet in 1872 and 1873. Seed, Yellow Globe; dibbled on ridges, rows 26 inches apart; plants 11 inches apart in the rows (?). Area under experiment about 8 acres. Roots all carted off; Leaves weighed, spread on the respective Plots, and ploughed in.

PLOTS.	MANURES PER ACRE PER ANNUM.									
	SERIES 1.		SERIES 2.		SERIES 3.		SERIES 4.		SERIES 5.	
			As Series 1, and Cross-dressed with 550 lbs. Nitrate Soda.		As Series 1, and Cross-dressed with 400 lbs. "Ammonia- salts."		As Series 1, and Cross-dressed with 2000 lbs. Rape-cake and 400 lbs. "Am- monia-salts."		As Series 1, and Cross-dressed with 2000 lbs. Rape-cake.	

FIRST SEASON, 1876. Seed dibbled, May 22-26. Crop taken up, Nov. 3-17.

		PRODUCE PER ACRE.									
		Roots.		Leaves.		Roots.		Leaves.		Roots.	
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)	19 12	4 9	25 2	7 5	29 19	7 12	31 9	10 5	24 9	5 19
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	19 13	4 6	27 13	7 3	29 8	7 10	30 18	9 16	29 19	6 12
3	Without Manure (1846, and since)	6 10	1 14	20 13	5 12	14 3	4 10	19 19	7 7	17 4	4 15
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride (Sodium (common salt), 200 lbs. Sulphate Magnesia)	8 8	1 15	25 1	6 0	19 19	4 9	30 8	8 13	25 8	5 10
5	3½ cwt. Superphosphate	7 10	1 14	21 0	5 14	13 10	5 1	17 2	7 14	17 17	5 17
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	6 16	1 12	21 2	5 8	17 15	4 13	26 8	9 0	20 10	5 4
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	8 13	2 3	22 11	5 14	19 2	5 11	27 2	9 9	20 12	5 15
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	5 9	1 10	15 16	5 3	11 17	4 16	18 2	7 11	15 12	4 18
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	..	..	..	..	25 14	7 6	..	..	..	..

SECOND SEASON, 1877. Seed dibbled, June 4-6 (Plots 8 and 9, June 11th). Crop taken up, Nov. 14-23.

		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)	15 7	2 1	24 13	3 14	27 1	4 4	30 5	5 5	25 13	3 4
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	16 14	1 19	26 8	3 12	26 18	4 6	28 15	5 9	24 12	2 19
3	Without Manure (1846, and since)	5 9	1 0	16 17	3 14	8 16	3 0	13 9	3 19	13 17	2 10
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride (Sodium (common salt), 200 lbs. Sulphate Magnesia)	6 16	1 3	21 10	3 10	16 10	2 2	27 9	3 8	21 14	1 17
5	3½ cwt. Superphosphate	6 1	0 19	20 5	3 1	12 2	2 10	15 3	3 8	15 3	2 2
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	5 8	0 18	20 19	2 18	15 6	1 16	24 18	3 16	19 3	1 12
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	7 0	1 3	22 2	3 16	16 13	2 7	25 15	5 0	20 13	2 8
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	3 19	1 3	9 17	5 4	7 4	3 10	11 9	4 11	10 3	3 3
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	..	..	..	..	13 17	4 0	..	..	..	..

THIRD SEASON, 1878.

		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)										
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..										
3	Without Manure (1846, and since)										
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride (Sodium (common salt), 200 lbs. Sulphate Magnesia)										
5	3½ cwt. Superphosphate										
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass										
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)										
8	Unmanured, 1853, and since; previously part Unman., part Superphos.										
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)										

FOURTH SEASON, 1879.

		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)										
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..										
3	Without Manure (1846, and since)										
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride (Sodium (common salt), 200 lbs. Sulphate Magnesia)										
5	3½ cwt. Superphosphate										
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass										
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)										
8	Unmanured, 1853, and since; previously part Unman., part Superphos.										
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)										

FIFTH SEASON, 1880.

		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure (14 tons)										
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..										
3	Without Manure (1846, and since)										
4	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride (Sodium (common salt), 200 lbs. Sulphate Magnesia)										
5	3½ cwt. Superphosphate										
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass										
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)										
8	Unmanured, 1853, and since; previously part Unman., part Superphos.										
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)										

(1) "Superphosphate of Lime"—in all cases made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid, sp. gr.; 1·7 (and water).

(2) "Ammonia-salts"—in each case equal parts Sulphate and Muriate of Ammonia of Commerce.

(3) Plot 9 sown on the flat instead of on ridges; plants ridged up afterwards; rows 22 inches apart, plants 10 inches apart in the rows.

(19)

EXPERIMENTS ON POTATOES.—HOOS FIELD; commencing 1876.

The Land had been under experiments with Wheat, differently manured, from 1856 to 1874; and was fallowed in 1875.

Plots 1, 2, 3, and 4 had been unmanured for the Wheat. Plots 5 and 6 had received the same quantity of Ammonia-salts alone every year for the Wheat, as Plot 5 now receives for potatoes: Plot 6 now receiving the same amount of nitrogen, but as Nitrate of Soda, instead of Ammonia-salts. Plots 7 and 8 received the same amount of complex mineral manure, and Ammonia-salts, for the Wheat, as Plot 7 now receives for potatoes; and Plot 8 now receives the same complex mineral manures, and the same amount of nitrogen, but as Nitrate of Soda instead of Ammonia-salts. Plots 9 and 10 received the same complex mineral manures alone for the Wheat as Plot 10 now receives for potatoes; Plot 9 now receives superphosphate only (?). Description of Potatoes, "Rock." Rows 25 inches apart; 12 inches from plant to plant in the rows.

PLOTS.	MANURES PER ACRE PER ANNUM.	PRODUCE PER ACRE.				
		Tubers.				Tops.
		Good.	Small.	Diseased.	TOTAL.	

FIRST SEASON, 1876. Potatoes planted, June 10-13; Crop taken up, Oct. 30-31.

1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	
2	Farmyard Manure (14 tons)	3 6½	0 5½	0 5½	3 17½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	3 18½	0 4	0 3½	4 5½	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	4 14½	0 6½	0 5½	5 6½	
5	400 lbs. Ammonia-salts (?)	5 5½	0 5½	0 19½	6 14½	
6	550 lbs. Nitrate of Soda	2 5½	0 6½	0 6	2 18	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	3 2	0 5½	0 9½	3 17½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	6 12½	0 9½	1 0	8 2	
9	3½ cwt. Superphosphate	6 17½	0 10	1 8½	8 15½	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	4 18½	0 8½	0 13½	6 1	
		5 3½	0 6½	0 19½	6 3½	

Withered, not weighed, each lot spread on its own Plot and ploughed in.

SECOND SEASON, 1877. Potatoes planted, April, 27-28; Crop taken up, Oct. 8-10.

1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	
2	Farmyard Manure (14 tons)	2 11½	0 6½	0 2½	3 0½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	5 0½	0 11½	0 6	5 18	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	4 13½	0 7½	0 4	5 4½	
5	400 lbs. Ammonia-salts (?)	6 18½	0 7	0 17½	8 3½	
6	550 lbs. Nitrate of Soda	3 9½	0 7½	0 4	4 1	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	4 14½	0 6½	0 5½	5 7½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	6 12	0 11½	0 14½	7 17½	
9	3½ cwt. Superphosphate	7 8½	0 8½	0 16½	8 13½	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	2 12½	0 11½	0 1½	3 6	
		3 6½	0 7½	0 1½	3 15½	

Withered, not weighed, each lot spread on its own Plot, but high wind (Oct. 14th) blew all off, before ploughing.

THIRD SEASON, 1878. Potatoes planted, April 29. Crop taken up.

1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	
2	Farmyard Manure (14 tons)	2 11½	0 8½	0 2	2 11½	0 13½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	4 11	0 11½	0 8½	5 11½	0 6½	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	5 18½	0 11½	0 13½	7 6	0 11	
5	400 lbs. Ammonia-salts (?)	6 11½	0 11½	1 6½	8 9½	1 6½	
6	550 lbs. Nitrate of Soda	3 11½	0 8½	0 5½	3 10½	0 11	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	3 11½	0 7½	0 4½	4 13½	0 10½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	4 11½	0 11½	0 11½	5 11½	0 13½	
9	3½ cwt. Superphosphate	4 11½	0 11½	0 11½	5 11½	0 13½	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	3 11½	0 8½	0 4½	4 11½	0 11½	

Withered, weighed, each lot spread on its own Plot and ploughed in.

FOURTH SEASON, 1879.

1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
2	Farmyard Manure (14 tons)					
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)					
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda					
5	400 lbs. Ammonia-salts (?)					
6	550 lbs. Nitrate of Soda					
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.					
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.					
9	3½ cwt. Superphosphate					
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia					

FIFTH SEASON, 1880.

1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
2	Farmyard Manure (14 tons)					
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)					
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda					
5	400 lbs. Ammonia-salts (?)					
6	550 lbs. Nitrate of Soda					
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.					
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.					
9	3½ cwt. Superphosphate					
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia					

(1) "Superphosphate of Lime"—in all cases made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid, sp. gr. 1.7 (and water).

(2) "Ammonia-salts"—in each case equal parts Sulphate and Muriate Ammonis of Commerce.

(3) The complex mineral manure having been sown in October, 1874, but the Wheat not put in, and therefore no crop taken in 1875, no mineral manures are sown afresh on Plots 7, 8, 9, and 10, for the first crop of potatoes, 1876.

[illegible]

DRESSED CORN PER ACRE. Bushels.									
1. White-chaff (Red)	..	..	40½	55½	40½	49½	55½	48½	46½
2. Rivett's (Red)	..	..	48½	67	48½	49½	66½	49½	51½
3. Chubb Wheat (Red)	..	..	35½	50½	38½	41½	50½	41½	39½
4. Red-chaff (White)	..	40	35½	48½	34½	41	45½	41	39
5. Brownick (Red)	..	37	35½	51½	38½	40½	49½	40½	40½
6. Red Wonder	..	40½	38½	51½	38½	39½	44½	40½	40½
7. Burwell (Old Red Lammus)	..	31½	43½	55½	37½	41½	44½	38½	38½
8. Bristol Red	..	44½	35½	47½	31½	42½	44½	39	37½
9. Red Nursery	..	41½	27½	53½	39½	37½	40½	37½	40½
10. Red Langham	..	43½	34½	53½	34½	42½	42½	40½	40½
11. Woolly Ear (White)	..	42½	37	51½	38½	37½	37½	40½	40½
12. Hardcastle (White)	..	49½	42	49½	33½	44½	42½	42½	43
13. Golden Drop (Red), Hallett's	..	38½	44½	51½	38½	44½	49½	45½	45½
14. Victoria White, Hallett's	..	35½	45½	44½	33½	41½	42½	42½	42½
15. Hunter's White, Hallett's	..	26½	39½	45½	26½	40½	44½	42½	42½
16. Original Red, Hallett's	..	30	36½	43½	26	40½	44½	42½	42½
17. White Chiddam	..	28½	31½	42	32½	37½	37½	42½	42½
18. Red Rostock	..	37	46½	53½	37½	40	46½	43	43
19. Casey's White	..	27½	37½	52½	38½	45½	43	41½	41½
20. Golden Rough-chaff (Red)	..	33	38½	52½	38½	38½	36½	39½	39½
21. Bole's Prolific (Red)	..	33½	43½	48½	44½	44½	44½	42½	42½
22. Club Wheat (Red)	..	36	45½	59½	46½	47½	49½	47½	47½
23. Main's Standing White	..	..	..	..	..	..	..	..	..
24. Main's Rough-chaff (White)	..	..	..	..	..	..	..	..	..
25. <i>Belgian (White)</i>	..	..	..	..	..	..	..	..	..
Means	32½	42½	38½	50½	36½	42½	42½	42½	41½

WEIGHT PER BUSHEL. Lbs.

	..	..	..	..	..	60½	Means	Mean.
1. White-chaff (Red)	58½	61½	61	63	60½	60½	61	61
2. Rivett's (Red)	57½	58½	58½	59	60½	58½	58½	58½
3. Chubb Wheat (Red)	59½	61½	60½	62½	60½	60½	61½	61½
4. Red-chaff (White)	62½	61½	61½	63½	60½	60½	61½	61½
5. Brownick (Red)	59½	61½	59½	62½	60½	60½	61½	61½
6. Red Wonder	60	62½	60½	63	61½	63	62½	62½
7. Burwell (Old Red Lammas) ..	61½	63½	61½	64½	61½	64	64	64
8. Bristol Red	60½	61½	60½	62½	59½	63½	63½	63½
9. Red Nursery	62	65½	62½	66	58½	62½	62½	62½
10. Red Langham	60½	63	60½	63½	59½	61½	61½	61½
11. Woolly Ear (White)	61½	62½	57½	63½	59½	62½	62½	62½
12. Hardcastle (White)	63	63	59½	64½	61½	63½	63½	63½
13. Golden Drop (Red), Hallett's ..	61½	62½	61½	63½	61	61½	61½	61½
14. Victoria White, Hallett's ..	62½	61½	60½	63½	59½	62½	62½	62½
15. Hunter's White, Hallett's ..	59½	61½	58½	62½	59	59	59	59
16. Original Red, Hallett's ..	58½	60½	61½	64½	61½	61½	61½	61½
17. White Childam	62½	62½	59½	63½	59½	60½	60½	60½
18. Red Rostock	56½	59½	60	63½	59½	60½	60½	60½
19. Casey's White	58½	60½	61½	63½	59½	60½	60½	60½
20. Golden Rough-chaff (Red) ..	59½	62½	65½	65½	60½	61½	61½	61½
21. Bole's Prolific (Red)	57½	62	60½	63½	59½	62½	62½	62½
22. Club Wheat (Red)	58½	61½	61½	63½	59½	61½	61½	61½
23. Main's Standing White	..	..	..	..	..	..	..	..
24. Main's Rough-chaff (White) ..	..	..	..	..	..	..	..	..
25. <i>Bulgarian (Wheat)</i>	..	..	..	..	..	56½	56½	56½
Means	59½	61½	60½	63½	60½	60½	61	61

ROTHAMSTED

MAY,

SUMMARY STATEMENT OF THE PRESENT AND PREVIOUS

(14 Years, 1865-1878,

PREVIOUS CROPPING

Name of Field.	Acres.	1865.	1866.	1867.	1868.	1869.	1870.	1871.	1872.
Thirty Acres	30	Oats, 1 cwt. Guano, 3 cwt. Corn Manure.	Tares and Swedes, Dung and Artificial.	Oats, after Sheep-Folding.	Clover.	Wheat, 2 cwt. Guano.	Oats, 2 cwt. Guano.	Barley, 2 cwt. superphos., 2 cwt. Nitrate Soda.	Barley, 2½ cwt. superphos., 2½ cwt. Nitr. Soda, (2½ acres experiment).
Harpenden	22	Wheat, Sheep-Folded.	Red Clover (peren.), Unmanured.	Wheat, 2½ cwt. Guano.	Oats, 2 cwt. Guano, & 1 cwt. Nitr. Soda. 1rd { 1 cwt. Nitr. Soda. and Sheep-folded.	Swedes, Dung and various Artificial Manures.	Wheat, 3 cwt. Guano.	Oats, 3 cwt. Guano, 1 cwt. Nitrate Soda. Tares, Dung.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda, Tares, Dung.
Little Hoos	9	Wheat, 1½ cwt. Guano, 1 cwt. Nitrate Soda, 1 cwt. Corn Manure.	Mangolds, Dung and Artificial.	Wheat, Unmanured.	Oats, 2 cwt. Guano, 1 cwt. Nitrate of Soda.	Barley, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia, 1 cwt. superphosphate.	Barley, 2½ cwt. Guano.	Barley, 3 cwt. superphos., 2½ cwt. Nitrate Soda.	Barley (with Clover). 2½ cwt. superphos., 2½ cwt. Nitr. Soda.
Fosters'	18	Oats, 1 cwt. Guano, 1 cwt. Corn Manure.	Red Clover, Unmanured.	Wheat, 2 cwt. Guano, ½ cwt. Corn Manure.	Oats, 2 cwt. Guano, 1 cwt. Nitrate of Soda.	Barley, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia, 1 cwt. superphosphate.	Oats, 2 cwt. Guano, 3 cwt. Blood Manure.	Roots, Tares, and Rape, Dung and Artificial.	Wheat, Varieties of Wheat, 2 cwt. superphos., 2 cwt. Nitr. Soda, ½ Sheep-folded.
Knott Wood	30	Wheat, Sheep-Folded, 1 cwt. Guano.	Oats, 2 cwt. Guano, 1 cwt. Sulph. Ammonia.	Oats, 2 cwt. Guano, 1 cwt. Sulph. Ammonia.	Swedes, 2 cwt. Guano, 2½ cwt. superphosphate and Dung.	Wheat, 3 cwt. Guano (one-half), Unmanured (one-half), after Swedes ploughed up and Fallowed.	Oats, 3 cwt. Guano.	Oats, 3 cwt. Guano, 1 cwt. Nitrate Soda.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.
Little Knott Wood	14	Red Clover (peren.), Unmanured.	Red Clover (peren.), Sheep-Folded.	Wheat, 1 cwt. Guano, ½ cwt. Corn Manure.	Oats, 2 cwt. Guano, 1 cwt. Nitrate Soda.	Mangolds, 12 tons Dung, 3 cwt. Guano.	Wheat, 3 cwt. Guano.	Oats, 3 cwt. Guano, 1 cwt. Nitrate Soda.	Oats, ½ Sheep-folded. All, 2½ cwt. super., 2½ cwt. Nitr. Soda.
Sawpit	14	Mangolds and Turnips, Dung and Artificial.	Wheat, Unmanured.	Red Clover, Unmanured.	Wheat, 1 cwt. Guano, 1 cwt. Wheat Manure.	Wheat, 3 cwt. Guano.	Mangolds, Dung and 3 cwt. Guano.	Wheat, 3 cwt. Guano.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.
Rick-yard	8	Barley, 2 cwt. Guano, 1½ cwt. Corn Manure.	Red Clover, Sheep-Folded.	Wheat, Guano.	Barley, 2 cwt. Wheat Manure.	Tares, Dung.	Barley, 1 cwt. Guano.	Mangolds, Dung and 4 cwt. Cotton Cake.	Wheat, Unmanured.
Six Acres	6	Red Clover, Unmanured.	Wheat, 2 cwt. Guano, 2 cwt. Corn Manure.	Oats, 3 cwt. Guano.	Beans, Dung.	Wheat, 2 cwt. Guano, 1 cwt. Nitrate of Soda.	Barley, 2½ cwt. Guano.	Barley, 3 cwt. superphos., 2½ cwt. Nitrate Soda.	Barley, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.
Clay-Croft	12	Oats, 2 cwt. Guano, 2 cwt. Corn Manure.	Oats, 2 cwt. Guano, 1 cwt. Sulph. Ammonia.	Beans, Dung.	Wheat, 2 cwt. Guano.	Oats, 2 cwt. Guano, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia.	Turnips, Dung and 3 cwt. super- phosphate.	Wheat, Unmanured.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.
Ten Acres	10	Tares, Dung.	Turnips, Artificial.	Wheat, Guano.	Red Clover.	Wheat, 2 cwt. Guano.	Oats, 3 cwt. Guano.	Mangolds, Dung and 4 cwt. Cotton Cake.	Wheat, Unmanured.
Agdell	9	Red Clover, Unmanured.	Wheat, 1½ cwt. Guano, 1½ cwt. Corn Manure.	Oats, 2 cwt. Guano.	Tares, Dung.	Barley, Unmanured.	Barley, 1½ cwt. Guano, 1½ cwt. super- phosphate.	Mangolds, Dung and 4 cwt. Cotton Cake.	Wheat, Unmanured (and part Roots).
Long Hoos	25	Barley, 1 cwt. Guano, 1 cwt. Corn Manure.	Barley, 1½ cwt. Guano, 1 cwt. Corn Manure.	Mangolds and Swedes, 15 tons Dung, 3 cwt. Guano.	Wheat, 1 cwt. Guano.	Oats, 2 cwt. Guano, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia.	Sainfoin, Unmanured.	Sainfoin, Unmanured. (Steam cultivated, July.)	Mangolds, Dung, (Carted off.)
Sawyers'	25	Swedes, Dung and Artificial.	Wheat and Barley, Sheep-Folded.	Red Clover, Unmanured.	Wheat, 3 cwt. Guano.	Fallow.	Wheat, 4 cwt. Guano.	Wheat, 4 cwt. Guano, 1 cwt. Nitrate Soda.	Barley, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.
West Barn	32	Red Clover (peren.), Sheep-Folded.	Wheat, 1½ cwt. Guano, 1½ cwt. Corn Manure.	Barley, 1 cwt. Blood Manure, 1 cwt. superphosphate, 1 cwt. Sulph. Ammonia.	Fallow.	Wheat, 3 cwt. Guano.	Sainfoin, Unmanured.	Sainfoin, Unmanured.	Sainfoin, Unmanured.

FARM.

1878.

CROPPING, &c., OF THE ARABLE LAND NOT UNDER EXPERIMENT.

inclusive.)

AND MANURING.

1873.	1874.	1875.	1876.	1877.	Crops, &c., Present Season, 1877-78.	Acres.	Name of Field.
Barley ($\frac{3}{4}$ with Grass-seeds). 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Grass ($\frac{3}{4}$). Folded, and 1 cwt. Nitrate. Barley ($\frac{1}{4}$). 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Grass ($\frac{3}{4}$). Sheep-folded. Tares ($\frac{1}{4}$). Dung.	Grass ($\frac{3}{4}$). Compost. Wheat ($\frac{1}{4}$). 1 cwt. Nitrate Soda.	Grass ($\frac{3}{4}$). Cattle Grazed. Barley ($\frac{1}{4}$). 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Grass ($\frac{3}{4}$). Cattle Grazed with Cotton-Cake. Tares ($\frac{1}{4}$). Dung.	30	Thirty Acres
Barley, After Oats—2 cwt. super- phosphate; 2 cwt. Nitrate. After Tares—1 cwt. super- phosphate; 1 cwt. Nitrate.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Mangolds, Dung, and 2 cwt. Guano. (Carted off.)	Wheat (Varieties). 2 cwt. Nitrate Soda.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Barley (with Clover). 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	22	Harpenden.
Barley ($\frac{1}{4}$). Unmanured. Clover ($\frac{1}{4}$). Unmanured.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda (1 acre Unmanured).	Barley, where Barley 1873, 2 cwt. superphosphate, 2 cwt. Nitrate of Soda. where Clover 1873, Half quantities.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda ($\frac{1}{4}$ with Clover).	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda ($\frac{1}{4}$ with Clover).	Barley ($\frac{1}{4}$). 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda. Clover ($\frac{1}{4}$). Unmanured.	9	Little Hoos.
Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda (2 acres experiment).	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, (1) 3 $\frac{1}{2}$ cwt. Guano, (4) 2 $\frac{1}{2}$ cwt. superphosphate, (4) 1 $\frac{1}{2}$ cwt. Guano, 1 $\frac{1}{2}$ Nitrate.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	White Turnips, Dung. Superphosphate, $\frac{1}{2}$ cwt. Nitrate Soda; part fed, part carted.	Wheat (Varieties). 2 cwt. Nitrate Soda.	18	Fosters'.
Tares ($\frac{1}{4}$). Dung. Swedes ($\frac{1}{4}$). Dung, 2 cwt. superphosph. 2 cwt. Nitrate Soda.	Barley, After Roots and Tares carted, 2 cwt. superphosphate, 2 cwt. Nitrate Soda, After Tares fed, 1 cwt. each.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Oats, 2 $\frac{1}{2}$ cwt. superphosphate, 3 cwt. Nitrate Soda.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Roots ($\frac{1}{4}$). Dung and Artificial. Fallow ($\frac{1}{4}$).	30	Knott Wood.
Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Mangolds, Dung. (Carted off.)	Wheat (Varieties). 1 $\frac{1}{2}$ cwt. Nitrate Soda.	Oats, 2 $\frac{1}{2}$ cwt. superphosphate, 3 cwt. Nitrate Soda.	Oats (with Clover). 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Clover, Unmanured.	14	Little Knott Wood.
Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Mangolds, 25 tons Dung. (Carted off.)	Wheat (Varieties). 1 $\frac{1}{2}$ cwt. Nitrate Soda.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	14	Sawpit.
Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Tares, Dung. $\frac{1}{4}$ followed by Turnips, 1 cwt. superphosphate, 1 cwt. Nitrate Soda.	Barley, 1 cwt. Nitrate Soda.	Swedes, Dung, and Superphosphate.	Barley, 1 cwt. Nitrate Soda.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	8	Rick-yard.
Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Barley (with Clover). 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Clover, Unmanured.	6	Six Acres.
Clover, Unmanured.	Wheat, 2 cwt. Nitrate Soda.	Oats, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Oats, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Fallow.	Wheat, 2 cwt. Nitrate Soda.	12	Clay-Croft.
Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda (5 acres experiment).	Oats, 2 cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Oats, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Fallow.	Wheat (with Clover). 2 cwt. Nitrate Soda.	Clover, Unmanured.	10	Ten Acres.
Clover, Unmanured. Barley, Experiment.	Wheat, 1 cwt. Nitrate Soda (3 acres Experiment, $\frac{1}{4}$ Clover, $\frac{1}{4}$ Fallow).	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda. Wheat, 3 acres, Experiment.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 3 cwt. Nitrate Soda. Swedes, 3 acres, Experiment.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda. Barley, 3 acres experiment.	Potatoes. Dung and Artificial. (3 acres experiment $\frac{1}{2}$ Beans, $\frac{1}{2}$ Fallow.)	9	Agdell.
Wheat. (1 Varieties of Wheat), 1 $\frac{1}{2}$ cwt. Nitrate Soda.	Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Oats, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Oats ($\frac{3}{4}$). 2 $\frac{1}{2}$ cwt. superphosphate, 3 cwt. Nitrate Soda. Tares ($\frac{1}{4}$). Dung.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	Barley, 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda.	25	Long Hoos.
Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Mangolds and Swedes, Dung.	Barley after Swedes ($\frac{3}{4}$). 2 cwt. superphosphate, 2 cwt. Nitrate Soda. Wheat after Mangolds ($\frac{1}{4}$). 1 $\frac{1}{2}$ cwt. Nitrate Soda.	Barley (with Clover). 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley ($\frac{1}{4}$). 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda. Tares ($\frac{1}{4}$). Dung.	Barley, ($\frac{3}{4}$) 2 $\frac{1}{2}$ cwt. superphosphate, 2 $\frac{1}{2}$ cwt. Nitrate Soda, ($\frac{1}{4}$) 2 $\frac{1}{2}$ cwt. Nit. Soda alone.	25	Sawyers'.
Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Wheat (Oats fed off 1873), 1 $\frac{1}{2}$ cwt. Nitrate Soda.	Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Oats, 2 cwt. superphosphate, ($\frac{3}{4}$) 1 $\frac{1}{2}$ Nitrate Soda, ($\frac{1}{4}$) 2 $\frac{1}{2}$ Nitrate Soda.	Fallow.	Wheat, 2 cwt. Nitrate Soda.	32	West Barn.