


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1939-1947 Volume 2



[Full Table of Content](#)

Q Carrots - Including Carrot Series

Rothamsted Research

Rothamsted Research (1948) *Q Carrots - Including Carrot Series* ; Yields Of The Field Experiments 1939-1947 Volume 2, pp 163 - 165 - DOI: <https://doi.org/10.23637/ERADOC-1-186>

Q/1

CARROTS

Woburn Lansome 1939 (6th year)

The effects of sulphate of ammonia, poultry manure, soot, rape dust and dung.

Design; 4 randomized blocks of 12 plots each.

Area of each plot; 0.00588 acre.

Treatments

None: Sulphate of ammonia at 0.4 and 0.8 cwt. N per acre (half in seed bed and half as top dressing) Poultry manure, soot and rape dust all at 0.8 cwt. N per acre: dung ploughed-in at 0.8 and 1.6 cwt. N per acre every second year including 1939, and every second year excluding 1939, and at 0.8 cwt. N per acre every year.

Basal manuring; superphosphate and muriate of potash applied to give a total of 1.0 cwt. P_2O_5 and 1.0 cwt. K_2O , per acre, including the P_2O_5 and K_2O in the organic fertilizer.

Crop Notes

Sown: May 22. Harvested: Oct. 27 and Nov. 11. Variety, Cooper's Early Market. Previous crop, kale (see 1938 Station Report, p.164).

Standard error per plot: 2.41 tons per acre or 20.5%, (34 d.f.)

Mean Yields: tons per acre

None	Sulphate of ammonia		Poultry Manure	Soot	Rape dust	0.8 cwt. N per acre
	0.4	0.8	0.8	0.8	0.8	
±0.852			±1.20			
10.56	11.04	11.88	13.83	12.92	13.03	
Dung						
1938	0.8	1.6	0.8	0	0	0 cwt. N
1939	0	0	0.8	0.8	1.6	per acre
			±1.20			Mean of all plots
	10.19	11.35	11.74	11.36	12.45	11.74

Q/2

CARROTS

Woburn - Lansome 1941

Woburn - Butt Close 1942

Effects of sulphate of ammonia, superphosphate, muriate of potash and agricultural salt. These two experiments form part of a country wide series of 25 similar experiments.

Design; 4 randomized blocks of 8 plots each, the third order interaction being confounded with block differences.

Area of each plot, 0.00875 acre.

Treatments

Sulphate of ammonia:	1941, None, 0.4 cwt. N per acre.
	1942, None, 0.3 cwt. N per acre.
Superphosphate:	1941, None, 0.8 cwt. P ₂ O ₅ per acre.
	1942, None, 0.6 cwt. P ₂ O ₅ per acre.
Muriate of potash:	1941, None, 1.0 cwt. K ₂ O per acre.
	1942, None, 0.9 cwt. K ₂ O per acre.
Agricultural salt:	1941 and 1942: None, 3.0 cwt. per acre.

Crop Notes

	Seed sown	Harvested	Variety	Previous crop
1941	June 23	Oct. 21, 31, Nov. 4, 12*	Cooper's Early Market	Oats
1942	April 14	Aug. 27, Sept. 3. Oct. 20 - Nov. 12	Cooper's Intermediate	Sugar beet

*On each day a strip was taken through all plots.

Standard errors per plot:

1941 Total roots:	0.655 tons per acre or 12.6%, 12 d.f.
1942 1sts:	5.39 tons per acre or 23.5%, 14 d.f.
2nds:	0.590 tons per acre or 52.5%, 14 d.f.

Note: 2nds consisted mostly of splits together with the smalls.

Q/3

1941 Differential responses; tons per acre

	Mean	Sulph. amm.		Super		Mur. Pot.		Salt	
		Abs.	Pres.	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Total Roots. Mean yield: 5.21 tons per acre									
	±0.231				±0.327				
Sulph. amm.	2.11	-	-	2.42	1.80	2.32	1.90	1.92	2.30
Super	0.11	0.42	-0.20	-	-	0.25	-0.03	0.57	-0.35
Mur. pot.	-0.22	-0.01	-0.43	-0.08	-0.36	-	-	-0.40	-0.04
Salt	0.38	0.19	0.57	0.84	-0.08	0.20	0.56	-	-

1942

1sts: Mean yield, 22.92 tons per acre

	±1.91				±2.70				
Sulph. amm.	0.52	-	-	-1.20	2.24	4.17	-3.13	-0.35	1.39
Super.	0.11	-1.61	1.83	-	-	-2.40	2.62	2.30	-2.08
Mur. pot.	1.51	5.16	-2.14	-1.00	4.02	-	-	1.29	1.73
Salt	4.22	3.35	5.09	6.41	2.03	4.00	4.44	-	-

2nds: Mean yield, 1.12 tons per acre

	±0.209				±0.296				
Sulph. amm.	0.14	-	-	0.05	0.23	0.18	0.10	-0.13	0.41
Super.	-0.12	-0.21	-0.03	-	-	-0.11	-0.13	0.22	-0.46
Mur. pot.	-0.03	0.01	-0.07	-0.02	-0.04	-	-	-0.10	0.04
Salt	0.42	0.15	0.69	0.76	0.08	0.35	0.49	-	-