


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1948

[Full Table of Content](#)



48/CC/1 Beans - Cultivations and Varieties

Rothamsted Research

Rothamsted Research (1949) *48/CC/1 Beans - Cultivations and Varieties* ; Yields Of The Field Experiments 1948, pp 74 - 76

48/Cc/1.1

BEANS

The comparison of four varieties of beans and the effects on them of two methods of placement, two rates and two times of sowing.

RE - Great Field I 1948

System of replication: 4 randomized blocks of 8 plots each, the effect of time of sowing and certain high order interactions being confounded with block differences.

Area of each plot: 0.0286 acre.

Treatments:

Varieties: Giant, Essex, Lines and Wilts.

Methods of placement: Seed broadcast before ploughing, dropped in furrow during ploughing.

Rate of sowing: 2 cwt, 3 cwt seed per acre.

Time of sowing: Early, Oct 28th. Late, Nov 18th.

Basal dressings: Dung: 10 tons per acre

Superphosphate 2 cwt per acre drilled across the plots

Muriate of potash 2 cwt per acre

Nitrochalk: 2 cwt per acre.

Cultivations etc: Superphosphate and Muriate of Potash applied: Oct 20.

Dung applied, ploughed in, beans placed in furrows, 1st sowing "Early":

Oct 27-29. Tractor rolled and disced each way after ploughing: Oct 29.

Chalk 45 cwt per acre applied: Nov 6. Dung spread, ploughed in beans

placed in furrows, 2nd sowing "Late": Nov 17-18. Harrowed "Early"

plots: Mar 16. Ring rolled all plots: Mar 30. Harrowed late plots:

Mar 31. Nitrochalk applied: May 4. Harvested: Aug 25. Previous

crop: Linseed.

Standard errors per plot :

Grain: 2.62 cwt per acre or 10.7% (11 d.f.)

Straw: 4.40 cwt per acre or 11.2% (11 d.f.)

4.8/Cc/1.2

Grain: cwt per acre

	Giant	Essex	Lincoln -shire	Wilts	Mean
(± 0.92) Mean	23.8	26.5	24.4	23.2	24.5
(± 1.31) Broadcast	23.2	25.3	24.4	24.1	24.3
Ploughed In	24.3	27.7	24.3	22.3	24.6 (± 0.65)
(± 1.85) Ploughed In -Broadcast	1.1	2.4	-0.1	-1.8	0.3 (± 0.92)
Seed rate per acre					
(± 1.31) 2 cwt	21.5	24.1	22.8	22.1	22.6 (± 0.65)
3 cwt	26.1	28.9	26.0	24.3	26.3 (± 0.65)
(± 1.85) 3 cwt - 2 cwt	4.6	4.8	3.2	2.2	3.7 (± 0.92)
(± 1.85) [*] Early-Late Sowing	4.1	8.0	7.2	6.0	

	Seed rate per acre		Difference	Early- Late Sowing
	2 cwt	3 cwt		
(± 0.92) Broadcast	22.4	26.1	3.7 (1)	7.1
Ploughed in	22.8	26.5	3.7 (1)	5.5
(± 1.31) Difference	0.4	0.4	0.0 (2)	-1.6 (2)
Early-Late Sowing	7.3	5.3	-2.0 (2)	

Standard errors (1) ± 1.31 (2) ± 1.85 ^{*} Standard error for comparison between main effects only

48/Cc/1.3

Straw: cwt per acre

	Giant	Essex	Lincoln -shire	Wilts	Mean
(±1.56) Mean	39.6	38.5	38.7	40.3	39.3
(±2.20) Broadcast	39.3	36.8	39.8	40.2	39.0 (±1.10)
Ploughed In	39.9	40.3	37.7	40.4	39.6 (±1.10)
(±3.11) Ploughed In -Broadcast	0.6	3.5	-2.1	0.2	0.6 (±1.56)
Seed rate per acre					
(±2.20) 2 cwt	36.4	34.9	36.9	37.3	36.4 (±1.10)
3 cwt	42.8	42.1	40.6	43.3	42.2 (±1.10)
(±3.11) 3 cwt - 2 cwt	6.4	7.2	3.7	6.0	5.8 (±1.56)
(±3.11)* Early-Late Sowing	7.5	8.2	9.2	11.6	

	Seed rate per acre 2 cwt	3 cwt	Difference	Early- Late Sowing
(±1.56) Broadcast	36.7	41.3	4.6(1)	10.8
Ploughed In	36.0	43.1	7.1(1)	7.4
(±2.20) Ploughed In -Broadcast	-0.7	1.8	2.5(2)	-3.4(2)
Early-Late Sowing	8.9	9.3	0.4(2)	

Standard Errors (1) (±2.20)

(2) (±3.11)

* Standard error for comparison between main effects only