


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1950

[Full Table of Content](#)



50/CH/1 Permanent Grass - Fertilizer Placement - Rothamsted

Rothamsted Research

Rothamsted Research (1951) *50/CH/1 Permanent Grass - Fertilizer Placement - Rothamsted* ; Yields Of The Field Experiments 1950, pp 83 - 84 - DOI: <https://doi.org/10.23637/ERADOC-1-185>

50/Ch/1.1

PERMANENT GRASS

Fertilizer placement - Highfield 1950

System of replication: 3 randomized blocks of 6 plots each

Area of each plot: 0.0137 acre

Treatments: No fertilizer; no fertilizer but plots drilled over;
and all combinations of:-

Compound granular PK fertilizer (13% P_2O_5 , 15% K_2O): 3.2, 6.4
cwt per acre.

Method of placement: Broadcast; drilled in bands 10" apart and
3" deep.

Basal manuring: 3 cwt sulphate of ammonia per acre

Cultivations, etc.:

Fertilizer broadcast and drilled: Apr 4. Sulphate of ammonia
applied: Apr 6. 1st cut: June 8. 2nd cut: Aug 2.

Standard errors per plot:

Dry matter, 1st cut: 2.26 cwt per acre or 8.4% (10 d.f.)
2nd cut: 1.36 cwt per acre or 11.4% (10 d.f.)

50/Ch/1.2

Summary of Results

Grass Dry Matter: cwt per acre

Method of Placement	Compound Fertilizer cwt per acre			Mean	Difference of levels
	None	1st Cut 3.2	6.4		
		(± 1.31)		(± 0.92)	(± 1.85)
Broadcast	25.2 ⁺	29.7	28.9	29.3*	-0.8
Drilled	23.5 ⁺	25.4	28.0	26.7*	2.6
Mean (± 0.92)	24.4	27.6	28.5	26.8	0.9
Difference (± 1.85)	-1.7	-4.3	-0.9	-2.6* (± 1.31)	(± 1.31)
2nd Cut					
		(± 0.79)		(± 0.56)	(± 1.11)
Broadcast	11.7 ⁺	11.8	12.5	12.1*	0.7
Drilled	11.8 ⁺	12.2	11.7	11.9*	-0.5
Mean (± 0.56)	11.7	12.0	12.1	11.9	0.1
Difference (± 1.11)	0.1	0.4	-0.8	-0.2* (± 0.79)	(± 0.79)

⁺ see Treatment descriptions.

* excluding no fertilizer.