


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1951

[Full Table of Content](#)



51/CG/1 Lucerne - Fertilizer Placement - Rothamsted

Rothamsted Research

Rothamsted Research (1952) *51/CG/1 Lucerne - Fertilizer Placement - Rothamsted* ; Yields Of The Field Experiments 1951, pp 86 - 87 - DOI: <https://doi.org/10.23637/ERADOC-1-171>

51/Cg/1.1

LUCERNE

Fertilizer placement - Long Hoos IV 1951

System of replication: 3 randomized blocks of 12 plots each.

Area of each plot: 0.00798 acre. Area harvested: 0.00644 acre.

Treatments - applied 1950 and 1951: No fertilizer (duplicate plots); no fertilizer but plots drilled over both in 1950 and 1951 (duplicate plots); and all combinations of:-
Compound granular fertilizer (10% P_2O_5 , 20% K_2O): 2.5; 5.0 cwt per acre.

Method of placement: Broadcast during preparation of seed bed 1950; drilled in bands 3" below surface and 2" to side of seed 1950; half broadcast during preparation of seed bed 1950, half broadcast on surface 1951; half drilled beside seed 1950, half drilled beside rows of plants 1951.

Basal manuring: None.

Cultivations, etc.: Fertilizers applied: Mar 30. 1st cut: June 20.
2nd cut: July 31. 3rd cut: Sept 26. Variety: Du Puits.

Standard errors per plot:

Dry matter, 1st cut: 2.95 cwt per acre or 5.6% (24 d.f.)
2nd cut: 2.37 cwt per acre or 11.2% (24 d.f.)
3rd cut: 2.41 cwt per acre or 9.4% (24 d.f.)

See. 1950 edition p.p. 50/Cg/1.1-1.2

51/Cg/1.2

Summary of Results

Dry matter: cwt per acre

Compound fertilizer cwt per acre	Method of placement				Mean
	Broadcast In seedbed '50	Drilled 3" below sur- face, 2" to side of seed	Broadcast $\frac{1}{2}$ on seedbed $\frac{1}{2}$ on sur- face '51	Drilled $\frac{1}{2}$ beside seed $\frac{1}{2}$ beside rows '51	
1st cut					(± 0.85)
None (± 1.20)	51.6 [†]	50.0 [†]			50.8
2.5 (± 1.70)	53.3	55.1	53.3	50.6	53.1
5.0 (± 1.70)	52.6	56.1	48.9	54.1	52.9
Mean (± 1.20)	53.0*	55.6*	51.1	52.3	52.3
2nd cut					(± 0.68)
None (± 0.97)	19.9 [†]	22.6 [†]			21.2
2.5 (± 1.37)	21.1	22.0	19.8	21.8	21.2
5.0 (± 1.37)	21.3	20.5	19.5	22.5	20.9
Mean (± 0.97)	21.2*	21.2*	19.6	22.2	21.1
3rd cut					(± 0.70)
None (± 0.98)	24.7 [†]	25.9 [†]			25.3
2.5 (± 1.39)	26.3	25.4	24.0	25.1	25.2
5.0 (± 1.39)	26.3	26.6	26.9	24.9	26.2
Mean (± 0.98)	26.3*	26.0*	25.5	25.0	25.6

† See treatment description

* Excluding 'None'.

Mean Dry Matter %.

1st cut: 22.0

2nd cut: 25.2

3rd cut: 19.8