


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1952

[Full Table of Content](#)



52/CB/2 Barley - Nitrophosphates - Rothamsted

Rothamsted Research

Rothamsted Research (1953) *52/CB/2 Barley - Nitrophosphates - Rothamsted* ; Yields Of The Field Experiments 1952, pp 79 - 79 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

52/Cb/2

BARLEY

Nitrophosphates placement - Highfield 3 1952.

System of replication: 4 randomized blocks of 6 plots each.

Area of each plot (average): 0.0143 acre.

Treatments: Nitrochalk at 0.5 cwt N per acre broadcast (two plots per block), and all combinations of:-

Manures: Nitrophosphate (British) 13.9% N, 14.6% P₂O₅; Nitrochalk and Granular Superphosphate. Each manure provided 0.5 cwt N and 0.53 cwt P₂O₅ per acre.

Method of placement: Broadcast; Combine drilled.

Basal dressing: None.

Cultivations, etc.: Ploughed: Sept 18. Seed drilled at 3½ bushels per acre and all fertilizers applied: Mar 15. Harvested: July 30. Variety: Plumage Archer. Previous crop: Linseed.

Standard errors per plot: Grain.

Yield: 2.37 cwt per acre or 11.0% (16 d.f.)

P₂O₅ uptake: 0.0180 cwt per acre or 16.0% (16 d.f.)Summary of Results

	Nitrochalk broadcast	Nitrophosphate broadcast	combine drilled	Nitrochalk and superphosphate broadcast	combine drilled	Mean
Yield: cwt per acre						
Grain (±1.18)	18.9 ⁽¹⁾	21.9	24.4	22.0	22.6	21.4
Increase (±1.45)		3.0	5.5	3.1	3.7	
Straw	32.6	30.5	36.7	33.2	32.9	33.1
Increase		-2.1	+4.1	+0.6	+0.3	
P ₂ O ₅ uptake: cwt per acre						
Grain (±0.0090)	0.098 ⁽²⁾	0.115	0.132	0.117	0.114	0.112
Increase (±0.0110)		0.017	0.034	0.019	0.016	
Straw	0.028	0.023	0.030	0.030	0.026	0.027
Increase		-0.005	+0.002	+0.002	-0.002	

(1) ±0.84
(2) ±0.0064