


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1953

[Full Table of Content](#)



53/CH/1 Broccoli - Control of Virus - Rothamsted

Rothamsted Research

Rothamsted Research (1954) *53/CH/1 Broccoli - Control of Virus - Rothamsted* ; Yields Of The Field Experiments 1953, pp 103 - 103 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/Ch/1.1

CARROTS

Residual effects of Krilium - Rothamsted, Fosters and Barnfield;
Woburn, Stackyard and Warren Field.

System of replication: 4 x 4 Latin Square.

Area of each plot: 0.00207 acre. Area harvested: 0.00138 acre.

Treatments, applied 1952:

Krilium per acre: None; 3 cwt broadcast and twice rotary cultivated
in; 6 cwt broadcast and twice rotary cultivated in;
3 cwt broadcast and raked into seed bed.

Basal dressing: 5 cwt compound fertilizer ($7\frac{1}{2}\%$ N, $7\frac{1}{2}\%$ P_2O_5 , $10.5\frac{1}{2}\%$ K_2O).

Cultivations, etc.:

Fosters: Ploughed: Mar 13. Roto tilled: Mar 13. Basal dressing
applied, seed drilled: Apr 22. Lifted: Sept 24 and Oct 23.

Barnfield: Ploughed: Jan 28. Basal dressing applied, seed
drilled: Apr 21. Lifted: Oct 20 and 26.

Stackyard: Hand dug: Jan 23. Seed sown: May 5. Basal dressing
applied: May 14. Lifted: Sept 22.

Warren Field: Hand dug: Jan 27. Seed sown: May 6. Basal
dressing applied: May 15. Singled: June 26. Lifted: Oct 5.

All fields, Variety: James' Scarlet Intermediate. Previous crop:
Clover, Carrots, Ryegrass and Red Beet in strips on all plots.

Standard errors per plot. Roots tons per acre:

Fosters:	0.995	tons	per	acre	or	4.5%	(6 d.f.)
Barnfield:	1.41	"	"	"	"	11.1%	(6 d.f.)
Stackyard:	1.46	"	"	"	"	17.2%	(6 d.f.)
Warren Field:	0.573	"	"	"	"	7.7%	(6 d.f.)