


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1953

[Full Table of Content](#)



53/A/3 Agdell - Rotation - Rothamsted

Rothamsted Research

Rothamsted Research (1954) *53/A/3 Agdell - Rotation - Rothamsted* ; Yields Of The Field Experiments 1953, pp 6 - 6 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1953

Without manure 1851 and since

For history, details of treatments etc. see "Results of the Field Experiments 1939-47", Vol. I, Section A/3.

Cultivations, etc.:

Cropped plots. Ploughed: Sept 6, 1952. Seed drilled at 3 bushels per acre: Oct 27. Harvested: Aug 20, 1953. Variety: Squareheads Master 13/4.

Fallow plots. Ploughed: Sept 6, 1952 and Sept 3, 1953.

Summary of Results

Plot No. of years of fallow	Mean yields: cwt per acre			Mean
	B ₂	B ₃	B ₄	
	1	1	3	
Grain	18.7	8.3	13.7	13.6
Straw	24.9	14.5	23.6	21.0

CROPS IN ROTATION - AGDELL 1953

Barley - 2nd crop of 27th course 1952-55

For history, details of treatments etc. see "Results of the Field Experiments 1939-47", Vol. I, Section A/4.

Cultivations, etc.: Ploughed: Sept 5, 1952, Seed drilled at 3 bushels per acre: Mar 2, 1953. Sprayed with MCPA, 2 pints per acre: May 26. Harvested: Aug 28. Variety: Plumage Archer.

Note: No turnips were grown and no manures were applied in 1952, the plots being fallowed.

Summary of Results

Manure to Turnips 1948	Mean yields: cwt per acre					
	None since 1848		Mineral manure No nitrogen		Complete mineral and nitrogenous manure	
	Fallow	Clover	Fallow	Clover	Fallow	Clover
Rotation Plot	5	6	3	4	1	2
Grain	5.7	11.1	7.6	16.2	3.6	5.0
Straw	5.8	11.5	8.3	16.4	4.8	8.9