


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1953

[Full Table of Content](#)



53/CA/4 Wheat - Residual of Dung - Rothamsted

Rothamsted Research

Rothamsted Research (1954) *53/CA/4 Wheat - Residual of Dung - Rothamsted* ; Yields Of The Field Experiments 1953, pp 74 - 74 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/Ca/4

WHEAT

Residual effect of dung - West Barnfield II 1953.

System of replication: 4 randomized blocks of 12 plots each.

Area of each plot: 0.0318 acre. Area harvested: 0.0129 acre.

Treatments: All combinations of:-

Dung: None; 5; 10; 15 tons per acre applied to potatoes 1951-52.

Methods of application: Ploughed in, in winter (1951); ploughed in, in spring (1952); placed in ridges (1952).

Basal dressing: 3 cwt sulphate of ammonia per acre as top dressing in spring.

Cultivations, etc.: Cultivated: Oct 22 and again Oct 30 1952. Seed drilled at 3 bushels per acre: Nov 4. Sulphate of ammonia applied: April 24, 1953. Sprayed with M.C.P.A. low volume, 2 pints per acre: May 21. Combine harvested: Aug 21. Variety: Cappelle. Previous crop: Potatoes.

Standard error per plot:

Grain*: 5.06 cwt per acre or 12.4% (35 d.f.)

Summary of Results

Method of application	Dung applied to potatoes 1951-52: tons per acre				Mean	
	None	5	10	15		
Grain*: cwt per acre						
		(±2.53)			(±1.46)	
Ploughed in, in winter		40.6	42.8	42.3	41.9	
Ploughed in, in spring		41.9	39.3	45.4	42.2	
Placed in ridges		37.7	41.4	42.8	40.6	
Mean (±1.46)		38.6	40.1	41.2	43.5	40.8

*Corrected to 85% D.M. Mean D.M. %: 84.3.

N.B. For details of original experiment see "Results of the Field Experiments 1952", Section Cc/1.