


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1953

[Full Table of Content](#)



53/CB/4 Barley - Methods of Harvesting - Rothamsted

Rothamsted Research

Rothamsted Research (1954) *53/CB/4 Barley - Methods of Harvesting - Rothamsted* ; Yields Of The Field Experiments 1953, pp 83 - 83 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/Cb/4

BARLEY

Methods of harvesting - Deacons Field 1953.

System of replication: 12 randomized blocks of 2 plots each.

Area of each plot: 0.0200 acre. Area harvested: Binder - full area;
Combine harvester - 0.0129 acre.

Treatments:

Methods of harvesting: Binder; Combine harvester (Massey Harris 726
8'6" cut).

Basal dressing per acre: 1 cwt superphosphate; $2\frac{1}{2}$ cwt sulphate of
ammonia.

Cultivations, etc.: Ploughed: Feb 5. Seed drilled with superphosphate:
Mar 7. Sulphate of ammonia applied: Mar 9. Harvested: Combine
plots - Aug 22, Binder - Aug 25. Variety: Herta. Previous crop:
Potatoes.

Standard error per plot:

Grain: 2.41 cwt per acre or 6.1% (11 d.f.)

Summary of Results

	Methods of Harvesting			
	Binder	Combine Harvester	Mean	Difference
Yields: cwt per acre				
Grain	36.8	41.6	39.2	4.8 (± 0.98)
Straw	39.4	-		

Mean Dry Matter %: Combine plots - 84.2, Binder plots - no value
available.