

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1954

[Full Table of Content](#)



54/R/CA/4 Wheat - Control Wireworm

Rothamsted Research

Rothamsted Research (1955) *54/R/CA/4 Wheat - Control Wireworm* ; Yields Of The Field Experiments 1954, pp 68 - 68 - DOI: <https://doi.org/10.23637/ERADOC-1-184>

54/Ca/4.

WHEAT

The residual effects of insecticides on the control of wireworm,
3rd year - Geescroft 1954.

System of replication: 3 randomized blocks of 8 plots each.

Area of each plot: 0.0289 acre. Area harvested: 0.0205 acre.

Treatments, applied autumn 1951 for wheat:

None (three plots per block)									(O)	
Gammexane seed dressing 2 oz per bushel									(S)	
Gammexane combine drilled with seed at	56 lb per acre	3.5%	dust						(G)	
Aldrin	"	"	"	"	"	200 lb	"	"	1.78%	(A)
Chlordane	"	"	"	"	"	100 lb	"	"	5%	(C)
D.D.T.	"	"	"	"	"	150 lb	"	"	5%	(D)

Basal dressing, per acre: $\frac{1}{4}$ cwt compound granular fertilizer (12% N, 12% P₂O₅, 15% K₂O) combine drilled with seed. $\frac{2}{2}$ cwt sulphate of ammonia in spring.

Cultivations, etc.: Ploughed: Sept 23, 1953. Seed treated with mercurial seed dressing, combine drilled at $\frac{1}{2}$ bushels per acre: Nov 5. Sulphate of ammonia applied: Apr 30, 1954. Sprayed with M.C.P.A. high volume, 2 pints per acre: May 24. Combine harvested: Sept 7. Variety: Cappelle. Previous crop: Wheat.

Standard error per plot:

Grain (at 85% D.M.): 1.79 cwt per acre or 6.1% (16 d.f.)

Summary of Results

	O	S	G	A	C	D	Mean
Grain (at 85% dry matter): cwt per acre							
Mean (± 1.03)	27.2 ⁽¹⁾	29.4	32.9	31.0	29.7	29.2	29.2
Increase (± 1.19)		2.2	5.7	3.8	2.5	2.0	
(1) ± 0.60	Mean dry matter % as harvested: 74.9.						

Note: Wireworm counts were made and are available.