


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/CA/5 Spring Wheat - Residuals of Dung, N P K

Rothamsted Research

Rothamsted Research (1956) *55/R/CA/5 Spring Wheat - Residuals of Dung, N P K* ; Yields Of The Field Experiments 1955, pp 77 - 77 - DOI: <https://doi.org/10.23637/ERADOC-1-175>

55/Ca/5

SPRING WHEAT

Residual effects of Dung, Nitrogen, Phosphate and Potash - Sawyers I 1955.

Design: 4 randomized blocks of 8 plots each, the interaction DNEK being confounded with block differences.

Area of each plot: 0.0210 acre. Area harvested: 0.0150 acre.

Treatments, applied to potatoes in 1954: All combinations of:-

Dung: None; 10 tons per acre.

Nitrogen: None; 0.6 cwt N per acre applied as sulphate of ammonia.

Phosphate: None; 0.6 cwt P₂O₅ per acre applied as superphosphate.

Potash: None; 1.0 cwt K₂O per acre applied as muriate of potash.

Basal dressing to wheat: 4 cwt nitrochalk per acre; 21 cwt ground chalk per acre.

Cultivations, etc.: Ploughed: Jan 21, 1955. Chalk applied: Mar 31.

Nitrogen applied, seed drilled at 2 bushels per acre: Apr 1.

Sprayed with DNOC at 8 lb per acre in 80 gallons: May 2. Combine

harvested: Sept 1. Variety: Koga II. Previous crop: Potatoes.

Standard error per plot:

Grain: 2.20 cwt per acre or 7.5% (18 d.f.)

For details of the preceding potato experiment see 54/Cd/1.

Summary of Results

Grain: Mean yield 29.5 cwt per acre

Responses to treatments

Response to	Mean	Dung: tons per acre		N		P ₂ O ₅		K ₂ O	
		None	10	None	0.6	None	0.6	None	1.0
	(±0.78)	(±1.10)							
Dung	+1.2	-	-	+1.1	+1.3	+1.5	+0.9	+0.5	+1.9
N	+1.7	+1.6	+1.8	-	-	+0.8	+2.6	+1.4	+2.0
P ₂ O ₅	+1.4	+1.7	+1.1	+0.5	+2.3	-	-	+2.0	+0.8
K ₂ O	-0.1	-0.8	+0.6	-0.4	+0.2	+0.5	-0.7	-	-

Mean dry matter % as harvested: 85.0