


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/CA/7 Spring Wheat - Varieties and N

Rothamsted Research

Rothamsted Research (1956) *55/R/CA/7 Spring Wheat - Varieties and N* ; Yields Of The Field Experiments 1955, pp 80 - 80 - DOI: <https://doi.org/10.23637/ERADOC-1-175>

55/Ca/7

SPRING WHEAT

Varieties and levels of nitrogen - Great Field I 1955.

Design: 3 randomized blocks of 5 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0083 acre. Area harvested: 0.0057 acre.

Treatments: All combinations of:

Whole plots. Varieties: Atle; Atson; Koga II; Peko; Progress.

Sub plots. Nitrogen: 0.3; 0.6 cwt N per acre applied as nitrochalk.

Basal dressing: 1.15 cwt compound granular fertilizer (13% P_2O_5 , 13% K_2O) per acre, combine drilled with seed.

Cultivations, etc.: Ploughed: Dec 29, 1954. Nitrogen applied, seed combine drilled at $2\frac{1}{2}$ bushels per acre: Apr 14, 1955. Sprayed with DNOC at 6 lb per acre in 80 gallons: May 12. Combine harvested: Aug 31. Previous crop: Potatoes.

Standard errors per plot, Grain:

Whole plot: 1.75 cwt per acre or 4.7% (8 d.f.)

Sub plot: 1.35 cwt per acre or 3.6% (10 d.f.)

Summary of Results

Grain: cwt per acre

	Atle	Atson	Variety Koga II	Peko	Progress	Mean
	(±1.15)*					
N: cwt per acre						
0.3	35.9	35.0	40.6	38.3	38.0	37.6
0.6	33.4	32.8	41.0	38.0	37.4	36.5
Mean (±1.01)	34.7	33.9	40.8	38.1	37.7	37.0
Diff. (±1.10)	-2.5	-2.2	+0.4	-0.3	-0.6	-1.1 (±0.49)

*for use in comparisons other than vertical.

Mean dry matter % as harvested: 83.5