


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/CB/1 Barley - Seed Rates, and Levels of N

Rothamsted Research

Rothamsted Research (1956) *55/R/CB/1 Barley - Seed Rates, and Levels of N ; Yields Of The Field Experiments 1955*, pp 81 - 81 - DOI: <https://doi.org/10.23637/ERADOC-1-175>

55/Cb/1

BARLEY

Seed rates and levels of nitrogen - Great Field I 1955.

Design: 3 randomized blocks of 12 plots each.

Area of each plot: 0.0140 acre.

Treatments: All combinations of:-

Seed rates: 1; 2; 3 bushels per acre.

Nitrogen: None; 0.3; 0.6; 0.9 cwt N per acre applied as sulphate of ammonia.

Basal dressing: 1.15 cwt compound granular fertilizer (13% P_2O_5 , 13% K_2O) per acre combine drilled. 1 ton ground chalk per acre.

Cultivations, etc.: Ploughed: Dec 29, 1954. Chalk and sulphate of ammonia applied: April 1, 1955. Seed combine drilled: April 2. Sprayed with D.N.O.C. at 6 lb. per acre in 80 gallons: May 11. Combine harvested: Aug 20. Variety: Proctor. Previous crop: Potatoes.

Standard error per plot:

Grain (at 85% D.M.): 2.43 cwt per acre or 5.7% (22 d.f.).

Summary of Results

Seed rate: bushels per acre	N: cwt per acre				Mean
	None	0.3	0.6	0.9	
Grain (at 85% dry matter): cwt per acre					
		(+1.40)			(+0.70)
1	41.4	44.7	41.7	42.9	42.7
2	43.3	44.6	44.7	40.7	43.3
3	41.4	42.4	43.2	41.6	42.1
Mean (± 0.81)	42.1	43.9	43.2	41.7	42.7
Straw: cwt per acre					
1	27.7	33.0	30.9	32.8	31.1
2	27.7	31.9	33.0	35.3	32.0
3	25.5	29.6	33.2	36.6	31.2
Mean	27.0	31.5	32.4	34.9	31.4

Mean dry matter % as harvested, Grain: 82.6

Records of incidence of disease (Eyespot and Take-All), estimates of % area lodged and counts of ear emergence and plant, straw and ear numbers, were made.