


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/CC/1 Spring Beans - Control of Aphids (Spraying and Times of Sowing)

Rothamsted Research

Rothamsted Research (1956) *55/R/CC/1 Spring Beans - Control of Aphids (Spraying and Times of Sowing)* ; Yields Of The Field Experiments 1955, pp 88 - 88 - DOI:

<https://doi.org/10.23637/ERADOC-1-175>

55/Cc/1

SPRING BEANS

The control of Black Aphids (*Aphis fabae*) by spraying and time of sowing - Fosters 1955.

Design: 4 x 4 Latin square, plots split into 2 for the application of spray.

Area of each sub plot: 0.0189 acre.

Treatments: All combinations of:-

Whole plots. Times of sowing: Mar 19; April 6; April 25; May 13.

Sub plots. Spray: None; "Metasystox" (0.05% active ingredient) at 100 gallons per acre.

Note: The beans sown on May 13 were sprayed twice.

Basal dressing: 6 cwt compound granular fertilizer (10% P₂O₅, 20% K₂O) per acre.

Cultivations, etc.: Ploughed: Oct 22, 1954. Basal fertilizer applied, seed drilled at 195 lb per acre: Mar 19, 1955, April 6, April 25, May 13 respectively. Sprayed with "Metasystox": June 23. Last sown crop sprayed with "Metasystox" for second time: July 22. Combine harvested first three sowings: Aug 27. Combine harvested last sowing: Sept 12. Variety: Garton's Tick. Previous crop: Barley.

Standard errors per plot. Grain (at 85% dry matter):

Whole plot: 1.49 cwt per acre or 12.0% (6 d.f.)

Sub plot: 1.54 cwt per acre or 12.4% (12 d.f.)

Counts of Black Aphids and of aphid predators were made at weekly intervals from June to August.

Summary of Results

Grain (at 85% dry matter): cwt per acre

Spray	Time of sowing				Mean
	Mar 19	Apr 6	Apr 25	May 13	
	(±0.92)*				
None	16.9	13.1	6.7	2.4	9.8
Metasystox	19.2	18.3	12.0	10.5	15.0
Mean (±0.74)	18.0	15.7	9.3	6.4	12.4
Difference (±1.09)	2.3	5.2	5.3	8.1	5.2
					(±0.54)

* for use in comparisons other than vertical.

Mean dry matter % as harvested: 82.4.