


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/CE/1 Lucerne - Levels and Times of K

Rothamsted Research

Rothamsted Research (1956) *55/R/CE/1 Lucerne - Levels and Times of K ; Yields Of The Field Experiments 1955*, pp 99 - 99 - DOI: <https://doi.org/10.23637/ERADOC-1-175>

55/Ce/1

LUCERNE

Single and repeated applications of potash - Great Harpenden II 1955 - the first year.

Design: 6 randomized blocks of 8 plots each.

Area of each plot: 0.0147 acre.

Treatments 1955:

Potash seedbed dressings: None (2 plots per block); 0.33; 0.66; 1.0 (2 plots per block); 2.0; 3.0 cwt K_2O per acre applied as muriate of potash.

Note: It is intended to repeat the following dressings annually for two further years: None; 0.33; 0.66; 1.0 (1 plot per block) cwt K_2O per acre.

Basal dressing: 14 cwt per acre ground chalk, 2.3 cwt per acre superphosphate placed beneath seed.

Cultivations, etc.: Ploughed: Oct 12, 1954. Chalk applied: Apr 6, 1955. Potash applied: May 6. Seed drilled at 174 lb. per acre with placed superphosphate: May 9. Cut: Sept 14. Variety: Du Puits. Previous crop: Barley.

Standard error per plot:

Dry Matter: 1.01 cwt per acre or 17.7% (36 d.f.)

Note: The yields have been corrected for a linear fertility trend along the blocks.

Summary of Results

	Dry matter: cwt per acre (1 cut)						
	K ₂ O: cwt per acre						
	None	0.33	0.66	1.0	2.0	3.0	Mean
Mean (±0.41)	5.2 ⁽¹⁾	5.7	5.8	5.6 ⁽¹⁾	6.3	6.1	5.7
Increase (±0.50)		0.5	0.6	0.4 ⁽²⁾	1.1	0.9	

(1) ± 0.29

(2) ± 0.41

Mean dry matter % as harvested: 36.9