


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/CA/2 Winter Wheat - Control of Bulb Fly (Insecticides)

Rothamsted Research

Rothamsted Research (1957) *56/R/CA/2 Winter Wheat - Control of Bulb Fly (Insecticides)* ; Yields Of The Field Experiments 1956, pp 72 - 72 - DOI: <https://doi.org/10.23637/ERADOC-1-176>

56/Ga/2

WINTER WHEAT

Control of wheat bulb fly by insecticides - Pennell's Piece 1956.

Design: 5 randomized blocks of 6 plots each.

Area of each plot: 0.00643 acres. Area harvested: 0.00621 acres.

Treatments: Insecticides:-

- None (2 plots per block). (0)
- 4% Dieldrin dust at 1 cwt per acre combine drilled with seed. (1)
- 4% Dieldrin dust at 1 cwt per acre broadcast on surface. (2)
- Sprayed early with Parathion 0.1% v/v at 100 gallons per acre. (3)
- Sprayed late with Parathion 0.1% v/v at 100 gallons per acre. (4)

Basal dressing: 3 cwt 'Nitro-Chalk' per acre.

Note: all seed dressed with organo-mercurial fungicide.

Cultivations, etc.:

Ploughed: Sept 5, 1955. Seed drilled at 2 bushels per acre: Nov 1.
Dieldrin dust broadcast: Feb 16, 1956. Sprayed with Parathion:
'3' plots - Mar 8, '4' plots - Apr 12. 'Nitro-Chalk' applied:
Apr 30. Combine harvested: Sept 4. Variety: Cappelle. Previous
crop: Bare fallow.

Standard error per plot:

Grain (at 85% dry matter): 4.29 cwt per acre or 9.2% (21 d.f.)

Counts of numbers of plants, tillers, damaged tillers and of wheat bulb fly larvae were made.

Summary of Results

Grain (at 85% dry matter): cwt per acre

		Insecticides					Mean
		0	1	2	3	4	
Mean	(± 1.92)	43.9 ⁽¹⁾	47.3	52.5	46.1	47.2	46.8
Increase	(± 2.35)		3.4	8.6	2.2	3.3	
	(1) ± 1.36						

Mean dry matter % as harvested: 69.8.