


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/CA/6 Spring Wheat - Varieties and N

Rothamsted Research

Rothamsted Research (1957) *56/R/CA/6 Spring Wheat - Varieties and N* ; Yields Of The Field Experiments 1956, pp 79 - 79 - DOI: <https://doi.org/10.23637/ERADOC-1-176>

56/Ca/6

SPRING WHEAT

Varieties and levels of nitrogen - Little Hoos 1956.

Design: 3 randomized blocks of 8 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0101 acres. Area harvested: 0.0067 acres.

Treatments: All combinations of:

Whole plots. Varieties: Atle (1); Atson (2); Bersee (3),
Koga II (4); Miana (5); Peko (6); Progress (7); Svenno (8).
Sub plots. Nitrogen: 0.3; 0.6 cwt N per acre applied as
'Nitro-Chalk'.

Basal dressing: 1 cwt superphosphate per acre combine drilled with seed.

Cultivations, etc.: Ploughed: Oct 14, 1955 and again Jan 24, 1956.
'Nitro-Chalk' applied, seed combine drilled at $2\frac{3}{4}$ bushels per acre: Mar 19. Sprayed with DNOG at 6 lb in 90 gallons: May 4.
Combine harvested: Sept 20. Previous crop: Potatoes.

Standard errors per plot, Grain (at 85% dry matter):

Whole plot: 1.74 cwt per acre or 5.7% (14 d.f.)
Sub plot: 1.76 cwt per acre or 5.8% (16 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre	Variety								Mean
	1	2	3	4	5	6	7	8	
	(± 1.24)*								
0.3	29.5	32.2	24.6	33.5	25.3	34.3	30.6	28.7	29.8
0.6	29.9	33.8	26.9	33.2	26.6	33.5	31.2	32.0	30.9
Mean (± 1.01)	29.7	33.0	25.8	33.4	25.9	33.9	30.9	30.3	30.4
Difference (± 1.43)	+0.4	+1.6	+2.3	-0.3	+1.3	-0.8	+0.6	+3.3	+1.1
									(± 0.51)

* for use in comparisons other than vertical.

Mean dry matter % as harvested: 80.5