


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/A/1 Broadbalk - Wheat

Rothamsted Research

Rothamsted Research (1959) *58/R/A/1 Broadbalk - Wheat* ; Yields Of The Field Experiments 1958, pp 4 - 5 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

58/A/1.1

WHEAT - BROADBALK 1958

The 115th year

For history, treatments, etc., see "Details of the Classical and Long Term Experiments" 1956.

Cultivations, etc.:

Cropped sections. Dung applied: Sept 4, 1957. Ploughed: Sept 3-17. Ground chalk applied: Sept 9-16. Autumn fertilizers applied, seed drilled at $2\frac{3}{4}$ bushels per acre: Nov 20. Section 1A under continuous wheat sprayed with CMPP at 6 pints in 40 gallons per acre, spring fertilizers applied: Apr 30, 1958. Second dressing of nitrate of soda applied to plot 16: May 15. Combine harvested: Sept 9. Variety: Squareheads Master 13/4.

Fallow section. (V) Ploughed: Sept 3-17, 1957, May 7-9, 1958, July 29-31.

On a few plots an estimate was made of the chaff, cavings, dust, etc., not picked up by the baler.

Broadbalk Wilderness. N.

Cultivations, etc.: Shrubs grubbed out: Dec 2-6, 1957. Part mown: Apr 17, 1958, May 1, May 20, June 11, July 16, Aug 8, Oct 1.

Summary of Results

Grain (at 85% dry matter): cwt per acre

Section	II	IB	III	IV	IA	
Years after fallow	1	2	3	4	7	Mean
2A	23.4	21.7	21.1	22.7	24.2	22.4
2B	24.1	23.5	27.6	26.5	27.7	25.8
3	20.5	11.5	11.7	12.8	17.3	14.7
5	14.1	7.5	14.8	18.6	21.0	15.0
6	18.9	13.3	19.8	21.5	22.3	19.3
7	21.8	21.9	23.5	21.6	25.0	22.4
8	24.9	24.7	24.7	23.3	26.0	24.4
9	26.1	20.7	19.2	19.6	24.1	20.7
10	18.5	21.1	18.1	18.9	19.0	18.9
11	14.5	19.5	16.0	17.8	15.9	16.6
12	15.5	21.8	17.4	20.2	21.2	18.5
13	26.1	22.3	21.0	20.5	22.0	22.5
14	16.8	22.0	19.0	20.4	22.7	19.5
15	27.8	19.3	19.7	16.9	20.9	21.1
16	24.9	21.3	22.5	20.9	20.3	22.4
17	27.2	18.4	22.8	22.4	25.8	23.4
18	21.4	10.5	11.8	12.4	11.7	14.3
19	25.3	20.9	17.4	15.5	20.3	19.7
20	21.2	17.3	-	-	21.0	19.7

58/A/1.2

Straw (at 85% dry matter): cwt per acre

Section Years after fallow	II	IB	III	IV	IA	Mean
	1	2	3	4	7	
2A	42.6	30.8	30.3	29.9	23.1	33.0
2B	54.4	38.6	38.6	38.3	14.0	41.1
3	23.1	14.6	18.0	16.5	42.1	19.7
5	20.4	16.5	16.2	18.7	23.2	18.5
6	34.1	19.6	26.0	29.0	26.8	28.1
7	36.0	41.3	32.2	34.8	20.5	34.4
8	25.5	43.1	37.8	39.5	17.2	34.3
9	46.8	20.5	31.2	29.6	48.8	33.3
10	22.7	26.3	23.3	23.2	23.5	23.7
11	20.2	23.3	19.5	22.2	18.0	21.0
12	22.8	27.7	21.5	28.3	25.3	24.7
13	43.1	35.0	28.8	33.6	29.8	34.8
14	21.1	19.6	23.1	27.7	32.8	23.9
15	43.5	32.0	30.3	34.3	31.0	35.2
16	37.1	34.4	33.2	30.2	24.3	33.0
17	41.4	24.8	31.6	31.4	37.2	33.6
18	33.6	17.5	19.7	20.6	14.8	21.9
19	34.3	31.5	27.7	28.7	24.6	30.0
20	27.9	26.0	-	-	27.2	27.1

Mean dry matter % as harvested: Grain 77.5
Straw 84.6