


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/CB/1 Barley - N P and Residual Dung N P K

Rothamsted Research

Rothamsted Research (1959) *58/R/CB/1 Barley - N P and Residual Dung N P K* ; Yields Of The Field Experiments 1958, pp 88 - 89 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

58/Cb/1.1

BARLEY

Residual effects of dung, N, P and K applied to potatoes 1957 and direct effects of N and P - West Barnfield II 1958.

Design: Half replicate of 4×2^5 arranged in 4 blocks of 16 plots each, the identity being $d(20 - 10 + 5 - 0)npkNP = 1$, with certain high order interactions confounded with block differences.

Area of each plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments. All combinations of:-

Applied to potatoes 1957.

Dung: None; 5; 10; 20 tons per acre ploughed in.

N: None; 0.9 cwt N per acre as sulphate of ammonia.

P: None; 0.75 cwt P_2O_5 per acre as superphosphate.

K: None; 1.5 cwt K_2O per acre as muriate of potash.

Applied to barley 1958.

N: None; 0.4 cwt N per acre as 'Nitro-Chalk'.

P: None; 0.4 cwt P_2O_5 as superphosphate.

Basal dressing: 0.8 cwt K_2O per acre as muriate of potash.

Cultivations, etc.: Ploughed: Nov 7, 1957. 'Nitro-Chalk' and basal fertilizers applied, seed drilled at 2 bushels per acre: Mar 25, 1958. Sprayed with MCPB at 5 pints in 40 gallons per acre: May 28. Combine harvested: Sept 2. Variety: Proctor. Previous crop: Potatoes.

Standard error per plot.

Grain (at 85% dry matter): 1.51 cwt per acre or 4.4% (27 d.f.)

58/Cb/1.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

		Dung to potatoes 1957: tons per acre				Mean
		None	5	10	20	
Mean	(± 0.38)	33.6	33.9	34.1	35.0	34.2
<u>Applied to potatoes 1957</u>						
N: cwt per acre						
None	(± 0.53)	34.1	34.3	34.1	34.8	34.3
0.9		33.2	33.6	34.1	35.2	34.0
Difference	(± 0.76)	-0.9	-0.7	0.0	+0.4	-0.3 (± 0.38)
P ₂₅ : cwt per acre						
None	(± 0.53)	32.1	32.7	33.7	34.1	33.1
0.75		35.2	35.2	34.5	35.9	35.2
Difference	(± 0.76)	+3.1	+2.5	+0.8	+1.8	+2.1 (± 0.38)
K ₂ O: cwt per acre						
None	(± 0.53)	33.6	34.3	33.4	35.7	34.2
1.5		33.7	33.6	34.8	34.3	34.1
Difference	(± 0.76)	+0.1	-0.7	+1.4	-1.4	-0.1 (± 0.38)
<u>Applied to barley 1958</u>						
N: cwt per acre						
None	(± 0.53)	33.2	32.9	33.6	34.6	33.6
0.4		34.1	35.0	34.6	35.4	34.8
Difference	(± 0.76)	+0.9	+2.1	+1.0	+0.8	+1.2 (± 0.38)
P ₂₅ : cwt per acre						
None	(± 0.53)	31.0	32.7	33.1	34.0	32.7
0.4		36.2	35.2	35.0	35.9	35.6
Difference	(± 0.76)	+5.2	+2.5	+1.9	+1.9	+2.9 (± 0.38)

Responses to treatments

cwt per acre

Response to	Applied to potatoes 1957						Applied to barley 1958			
	N	P ₂₅	K ₂ O	N	P ₂₅	K ₂ O	N	P ₂₅	K ₂ O	
	None	0.9	None	0.75	None	1.5	None	0.4	None	0.4
<u>Applied to potatoes 1957</u>										
N	-	-	+0.3	-0.9	-1.0	+0.4	+0.8	-1.4	-0.6	0.0
P ₂₅	+2.6	+1.4	-	-	+1.5	+2.5	+2.4	+1.6	+2.9	+1.1
K ₂ O	-0.9	+0.5	-0.7	+0.3	-	-	-0.2	-0.2	0.0	-0.4
<u>Applied to barley 1958</u>										
N	+2.3	+0.1	+1.6	+0.8	+1.2	+1.2	-	-	+1.0	+1.4
P ₂₅	+2.6	+3.2	+3.8	+2.0	+3.1	+2.7	+2.7	+3.1	-	-
Mean dry matter % as harvested: 80.9										