


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1959

[Full Table of Content](#)



59/R/CG/3 Grass - Species and N

Rothamsted Research

Rothamsted Research (1960) *59/R/CG/3 Grass - Species and N* ; Yields Of The Field Experiments 1959, pp 105 - 106 - DOI: <https://doi.org/10.23637/ERADOC-1-179>

59/Cg/3.2

Summary of ResultsDry matter: cwt per acre

N: cwt per acre*	Species				Mean
	C	M	R	T	
<u>1st cut</u>					
	(±0.85)				(±0.43)
None	9.1	19.6	20.7	12.9	15.5
0.3	25.1	34.8	39.9	30.1	32.5
0.6	34.0	40.8	45.2	36.0	39.0
Mean (±0.49)	22.7	31.7	35.2	26.3	28.9
<u>2nd cut</u>					
	(±1.49)				(±0.74)
None	6.2	5.5	1.7	10.8	6.0
0.3	24.0	15.3	12.2	27.9	19.9
0.6	31.4	21.5	21.0	33.6	26.8
Mean (±0.85)	20.6	14.1	11.6	24.1	17.5
<u>3rd cut</u>					
	(±1.01)				(±0.50)
None	2.0	1.9	1.3	1.6	1.7
0.3	17.4	9.8	4.2	13.0	11.1
0.6	30.1	17.5	7.8	21.1	19.1
Mean (±0.59)	16.5	9.7	4.4	11.9	10.6
<u>Total of 3 cuts</u>					
	(±2.18)				(±1.09)
None	17.2	26.9	23.6	25.2	23.2
0.3	66.5	59.8	56.3	71.1	63.4
0.6	95.5	79.7	73.9	90.7	84.9
Mean (±1.25)	59.7	55.5	51.2	62.3	57.1

Mean dry matter % as cut:
 1st cut: 22.7
 2nd cut: 36.7
 3rd cut: 30.5
 Total of 3 cuts: 30.0

Species
 C S37 Cocksfoot
 M S215 Meadow Fescue
 R S24 Perennial Ryegrass
 T Timothy "Scotia"

* Applied for each cut.

59/Cg/4.1

CLOVER AND GRASS LEYS

The comparison of clover and grass leys as a preparation for wheat - West Barnfield II, 1959.

Design: 4 randomized blocks of 16 plots each.

Area of each plot: 0.0159 acres. Area harvested: 0.0068 acres.

Treatments:

Nitrogen to Leys 1959:-

To clover: none (4 plots per block)

To ryegrass: none, R1 and R2 (4 plots per block in each case)

Where R1 = 0.6 cwt N in spring, 0.15 cwt N after 1st hay cut.

R2 = 1.2 cwt N in spring, 0.30 cwt N after 1st hay cut.

The Nitrogen was applied as 'Nitro-chalk'

Note: the experiment is designed to include four rates of N applied to wheat in 1960/61.

Basal Dressings per acre:

To barley nurse crop 1958: 3 cwt compound fertilizer (10% P_2O_5 , 20% K_2O) combine drilled; 2 cwt sulphate of ammonia in seedbed.
To leys combine drilled in seedbed 1958: 1 cwt superphosphate.

Cultivations, etc., barley drilled March 25th, 1958: superphosphate applied, leys undersown in barley, ryegrass at 30 lb. and clover at 20 lb. per acre: April 22nd.

'Nitro-chalk' dressings applied: March 12th and May 25th, 1959.

Cut twice for hay: May 20th and July 20th. Varieties: Italian ryegrass S22 and Double cut red clover S151.

Standard errors per plot:

Ryegrass. Dry matter:

1st cut:	3.41 cwt per acre or 7.5%	(42 d.f.)
2nd cut:	0.97 cwt per acre or 7.5%	(42 d.f.)
Total of 2 cuts:	3.78 cwt per acre or 6.5%	(42 d.f.)