


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1959

[Full Table of Content](#)



59/R/CA/2 Winter Wheat - Seed Rates, Sowing Dates and N (After Non-cereal Crop)

Rothamsted Research

Rothamsted Research (1960) *59/R/CA/2 Winter Wheat - Seed Rates, Sowing Dates and N (After Non-cereal Crop)* ; Yields Of The Field Experiments 1959, pp 69 - 70 - DOI:

<https://doi.org/10.23637/ERADOC-1-179>

59/Ca/2.2

Summary of ResultsGrain (at 85% dry matter): cwt per acre

Seed rate: bushels. per acre	Date of sowing				N: cwt per acre (including basal)		Diff.	Mean
	Oct 21st	Nov 11th	Nov 25th	Jan 8th	0.6	1.1		
	(±1.50)				(±0.81)*		(±0.61)	(±0.75)
2	40.0	41.6	42.4	35.3	36.1	43.6	+7.5	39.8
4	43.1	42.4	43.4	37.3	38.0	45.1	+7.1	41.6
	Date of sowing				(±1.14)*		(±0.86)	(±1.06)
	Oct 21st				37.7	45.3	+7.6	41.6
	Nov 11th				38.5	45.5	+7.0	42.0
	Nov 25th				39.8	46.0	+6.2	42.9
	Jan 8th				32.1	40.4	+8.3	36.3
	Mean				37.1	44.3	+7.2 (±0.43)	40.7

*For use in vertical and diagonal comparisons.

Mean dry matter % as harvested: 78.4

59/Ca/3.1

SPRING WHEAT

Row spacing, seed rates and nitrogen - Long Hoos I, II and III 1959.

Design: 2 randomized blocks of 10 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot (acres): 0.0148. Area harvested: 0.01 acre approximately (varying with row spacing).

Treatments. All combinations of:-

Whole plots. Row spacing, inches/seed rate, bushels per acre:
7/2; 7/4; 14/4; 14/2; 7B/3, where B = every 4th row blank.

Types of drill: Standard; precision.

Sub plots. Nitrogen: 0.6; 1.2 cwt N per acre as 'Nitra-Shell' in seed bed.

Basal dressing: None.

Cultivations, etc.: Ploughed: Oct 21, 1958. 'Nitra-Shell' applied, seed drilled: Mar 2, 1959. Sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 14 and 27. Combine harvested: Aug 22. Variety: Koga II. Previous crop: Oats.

Standard error per plot, Grain (at 85% dry matter):

Whole plot: 1.35 cwt per acre or 4.3% (9 d.f.)

Sub plot: 2.68 cwt per acre or 8.5% (10 d.f.)