


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1959

[Full Table of Content](#)



59/R/CE/3 and 59/W/CE/3 Beans - Time of Sowing, Spraying P and K

Rothamsted Research

Rothamsted Research (1960) *59/R/CE/3 and 59/W/CE/3 Beans - Time of Sowing, Spraying P and K ; Yields Of The Field Experiments 1959*, pp 89 - 91 - DOI: <https://doi.org/10.23637/ERADOC-1-179>

59/Ce/3.2

Summary of ResultsGrain (at 85% dry matter): cwt per acre

Sown			P ₂ O ₅ : cwt per acre			K ₂ O: cwt per acre			Mean	
Autumn	Spring	Diff.	None	0.5	1.0	None	1.0	2.0		
<u>Great Knott III, Rothamsted</u>										
<u>Spray</u>				(±0.66)*			(±0.66)*			
None	26.6	13.9	-12.7	19.8	20.0	20.8	20.1	19.8	20.8	20.2
Demeton-methyl	28.8	18.8	-10.0	23.7	23.4	24.3	24.2	24.5	22.7	23.8

Diff.	+2.2	+4.9	+2.7	+3.9	+3.4	+3.5	+4.1	+4.7	+1.9	+3.6
			(±2.36)	(±0.94)**			(±0.94)**			
<u>Sown</u>				(1) and (2)			(1) and (2)			
Autumn				26.8	27.9	28.4	27.4	27.9	27.9	27.7
Spring				16.7	15.5	16.8	16.9	16.5	15.6	16.3
Mean (±0.47)				21.8	21.7	22.6	22.1	22.2	21.7	22.0
Diff.(±1.41)				-10.1	-12.4	-11.6	-10.5	-11.4	-12.3	-11.4
										(±1.18)

(1) ±0.99 for use in diagonal comparisons only.

(2) ±0.66 for use in horizontal and interaction comparisons only.

*For use in horizontal comparisons only.

**For use only in testing the difference of two differences.

Mean dry matter % as harvested: Autumn sown, 85.2; spring sown, 85.1.

59/Ca/3.3

Grain (at 85% dry matter): cwt per acre

Sown			P ₂ O ₅ : cwt per acre			K ₂ O: cwt per acre			Mean	
Autumn	Spring	Diff.	None	0.5	1.0	None	1.0	2.0		
<u>Mill Dam Close, Woburn</u>										
<u>Spray</u>				(±2.06)*			(±2.06)*			
None	27.4	23.3	-4.1	24.4	27.9	23.8	25.8	24.3	25.9	25.4
Demeton-methyl	30.0	17.8	-12.2	25.6	26.2	19.9	19.5	24.9	27.2	23.9
-----										----
Diff.	+2.6	-5.5	-8.1	+1.2	-1.7	-3.9	-6.3	+0.6	+1.3	-1.5
			(±3.26)	(±2.91)**			(±2.91)**			
<u>Sown</u>				(1) and (2)			(1) and (2)			
Autumn				27.4	34.7	24.1	27.3	28.1	30.8	28.7
Spring				22.6	19.4	19.7	18.1	21.2	22.3	20.5
Mean (±1.46)				25.0	27.0	21.9	22.7	24.6	26.6	24.6
Diff.(±2.88)				-4.8	-15.3	-4.4	-9.2	-6.9	-8.5	-8.2
										(±1.63)

(1) ±2.04 for use in diagonal comparisons only.

(2) ±2.06 for use in horizontal and interaction comparisons only.

*For use in horizontal comparisons only.

**For use only in testing the difference of two differences.

Mean dry matter % as harvested: Autumn sown, 75.5; spring sown, 82.9.

59/Cf/1.1

POTATOES

Forms and levels of K - Rothamsted (R) Sawyers I and Woburn (W)
Lansome Field 1959.

Design (each field): 4 blocks of 8 plots with certain high order
interactions partially confounded with block differences.

Area of each plot: 0.0141 acres. Area harvested: 0.0071 acres.

Treatments: No potash (2 plots per block) and all combinations of:-

Forms of K: Potassium bi-carbonate, KHCO_3 (C);
Potassium sulphate, K_2SO_4 (S);
Potassium chloride, KCl (M);

Levels of K: 1.25; 2.50 cwt K_2O per acre.

All the above in combination with:-

Levels of N: 0.75; 1.50 cwt N per acre as sulphate of ammonia.

Basal dressing (each field): 1.0 cwt P_2O_5 per acre as superphosphate.

Cultivations, etc.:

Sawyers I (R): Ploughed: Feb 19, 1959. Ridged, fertilizers
applied by hand: Apr 20. Potatoes hand planted: Apr 21.
Earthed up: July 6. Sprayed with copper fungicide, 5 lb in
40 gallons per acre: Aug 24. Sprayed with sulphuric acid at
15% BOV in 100 gallons per acre: Sept 29. Lifted: Oct 12.
Variety: Majestic. Previous crop: Barley.

Lansome Field (W): Sprayed with dalapon at 10 lb and 5 lb in
40 gallons per acre: Nov 8, 1958 and Nov 22 respectively.
Ploughed: Jan 23 - Feb 2, 1959. Ridged: Apr 4. Fertilizers
applied and potatoes hand planted: Apr 10. Earthed up: June 22.
Sprayed with zineb at 2 lb and demeton methyl (against aphids)
at 12 fluid oz (50% active ingredients) in 40 gallons per acre:
Aug 15. Tops destroyed mechanically: Sept 22. Lifted:
Sept 29. Variety: Majestic. Previous crop: Barley.

Standard errors per plot. Total tubers:

Sawyers I (R): 1.215 tons per acre or 12.5% (14 d.f.)

Lansome (W): 1.076 tons per acre or 16.4% (13 d.f.)

Note (1): Chemical analyses were made of % N, P, K, Cl and S on
haulm and tubers at Rothamsted and on tubers at Woburn.

Note (2): At Woburn the plant was poor and irregular and the
yields were adjusted for plant number.