

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/DA/2 Winter Wheat - Varieties and N

Rothamsted Research

Rothamsted Research (1962) *61/R/DA/2 Winter Wheat - Varieties and N* ; Yields Of The Field Experiments 1961, pp 108 - 108 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/Da/2

WINTER WHEAT

Varieties and levels of nitrogen - Deacons Field 1961.

Design: 4 randomised blocks of 12 plots each.

Area of each plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments. All combinations of:-

Varieties: Cappelle (C); Professeur Marchal (M); Viking (V).

Levels of nitrogen (in addition to basal): 0.0; 0.3; 0.6; 0.9 cwt

N per acre as 'Nitro-Chalk'.

Basal dressing: 2 cwt compound fertiliser (6% N, 15% P₂O₅, 15% K₂O) per acre combine drilled.

Cultivations, etc.: Ploughed: Nov 15 - 25, 1960. Seed combine drilled at 3 bushels per acre: Feb 13, 1961. 'Nitro-Chalk' applied: Apr 15. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 8. Combine harvested: Sept 8. Previous crop: Spring beans.

Standard error. per plot.

Grain (at 85% dry matter): 2.03 cwt per acre or 4.9% (33 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre	C	Variety M	V	Mean
		(±1.01)		(±0.59)
None	31.1	31.7	28.2	30.3
0.3	37.3	37.5	42.2	39.0
0.6	43.1	46.5	49.1	46.2
0.9	50.0	50.4	50.9	50.4
Mean (±0.50)	40.3	41.5	42.6	41.4

Mean dry matter % as harvested: 79.6