


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/DD/2 Spring Beans - Control of Weeds

Rothamsted Research

Rothamsted Research (1962) *61/R/DD/2 Spring Beans - Control of Weeds* ; Yields Of The Field Experiments 1961, pp 122 - 123 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/Da/2.1

SPRING BEANS

Control of weeds by residual herbicides - Great Field I 1961.

Design: 3 randomised blocks of 23 plots each.

Area of each plot: 0.0176 acres. Area harvested: 0.0110 acres.

Treatments:

Beans drilled at normal (21") spacing, no inter row cultivations (O)

Beans drilled at narrow (10½") spacing, no inter row cultivations (A)

Beans drilled at normal (21") spacing, inter cultivated (B)

Treatments O, A, B taken factorially with:

No spray:

Simazine: ½; 1 lb active material per acre (S₁S₂) as pre-emergence spray

2,6-DBN: 1; 3 lb active material per acre (D₁D₂) as pre-emergence spray

Treatments O and A taken factorially with

2,6-DBN: 1½; 3 lb active material per acre as post-emergence spray.

Basal dressing: 3½ cwt per acre compound fertiliser (14% P₂O₅, 28% K₂O) placement drilled.

Cultivations, etc.: Ground chalk applied at 54 cwt per acre: Oct 4, 1960. Ploughed: Oct 14. Seedbed spray applied and worked in: Mar 4, 1961. All plots rotary cultivated except those already sprayed, seed placement drilled at 200 lb per acre: Mar 17. Pre-emergence spray applied: Mar 23. Post-emergence spray applied: May 1. Sprayed with demeton-methyl at 12 fluid oz in 60 gallons per acre: June 10. Combine harvested: Sept 4. Variety: Garton's Spring Tick. Previous crop: Winter wheat.

Note (1). The plots on which 2,6-DBN was worked into the seedbed failed reducing the experiments to 3 blocks of 19 plots each.

Note (2). Weed counts were taken on all wide row plots.

Standard error per plot.

Grain (at 85% dry matter): 2.34 cwt per acre or 9.7% (36 d.f.)

61/Da/2.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

	No inter row cultivation		Inter cultivated
	21" row	10 $\frac{1}{2}$ " row (± 1.35)	21" row
No spray	22.5	21.0	26.8
Pre-emergence S ₁	23.1	21.8	27.0
S ₂	27.2	24.5	26.5
D ₁	25.9	23.0	25.6
D ₂	27.8	23.8	28.3
Post-emergence D ₁	21.5	20.6	
D ₂	20.0	21.2	
General mean		24.1	

Mean dry matter % as harvested: 79.8