


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/DI/1 Carrots - Motley Dwarf Virus

Rothamsted Research

Rothamsted Research (1962) *61/R/DI/1 Carrots - Motley Dwarf Virus* ; Yields Of The Field Experiments 1961, pp 140 - 142 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/Di/1.1

CARROTS

The effect of a systemic insecticide on yield through control of motley dwarf virus - Woburn Lansome Field 1961.

Design: A plaid rectangle of 4 rows and 8 columns.

Area of each sub plot: 0.0048 acres. Area harvested: 0.0030 acres.

Treatments. All combinations of:-

Sowing dates (to columns): Apr 13 (D1); May 16 (D2).

Times of spraying with demeton methyl at 12 fluid oz in 40 gallons per acre:

To treatment D1: None (0); May 15 (S1); May 29 (S2);
June 19 (S3).

To treatment D2: None (0); June 5 (S1); June 19 (S2);
July 7 (S3).

Infection dates: To treatment D1: May 23 (I1); June 14 (I2).

To treatment D2: June 15 (I1); July 3 (I2).

Basal dressing: 10 cwt per acre compound fertiliser (10% N, 10% P₂O₅, 18% K₂O).

Cultivations, etc.: Sprayed with dalapon at 8 lb in 20 gallons per acre: Oct 13, 1960; and at 4 lb in 20 gallons per acre: Nov 9. Ploughed: Dec 5. Basal dressing applied: Apr 10, 1961. Seed drilled at 5 lb per acre: Apr 13 and May 16. Thinned: 1st sowing - June 19; 2nd sowing - July 24. Lifted: Sept. 26. Variety: Scarlet Intermediate. Previous crop: Spring wheat.

Note: Aphid counts and estimates of virus infection were made.

Standard error per plot.

Saleable roots: 1.210 tons per acre or 20.4% (10 d.f.)

Tops from saleable roots: 0.467 tons per acre or 18.4% (10 d.f.)

61/Di/1.2

Summary of Results

	<u>Time of spraying</u>				
	S ₀	S ₁	S ₂	S ₃	Mean
<u>Saleable roots: tons per acre</u>					
Mean (± 0.428)	6.85	6.42	5.71	4.79	5.94
<u>Sowing date</u>					
Apr 13	7.14	6.85	6.83	6.14	6.74
May 16	6.56	5.98	4.60	3.45	5.15
Diff. (± 0.856)**	-0.58	-0.87	-2.23	-2.69	-1.59
<u>Infection date</u>					
I ₁	6.75	6.45	6.02	4.40	5.90
I ₂	6.95	6.38	5.41	5.19	5.98
Diff. (± 0.856)	+0.20	-0.07	-0.61	+0.79	+0.08 (± 0.428)

Infection date	<u>Sowing date</u>	
	Apr 13	May 16
	<u>(± 0.428)*</u>	
I ₁	6.56	5.25
I ₂	6.93	5.04

<u>Tops from saleable roots: tons per acre</u>					
Mean (± 0.165)	2.76	2.83	2.45	2.09	2.53
<u>Sowing date</u>					
Apr 13	1.95	1.96	1.98	1.80	1.92
May 16	3.57	3.70	2.93	2.39	3.14
Diff. (± 0.330)**	+1.62	+1.74	+0.95	+0.59	+1.22
<u>Infection date</u>					
I ₁	2.88	2.84	2.62	1.92	2.57
I ₂	2.64	2.81	2.29	2.26	2.50
Diff. (± 0.330)	-0.24	-0.03	-0.33	+0.34	-0.07 (± 0.165)

Infection date	<u>Sowing date</u>	
	Apr 13	May 16
	<u>(± 0.165)*</u>	
I ₁	1.91	3.23
I ₂	1.94	3.06

* For use in vertical and interaction comparisons only

** For use only in testing the difference of 2 differences

61/Di/1.3

	<u>Time of spraying</u>				
	S ₀	S ₁	S ₂	S ₃	Mean
<u>Saleable numbers: thousands per acre</u>					
Mean	109.9	113.6	111.6	104.5	109.9
<u>Sowing date</u>					
Apr 13	107.7	115.8	115.0	107.3	111.5
May 16	112.0	111.5	108.3	101.7	108.4
Diff.	+4.3	-4.3	-6.7	-5.6	-3.1
<u>Infection date</u>					
I ₁	107.7	115.8	112.9	96.6	108.2
I ₂	112.1	111.5	110.4	112.4	111.6
Diff.	+4.4	-4.3	-2.5	+15.8	+3.4

Infection date	<u>Sowing date</u>	
	Apr 13	May 16
I ₁	109.9	106.6
I ₂	113.0	110.1