


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/W/B/6 Irrigation

Rothamsted Research

Rothamsted Research (1962) *61/W/B/6 Irrigation* ; Yields Of The Field Experiments 1961, pp 70 - 73
- DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/B/6.1

IRRIGATION EXPERIMENT

The 11th year - revised 1960

The effects of irrigation and nitrogen - Woburn Butt Close 1961.

For details of previous cropping, treatments etc. see "Details of the Classical and Long Term Experiments" 1956.

Cut grass: The basal PK compound is now applied in winter. Applications of muriate of potash are now made in spring and after each cut except the last at the rates 0.3; 0.6 cwt K_2O per acre. Plots of the different treatments may be cut and manured independently if rates of growth vary.

(applied to 1/2 plot)

Beans: Owing to weather conditions the third year crop in the rotation was spring, and not winter beans.

Area of each whole plot (acres): Spring beans: 0.0555.

Sub plots (acres): Grass: 0.0264; remainder: 0.0277.

Area harvested (acres): Early potatoes: 0.0171; barley: 0.0098; spring beans: 0.0181; Cut grass: 0.0165.

Rainfall and Irrigation: inches

Week ending	Rain-fall	Grass C	Barley G	Potatoes C	A	Beans B	C
May 1	0.36						
8	0.62						
15	-						
22	0.01	0.50	0.50	0.50	-	0.50	0.50
29	0.01	-	-	-	-	-	-
June 5	0.23	0.50	0.50	0.50	-	0.50	0.50
12	0.03	0.75	0.50	0.75	-	0.50	0.50
19	0.85	-	0.23	-	-	-	-
26	0.02	0.75	0.50	0.75	0.50	-	0.50
July 3	0.33	0.75	0.50	0.75	0.75	-	0.75
10	-	0.75	0.50	0.75	0.75	-	0.75
17	1.03	0.75	-	-	-	-	-
24	0.02	-	-	-	-	-	-
31	0.10	0.75	-	-	0.50	-	0.50
Aug 7	0.71	0.50	-	-	0.50	-	0.50
14	0.86						
21	0.21						
28	0.55						
Sept 4	0.32						
11	0.30						
Total	6.56	6.00	3.23	4.00	3.00	1.50	4.50

61/B/6.2

Cultivations, etc.:

Early potatoes: Ploughed twice: Sept 24, 1960 and Feb 14, 1961.
Machine planted, fertilisers applied: Mar 17. Appropriate plots
sprayed with simazine: Mar 31. Earthed up (except the simazine
plots): May 23. Lifted: July 18. Simazine plots ploughed:
July 19. Trefoil (inoculated seed) sown at 30 lb per acre:
July 21. Variety: Arran Pilot.
Barley: Ploughed: Feb 7, 1961. Fertilisers applied: Mar 7. Seed
drilled at $2\frac{1}{4}$ bushels per acre: Mar 8. Sprayed with CMFP at
5 pints in 40 gallons per acre: May 10. Combine harvested:
Aug 17. Variety: Proctor.
Spring beans: Ploughed 3 times: Aug 16 and Sept 26, 1960;
Jan 19, 1961. Seed placement drilled at 200 lb per acre with
PK compound: Feb 22. Sprayed with demeton methyl at 12 fluid
oz in 40 gallons per acre: May 29. Combine harvested: O and A -
Aug 12, B and C - Aug 21. Variety: Tick
Grass: PK compound applied: Nov 9, 1960. Muriate of potash and
'Nitro-Chalk' applied: Mar 21, 1961. Cut six times: Apr 18,
May 15, June 15, July 14, Aug 14, Sept 11. Muriate of potash
and 'Nitro-Chalk' applied to appropriate plots after each cut
except the last. One application, however (immediately following
the fourth cut), was to appropriate irrigated plots only.
Variety: S22 Italian ryegrass.

Standard errors per plot.

Early potatoes, total tubers
Whole plot: 1.338 tons per acre or 15.7% (4 d.f.)
 $\frac{1}{2}$ plot: 0.554 tons per acre or 6.5% (4 d.f.)
 $\frac{1}{4}$ plot: 1.012 tons per acre or 11.9% (8 d.f.)
Barley, (grain at 85% dry matter)
Whole plot: 2.06 cwt per acre or 6.9% (5 d.f.)
Sub plot: 2.16 cwt per acre or 7.3% (10 d.f.)
Spring beans, (grain at 85% dry matter)
Whole plot: 1.98 cwt per acre or 9.6% (6 d.f.)
Cut grass, dry matter. Total of 6 cuts
Whole plot: 3.38 cwt per acre or 5.4% (6 d.f.)
Sub plot: 4.60 cwt per acre or 7.4% (8 d.f.)

61/B/6.3

Summary of ResultsEarly potatoes, Total tubers: tons per acre

Weed control	Irrigation		Weed control Normal Simazine cultivation spray		Mean
	0	C			
	(± 0.320) ⁽¹⁾	(± 0.805) ⁽²⁾			
Normal cultivation	6.01	14.13			
Simazine spray	5.23	8.77			
N: cwt per acre including basal					
	(± 0.413) ⁽³⁾	(± 0.369) ⁽⁴⁾	(± 0.413) ⁽³⁾	(± 0.805) ⁽⁴⁾	
0.6	5.31	10.26	9.58	5.98	7.78
1.2	5.93	12.64	10.55	8.02	9.29
Mean	5.62	11.45	10.07	7.00	8.54
Difference	0.62	2.38	0.97	2.04	1.51
	(± 0.584)		(± 0.584)		(± 0.413)

Barley, (Grain at 85% dry matter): cwt per acre

N: cwt per acre including basal	Irrigation		Mean
	0	C	
	(± 0.88) ⁽³⁾	(± 1.05) ⁽⁴⁾	
0.2	23.3	29.0	26.2
0.4	31.3	35.4	33.4
Mean	(± 0.84)	27.3 32.2	29.7
Difference	(± 1.25)	+8.0 +6.4	+7.2
			(± 1.76)

Mean dry matter % as harvested: 82.3

Spring beans, (Grain at 85% dry matter): cwt per acre

Irrigation				
O	A	B	C	Mean
13.3	21.3	18.7	28.8	20.5
	(±1.14)			

Mean dry matter % as harvested: 77.9

- (1) For use in horizontal and interaction comparisons
- (2) For use in vertical and diagonal comparisons
- (3) For use in vertical and interaction comparisons
- (4) For use in horizontal and diagonal comparisons.

61/B/6.4

Cut grass, Dry matter: cwt per acre

Total of 6 cuts

For each cut		Irrigation		Mean
K ₂ O cwt per acre	N cwt per acre	0	C	
		(±2.66) ⁽¹⁾ (±2.71) ⁽²⁾		
None	0.3	46.0	68.9	57.4
0.3	0.3	44.7	69.4	57.1
None	0.6	46.2	81.5	63.8
0.6	0.6	45.4	97.9	71.6
Mean (±1.38)		45.6	79.4	62.5

Mean dry matter % as cut: 22.9

- (1) For use in comparisons within the same irrigation or absence or presence of K.
- (2) For use in comparisons involving different irrigations or absence or presence of K.