


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/B/8 Residual Phosphate Rotation

Rothamsted Research

Rothamsted Research (1962) *61/R/B/8 Residual Phosphate Rotation ; Yields Of The Field Experiments 1961*, pp 76 - 78 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/B/8.1

RESIDUAL PHOSPHATE ROTATIONS

The long term and residual effects of a number of phosphate fertilisers compared with superphosphate - Great Field IV and Sawyers I 1961, the second year.

Design: Great Field IV: 1 randomised block of 12 plots per crop.
Sawyers I: 2 randomised blocks of 12 plots per crop.

Rotation: Potatoes, Barley, Swedes.

Area of each plot (acres):

Great Field IV: 0.0193. Area harvested: Potatoes and Barley - 0.0129, Swedes - 0.0096.

Sawyers I: 0.0212. Area harvested: Potatoes and Barley - 0.0141, Swedes - 0.0106.

Treatments:

Granular superphosphate treatments broadcast in spring before sowing or ridging:-

1. No phosphate.
2. 0.25 cwt P_{205} per acre per year.
3. 0.50 cwt P_{205} per acre per year.
- 4 & 5. No phosphatic fertiliser in 1960 or 1961, but later at rates to be decided.

Phosphate fertilisers ploughed in (to a depth not exceeding 6 inches) at 3.0 cwt P_{205} per acre in September 1959 and rotary hoed in in spring.

6. Nitrophosphate I (17.1% P_{205} , none water soluble)
7. Nitrophosphate II (18.8% P_{205} , one quarter water soluble)
8. Nitrophosphate III (22.4% P_{205} , half water soluble)
9. Gafsa rock phosphate (28.9% P_{205})
10. Bessemer basic slag (15.2% P_{205})
11. Potassium metaphosphate* (57.9% P_{205} , 38.8% K_2O)
12. Granular superphosphate (20.4% P_{205})

*Note. To balance the K_2O content of potassium metaphosphate, all the other treatments included 2.0 cwt K_2O per acre as sulphate of potash in autumn 1959.

Basal dressings per acre: Broadcast in spring before sowing or ridging:

N as 'Nitro-Chalk' 21:-

To potatoes: 1.2 cwt; to barley: 0.6 cwt; to swedes: 0.5 cwt.

K_2O as sulphate of potash:-

To potatoes: 1.0 cwt; to barley: 1.0 cwt; to swedes: 1.0 cwt.

Cultivations, etc. (both fields, except as indicated):

Ploughed: Dec 28, 1960 - Jan 7, 1961. Ground chalk applied to Sawyers I at 9 cwt per acre: Feb 15.

61/B/8.2

Potatoes: Rotary cultivated twice, fertilisers applied, planted: Apr 20, 1961. Earthed up: July 5. Lifted: Sept 26.

Variety: Majestic. Previous crop: Swedes.

Barley: Fertilisers applied, seed drilled at 2 bushels per acre on Great Field IV: Mar 14, 1961. Sawyers I rotary cultivated twice, fertilisers applied, seed drilled at 2 bushels per acre: Mar 16. Sprayed with CMPP at 6 pints in 40 gallons per acre: May 12. Combine harvested: Aug 17. Variety: Proctor. Previous crop: Potatoes.

Swedes: Rotary cultivated twice: May 15, 1961. Fertilisers applied: May 16. Seed hand drilled at $2\frac{3}{4}$ lb per acre: May 17.

Singled: Great Field IV - June 30; Sawyers I - July 10.

Lifted: Nov 2. Variety: Wilhelmsburger. Previous crop: Barley.

Standard errors per plot:

Sawyers I

Potatoes, Total tubers: 0.705 tons per acre or 9.9% (13 d.f.)

Barley, Grain (at 85% dry matter): 3.41 cwt per acre or 10.5% (13 d.f.)

Swedes, Roots: 1.438 tons per acre or 11.4% (13 d.f.)

Note. For details for the previous year's results see 'Results of the Field Experiments' 1960 pages 60/B/9.1 to 60/B/9.3.

Summary of Results

Potatoes

Phosphate	Total tubers: tons per acre				Percentage ware ($1\frac{1}{2}$ " riddle)			
	Great Field IV		Sawyers I		Great field IV		Sawyers I	
	Mean	Increase	Mean	Increase	Mean	Increase	Mean	Increase
			$(\pm 0.499)(\pm 0.576)$					
None(1,4,5)	10.23		6.60 ⁽¹⁾		96.0		93.1	
2	9.24	-0.99	6.53	-0.07	95.2	-0.8	94.1	+1.0
3	11.53	+1.30	8.09	+1.49	94.7	-1.3	95.0	+1.9
6	11.78	+1.55	7.34	+0.74	96.8	+0.8	95.3	+2.2
7	11.53	+1.30	7.93	+1.33	96.8	+0.8	94.4	+1.3
8	11.89	+1.66	6.35	-0.25	95.7	-0.3	93.9	+0.8
9	11.49	+1.26	6.26	-0.34	96.3	+0.3	94.5	+1.4
10	10.83	+0.60	7.62	+1.02	96.4	+0.4	95.6	+2.5
11	10.98	+0.75	7.55	+0.95	95.2	-0.8	94.3	+1.2
12	11.52	+1.29	8.17	+1.57	94.8	-1.2	95.3	+2.2
Mean	10.96		7.13		95.8		94.3	
(1) (± 0.288)								

61/B/8.3

Phosphate	Great Field IV		Sawyers I		Great Field IV		Sawyers I	
	Mean	Increase	Mean	Increase	Mean	Increase	Mean	Increase
<u>Barley</u>								
<u>Grain (at 85% dry matter)</u>				<u>Straw (at 85% dry matter)</u>				
<u>cwt per acre</u>				<u>cwt per acre</u>				
			$(\pm 2.41)(\pm 2.78)$					
None(1,4,5)	41.0		31.0	(1)	31.1		17.1	
2	44.6	+3.6	32.4	+1.4	31.9	+0.8	15.2	-1.9
3	45.1	+4.1	30.0	-1.0	35.2	+4.1	15.1	-2.0
6	45.1	+4.1	29.2	-1.8	35.3	+4.2	20.7	+3.6
7	46.0	+5.0	30.0	-1.0	34.8	+3.7	19.4	+2.3
8	43.0	+2.0	37.0	+6.0	33.6	+2.5	20.1	+3.0
9	47.1	+6.1	35.5	+4.5	37.4	+6.3	20.5	+3.4
10	45.7	+4.7	34.1	+3.1	41.8	+10.7	20.1	+3.0
11	45.0	+4.0	33.2	+2.2	38.2	+7.1	19.9	+2.8
12	41.5	+0.5	34.4	+3.4	37.9	+6.8	19.9	+2.8
Mean	43.8		32.4		35.0		18.5	
Mean dry matter								
% as harvested:	81.9		80.5		72.0		71.5	
(1) (± 1.39)								

Swedes, Roots: tons per acre

			$(\pm 1.017)(\pm 1.174)$	
			6.20(1)	
None(1,4,5)	10.28		11.87	+5.67
2	12.83	+2.55	12.00	+5.80
3	17.04	+6.76	17.58	+11.38
6	18.57	+8.29	16.31	+10.11
7	22.09	+11.81	16.82	+10.62
8	22.18	+11.90	14.12	+7.92
9	19.59	+9.31	13.64	+7.44
10	18.89	+8.61	14.84	+8.64
11	19.03	+8.75	15.28	+9.08
12	22.73	+12.45		
Mean	16.98		12.59	
(1) (± 0.587)				