

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/C/1 Cereal-bean Rotations - Wheat and Barley (Test Crops)

Rothamsted Research

Rothamsted Research (1962) *61/R/C/1 Cereal-bean Rotations - Wheat and Barley (Test Crops)* ;
Yields Of The Field Experiments 1961, pp 88 - 89 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/C/1.1

CEREALS AND BEANS ROTATIONS

The effect of crop sequences on the incidence of cereal foot and root rot diseases - Great Field I 1961 - the 5th year.

Design: Three series each of 3 randomised blocks of 6 plots, starting in each of the years 1957, 1958 and 1959.

Area of each plot (acres): 0.0305. Area harvested: Winter wheat, series starting 1958 - 0.0095; series starting 1959, Spring wheat, Barley - 0.0200; Beans - 0.0191.

Treatments:

Crop sequences for each series:

1st year:	WW	SW	O	WW	B	WW
2nd year:	WW	WW	WW	O	WW	O
3rd year:	SW	SW	SW	SW	B	Be

WW = Winter wheat, SW = Spring wheat, O = Oats, B = Barley, Be = Beans.

In the 4th year the plots are split for N and all cropped with winter wheat, the series starting in 1958 falling due for this treatment this year, and receiving N at 0.5, 1.0 cwt per acre in 2 doses on Mar 22 and May 8, 1961 as 'Nitro-Chalk'.

Basal dressing: 2 cwt compound fertiliser (16% P_2O_5 , 16% K_2O) per acre combine drilled with seed (placed in sideband for beans); all blocks received 23 cwt ground chalk per acre in Nov 1956 and 54 cwt per acre in Oct 1960.

Nitrogen for cereals: 0.46 cwt N as 'Nitro-Chalk' 21 per acre to spring wheat and 0.31 cwt N as 'Nitro-Chalk' 21 per acre to barley, all in seedbed.

Cultivations, etc.: Ground chalk applied: Oct 4 - 14, 1960. Ploughed: Oct 14. Winter wheat combine drilled at 3 bushels per acre: Jan 23, 1961. 'Nitro-Chalk' applied to spring wheat and barley: Mar 7. Beans placement drilled at 200 lb per acre, barley combine drilled at 2 bushels per acre, and spring wheat at 3 bushels per acre: Mar 8. Winter wheat sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 10. Spring wheat and barley sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 11. Beans sprayed with demeton-methyl at 12 fluid oz in 60 gallons per acre: June 14. Combine harvested: Barley - Aug 23; winter and spring wheat - Aug 31; beans - Sept 4. Varieties: Beans - Gartons Tick; winter wheat - Cappelle; spring wheat - Koga II; barley - Proctor.

Note. Estimates of plant height, % area lodged, incidence of Eyespot (*Cercospora herpotrichoides*) and Take-all (*Ophiobolus graminis*) and counts of plants and shoots were made.

61/C/1.2

Errata to "Results of the Field Experiments" 1960.

Page 60/Cd/1.1 the 6th line should read ".... series starting 1958, and 1959, all cereals - 0.0200".

Standard errors per plot, Grain (at 85% dry matter):

Series starting:

1958 Winter wheat

Whole plot: 2.03 cwt per acre or 7.1% (10 d.f.)

Sub plot: 2.53 cwt per acre or 8.8% (12 d.f.)

1959 Spring wheat 1.58 cwt per acre or 4.7% (6 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

Series starting in 1958

Winter wheat

Crop in 1958	WW	SW	O	WW	B	WW	
1959	WW	WW	WW	O	WW	O	
1960	SW	SW	SW	SW	B	Be	Mean
N cwt per acre	(±1.45) ⁽¹⁾ (±1.56) ⁽²⁾						
0.5	28.2	26.4	20.0	22.9	19.6	39.7	26.2
1.0	30.3	28.7	27.4	27.5	24.4	47.1	30.9
Mean (±1.18)	29.3	27.6	23.7	25.2	22.0	43.4	28.5
Diff. (±2.07)	+2.1	+2.3	+7.4	+4.6	+4.8	+7.4	+4.7 (±0.84)
Mean dry matter % as harvested:	85.1						

(1) for use in vertical and interaction comparisons

(2) for use in horizontal and diagonal comparisons

Grain (at 85% dry matter): cwt per acre

Series starting in 1959

Spring wheat

Crop in 1959	WW	SW	O	WW		Barley	Spring beans
1960	WW	WW	WW	O	Mean	B W	W O
	32.7	33.4	32.3	36.7	33.8	35.3	23.9
	(±0.90)						
Mean dry matter % as harvested:	85.9					84.7	78.1