


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/C/2 One Year Leys for Wheat - Ryegrass Ley

Rothamsted Research

Rothamsted Research (1962) *61/R/C/2 One Year Leys for Wheat - Ryegrass Ley* ; Yields Of The Field Experiments 1961, pp 90 - 91 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/C/2.1

ONE YEAR LEYS FOR WHEAT

The comparison of various clover and grass leys as a preparation for wheat - Stackyard 1961.

Design: 4 randomised blocks of 18 plots each.

Area of each plot (acres): 0.0159. Area harvested: 1st cut - 0.0045; 2nd and 3rd cuts - 0.0114.

Treatments:

Nitrogen to leys 1961:- 3 plots per block of each of the following 6 treatments:

To clover: None (Co)

To ryegrass: None (Ro); 1 cwt (R1); 2 cwt (R2) N per acre.

To clover-ryegrass: None (CRo); 1cwt (CR1) N per acre.

The nitrogen was applied as 'Nitro-Chalk', 0.625 and 1.25 cwt N in spring, and 0.375 and 0.75 cwt N after 1st silage cut.

Note: The experiment is designed to include three rates of N applied to wheat in 1961/62.

Basal dressings per acre:

To barley nurse crop 1960: 3 cwt compound fertiliser (16% N, 9% P₂O₅, 9% K₂O) combine drilled.

To leys, combine drilled in seedbed 1960: 1½ cwt superphosphate.

Cultivations, etc.: Barley combine drilled, leys combine drilled, ryegrass at 30 lb, clover at 12 lb and clover-ryegrass at 10 lb clover and 20 lb ryegrass, per acre: Apr 19, 1960. 'Nitro-Chalk' dressings applied: Mar 13 and May 17, 1961. Cut three times for silage: May 15, July 6 and Sept 18. Varieties: Italian ryegrass S22 and Dorset Marl broad red clover.

Standard error per plot, Grass dry matter cwt per acre:*

1st cut: 3.34 cwt per acre or 6.7% (62 d.f.)*

2nd cut: 2.25 cwt per acre or 9.7% (63 d.f.)

3rd cut: 1.42 cwt per acre or 11.3% (63 d.f.)

Total of 3 cuts: 3.88 cwt per acre or 4.5% (63 d.f.)

*₁ missing value.

61/C/2.2

Summary of Results

Grass dry matter: cwt per acre

<u>Nitrogen to leys 1961</u>						
G_0	R_0	R_1	R_2	CR_0	CR_1	Mean
<u>1st cut</u>						
28.5	39.6	62.5	59.6 ⁺	51.3	59.1	50.1
		(±0.97)				
<u>2nd cut</u>						
28.2	9.9	25.2	32.9	15.8	26.7	23.1
		(±0.65)				
<u>3rd cut</u>						
23.7	4.9	6.7	10.5	19.1	10.9	12.6
		(±0.41)				
<u>Total of 3 cuts</u>						
80.4	54.3	94.3	102.9	86.2	96.8	85.8
		(±1.12)				

+ Includes 1 estimated value.

Mean dry matter % as harvested: 1st cut: 21.8
 2nd cut: 30.7
 3rd cut: 31.6
 Total of 3 cuts: 28.0