


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1962

[Full Table of Content](#)



62/R/C/8 and 62/W/C/8 Long-term Liming - Spring Beans (Test Crop)

Rothamsted Research

Rothamsted Research (1963) *62/R/C/8 and 62/W/C/8 Long-term Liming - Spring Beans (Test Crop)* ;
Yields Of The Field Experiments 1962, pp 115 - 117 - DOI:
<https://doi.org/10.23637/ERADOC-1-164>

62/0/8.1

LONG TERM LIMING EXPERIMENT - SPRING BEANS 1962

The effect of lime on the yield and composition of crops and on the status of P and K in soils - Rothamsted (R) Sawyers I and Woburn (W) Stackyard Series C 1962, the first year.

Design (each field): 2 blocks of 16 plots each, columns of 4 plots being split for uninoculated v. inoculated seed.

Area of each sub plot: 0.0145 acres. Area harvested: 0.0093 acres.

Treatments. All combinations of:-

Ground chalk: None; 2; 4; 6 tons CaCO_3 per acre applied in March 1962.

P: None; 0.5 cwt P_2O_5 per acre as superphosphate.

K: None; 1.0 cwt K_2O per acre as muriate of potash.

Inoculation of seed: None; Inoculated with *Rhizobium leguminosarum*.

The maximum pH range between plots was 4.8 - 5.6 on Sawyers I (R) and 5.6 - 6.3 on Stackyard Series C (W).

Basal dressing: None.

Cultivations, etc.:-

Sawyers I (R): Ploughed: Nov 17, 1961. Ground chalk applied: Mar 5, 1962. Rotary cultivated: Mar 14. Superphosphate and muriate of potash applied: Mar 15. Seed drilled at 200 lb per acre: Mar 16. Sprayed with simazine at 1 lb in 40 gallons per acre: Apr 11. Sprayed with demeton methyl at 6 fluid oz in 60 gallons per acre: July 6. Combine harvested: Sept 20. Variety: Tick 30B. Previous crops: Potatoes and fallow 1960, potatoes and fallow 1961.

Stackyard Series C (W): Ploughed: Nov 24, 1961. Ground chalk applied: Mar 9, 1962. Superphosphate and muriate of potash applied: Mar 15. Seed drilled at 200 lb per acre: Mar 19. Sprayed with simazine at 1 lb in 40 gallons per acre: Apr 12. Sprayed with demeton methyl at 6 fluid oz in 60 gallons per acre: July 12. Combine harvested: Sept 20. Variety: Tick 30B. Previous crops: Barley 1960, sugar beet 1961.

Note: Samples were taken for counts of pods and beans.

Standard errors per plot. Grain (at 85% dry matter):

Sawyers I (R) Whole plot: 4.26 cwt per acre or 25.4% (15 d.f.)
Sub plot: 1.42 cwt per acre or 8.4% (16 d.f.)

Stackyard Series C (W) Whole plot: 2.44 cwt per acre or 13.1% (15 d.f.)
Sub plot: 2.79 cwt per acre or 14.9% (16 d.f.)

62/C/8.2

Summary of ResultsSawyers I (R)Grain (at 85% dry matter): cwt per acre

	Ground chalk: tons per acre				Mean
	None	2	4	6	
Mean (± 1.51)	12.3	16.0	20.3	18.6	16.8
P ₂ O ₅ : cwt per acre		(± 2.13)			
None	12.7	17.2	21.2	17.5	17.2
0.5	12.0	14.7	19.5	19.7	16.5
Diff. (± 3.01)	-0.7	-2.5	-1.7	+2.2	-0.7
					(± 1.51)
K ₂ O: cwt per acre					
None	13.8	15.2	19.1	18.7	16.7
1.0	10.9	16.8	21.6	18.5	16.9
Diff. (± 3.01)	-2.9	+1.6	+2.5	-0.2	+0.2
					(± 1.51)
Inoculation of seed		(± 1.55) ¹			
None	12.1	15.9	19.7	18.4	16.5
Inoculated	12.5	16.1	20.9	18.8	17.1
Diff. (± 0.71) ²	+0.4	+0.2	+1.2	+0.4	+0.6
	P ₂ O ₅ : cwt per acre		K ₂ O: cwt per acre		
	None	0.5	None	1.0	
K ₂ O: cwt per acre	(± 1.51)				
None	16.7	16.7			
1.0	17.6	16.3			
Inoculation of seed	(± 1.10) ¹		(± 1.10) ¹		
None	17.0	16.1	16.5	16.6	
Inoculated	17.3	16.8	16.9	17.3	

Mean dry matter % as harvested: 76.2

(1) For use only in horizontal comparisons.

(2) For use only in testing the difference of 2 differences.

62/C/8.3

Stackyard Series C (W)Grain (at 85% dry matter): cwt per acre

	Ground chalk: tons per acre				Mean
	None	2	4	6	
Mean (± 0.86)	14.8	19.0	19.1	22.0	18.7
P ₂₅ ⁰ : cwt per acre		(± 1.22)			
None	13.1	19.2	19.4	21.7	18.3
0.5	16.6	18.8	18.8	22.3	19.1
Diff. (± 1.73)	+3.5	-0.4	-0.6	+0.6	+0.8
K ₂₀ : cwt per acre					(± 0.86)
None	14.9	19.1	20.2	23.8	19.5
1.0	14.8	19.0	18.0	20.2	18.0
Diff. (± 1.73)	-0.1	-0.1	-2.2	-3.6	-1.5
Inoculation of seed		(± 1.11) ¹			(± 0.86)
None	14.7	20.2	18.0	21.6	18.6
Inoculated	14.9	17.9	20.2	22.4	18.8
Diff. (± 1.40) ²	+0.2	-2.3	+2.2	+0.8	+0.2
P ₂₅ ⁰ : cwt per acre K ₂₀ : cwt per acre					
	None	0.5	None	1.0	
K ₂₀ : cwt per acre		(± 0.86)			
None	20.2	18.8			
1.0	16.5	19.4			
Inoculation of seed		(± 0.79) ¹		(± 0.79) ¹	
None	18.7	18.5	18.8	18.4	
Inoculated	18.0	19.7	20.2	17.5	

Mean dry matter % as harvested: 77.6

(1) For use only in horizontal comparisons

(2) For use only in testing the differences of 2 differences