


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1962

[Full Table of Content](#)



62/R/DD/1 and 62/W/DD/1 Potatoes - Control of Weeds

Rothamsted Research

Rothamsted Research (1963) *62/R/DD/1 and 62/W/DD/1 Potatoes - Control of Weeds ; Yields Of The Field Experiments 1962*, pp 134 - 136 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/Da/1.1

POTATOES

Control of weeds by residual herbicides - Rothamsted (R) Great Knott I and Woburn (W) Great Hill 1962.

Design: 4 randomised blocks of 10 plots each, plots on Great Knott I (R) being split for 0 v final earthing up.

Area of each plot (acres): 0.0216 Area harvested:
R - 0.0072
W - 0.0096

Treatments (in lb active material in 40 gallons per acre):-

None	(O)
Mechanical cultivations	(M)
2,6-DBN: 1, 2 lb	(DB1; DB2)
Dinoseb: 4, 6 lb	(DS1; DS2)
Prometryne: $1\frac{3}{4}$; $2\frac{1}{2}$ lb	(Ta1; Ta2)
Trietazine: $1\frac{1}{2}$ lb	(Te)
Trietazine: $1\frac{1}{2}$ lb + paraquat: $\frac{3}{4}$ lb**	(TeP)

In addition plots on Great Knott I (R) were split for 0 v final earthing up (E).

* $1\frac{1}{4}$ lb on Great Hill (W)

** paraquat spray repeated on Great Hill (W).

Note: On Great Knott I (R) weeds and haulm on 0, DB1 and DB2 plots were mown on July 31, and the yields are omitted from the analysis.

Basal dressing: 8 cwt compound fertiliser (17% N, 11% P₂O₅, 22% K₂O).

Cultivations, etc.:

Great Knott I (R): Ploughed: Dec 26, 1961. Rotary cultivated, basal dressing applied: Apr 16, 1962. Potatoes planted: Apr 26. Ridges rolled: Apr 28. DB and Te sprays applied: May 9. M plots chain harrowed: May 11. M plots grubbed: May 14. DS, Ta and TeP sprays applied: May 30. M plots re-ridged: June 6. M plots grubbed: June 18. E sub-plots grubbed and earthed up: July 5. Sprayed with maneb at $1\frac{1}{2}$ lb active material in 18 gallons per acre: July 18. Weeds and haulm mown on 0 and DB plots: July 31. Sprayed with copper fungicide at 5 lb in 20 gallons per acre: Aug 9 and again at 5 lb in 40 gallons per acre: Aug 24. Sprayed with undiluted BOV at 18 gallons per acre: Sept 18. Lifted: Oct 8. Variety: King Edward. Previous crops: Winter wheat 1960, barley 1961.

62/Da/1.2

Great Hill (W): Subsoiled: Oct 26, 1961. Ploughed: Oct 31.
Basal dressing applied: Apr 7, 1962. Potatoes planted: Apr 9.
Ridges rolled: Apr 12. DB, Te and TeP sprays applied: Apr 19.
Ta, DS sprays applied and second paraquat spray of $\frac{3}{4}$ lb in 40
gallons per acre applied to TeP plots: May 16. Cultivations to
M plots: chain harrowed - May 1; harrowed with weeder - May 5;
grubbed - May 14; harrowed - May 26; re-ridged - May 24;
grubbed - May 31 and June 15; earthed up - June 21. Sprayed
with copper oxychloride at 2.3 lb copper in 40 gallons per acre:
July 24. Tops and weeds burnt off with paraquat at 0.9 lb
in 40 gallons per acre: Aug 9. Haulm destroyed mechanically:
Aug 22. Lifted: Sept 5. Variety: Majestic. Previous crops:
Potatoes 1960, barley 1961.

Note: At Woburn dry weights of weeds were estimated on plots of
certain treatments.

Standard errors per plot. Total tubers:

Great Knott I (R): Whole plot: 1.414 tons per acre or 10.4% (18 d.f.)
Sub plot: 1.311 tons per acre or 9.6% (21 d.f.)
Great Hill (W): 1.086 tons per acre or 18.1% (27 d.f.)

62/Da/1.3

Summary of ResultsTreatments

Final cultivation	O	M	DB1	DB2	DS1	DS2	Ta1	Ta2	Te	TeP	Mean
<u>Total tubers: tons per acre</u>											
<u>Great Knott I (R)</u>											
$(\pm 0.655)^1 (\pm 0.845)^2$											
None	+	14.66			12.88	14.62	12.34	14.40	12.37	13.77	13.57
Earthed up		14.22			13.83	14.48	12.75	14.18	12.46	13.44	13.62
Mean (± 0.736)		14.44			13.35	14.55	12.54	14.29	12.42	13.60	13.59
Diff. (± 0.927)		-0.44			+0.95	-0.14	+0.41	-0.22	+0.09	-0.33	+0.05
<u>Great Hill (W)</u>											
Mean (± 0.543)	2.21	7.49	5.75	6.36	6.32	6.50	4.91	5.71	5.79	8.98	6.00
<u>Percentage ware $(1\frac{1}{2}"$ riddle)</u>											
<u>Great Knott I (R)</u>											
None	+	92.9			92.9	94.7	93.4	94.4	93.3	92.9	93.5
Earthed up		94.2			94.6	94.0	92.6	93.9	92.7	92.9	93.5
Mean		93.6			93.8	94.4	93.0	94.1	93.0	92.9	93.5
Diff.		+1.3			+1.7	-0.7	-0.8	-0.5	-0.6	0.0	0.0
<u>Great Hill (W)</u>											

Not recorded

(1) for use in vertical and interaction comparisons

(2) for use in horizontal and diagonal comparisons

* $\frac{1}{4}$ lb on Great Hill (W)

** paraquat spray repeated on Great Hill (W)

Treatments

O = None

M = Mechanical cultivations

DB1; DB2 = 2,6-DBN: 1, 2 lb

DS1; DS2 = Dinoseb: 4, 6 lb

Ta1; Ta2 = Prometryne: $\frac{3}{4}$; $2\frac{1}{2}$ lbTe = Trietazine: $1\frac{1}{2}$ lbTeP = Trietazine: $1\frac{1}{2}$ lb + paraquat: $\frac{3}{4}$ lb

+ crop failed