

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1962

[Full Table of Content](#)



62/R/B/7 Concentrated Fertilisers Rotation

Rothamsted Research

Rothamsted Research (1963) *62/R/B/7 Concentrated Fertilisers Rotation ; Yields Of The Field Experiments 1962*, pp 86 - 87 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/B/7.1

CONCENTRATED FERTILISER ROTATION

Concentrated compound fertiliser and forms of N - West Barnfield I
1962, the third year.

Rotation: Kale, ryegrass, barley.

Design (each crop): 2 randomised blocks of 14 plots each.

Area of each plot (acres): 0.0174. Area harvested: Kale - 0.0084,
ryegrass - 0.0056, barley - 0.0114.

Treatments (per acre): No fertiliser (O)
P₂O₅ and K₂O each at 0.3 cwt to barley and each at 1.0 cwt to
kale and ryegrass, as triple superphosphate and potassium
bicarbonate. (B)
Compound fertiliser, 20% N, 10% P₂O₅, 10% K₂O at 0.3(1),
0.6(2) cwt N to barley and 1.0(1), 2.0(2) to kale and
ryegrass. (F)
Sulphate of ammonia, granular superphosphate and muriate of
potash at rates equivalent to treatments F(1) and (2). (P)
PK as treatment B plus
Sulphate of ammonia (S)
Calcium nitrate (C)
Urea (U)
Ammonium nitrate (A)
each at rates 1 and 2 of N.

Basal dressing: None.

Cultivations, etc.: Ploughed: Dec 13, 1961. Fertilisers broadcast
for barley, barley drilled at 2½ bushels per acre: Feb 21, 1962.
Fertilisers broadcast for ryegrass, ryegrass sown at 30 lb per acre:
Apr 13. Fertilisers applied for kale: Apr 24. Kale drilled at
3 lb per acre: Apr 25. Barley sprayed with MCPA/TBA at 4 pints
in 40 gallons per acre: May 30. Grass cut twice: July 30 and
Oct 1. Barley combine harvested: Aug 27. Kale harvested:
Nov 6 and 15. Varieties: Kale - Thousand head (Canson);
ryegrass - S22; barley - Proctor.

Note: For previous year's results see "Results of the Field Experiments"
60/B/8 and 61/B/7.

Standard errors per plot.

Kale, fresh weight: 0.973 tons per acre or 4.6% (13 d.f.)

Ryegrass, dry matter:

1st cut: 2.53 cwt per acre or 11.3% (13 d.f.)

2nd cut: 2.71 cwt per acre or 13.6% (13 d.f.)

Total of 2 cuts: 3.06 cwt per acre or 7.2% (13 d.f.)

Barley, grain (at 85% dry matter): 1.86 cwt per acre or 6.0%
(13 d.f.)

62/B/7.2

Summary of Results

Fertiliser	Kale fresh weight: tons per acre	Ryegrass dry matter: cwt per acre			Barley (at 85% dry matter): cwt per acre	
		1st cut	2nd cut	Total of 2 cuts	Grain	Straw
	(±0.688)	(±1.79)	(±1.91)	(±2.16)	(±1.31)	
O	10.69	5.3	4.6	9.9	12.1	5.8
B	12.73	8.0	5.4	13.4	13.2	7.0
F ₁	21.82	21.7	19.3	41.0	31.3	20.1
F ₂	25.11	28.1	25.8	53.8	38.1	29.3
P ₁	21.08	20.2	20.9	41.1	30.6	20.6
P ₂	24.87	24.0	23.2	47.2	36.5	29.4
S ₁	17.95	24.7	15.5	40.2	27.1	18.1
S ₂	23.24	24.7	31.7	56.4	38.2	29.9
C ₁	20.75	23.2	18.3	41.5	30.7	22.3
C ₂	26.31	27.3	26.7	54.0	37.4	28.8
U ₁	21.10	23.7	17.3	41.0	27.7	19.1
U ₂	22.68	25.4	25.6	51.0	39.7	28.8
A ₁	20.74	27.0	18.7	45.7	30.3	20.9
A ₂	24.08	29.3	27.3	56.6	40.8	30.1
Mean	20.94	22.3	20.0	42.3	30.9	22.1
Mean dry matter % as harvested:		17.6	20.7	19.2	81.0	67.5

Treatments

O = No fertiliser

B = P₂O₅ and K₂O each at 0.3 cwt to barley and each at 1.0 cwt to kale and ryegrass, as triple superphosphate and potassium bicarbonate.F = Compound fertiliser, 20% N, 10% P₂O₅, 10% K₂O at 0.3(1), 0.6(2) cwt N to barley and 1.0(1), 2.0(2) to kale and ryegrass.

P = Sulphate of ammonia, granular superphosphate and muriate of potash at rates equivalent to treatments F (1) and (2).

S = Sulphate of ammonia Plus PK as treatment B.

C = Calcium nitrate " " " " "

U = Urea " " " " "

A = Ammonium nitrate " " " " "