


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1963

[Full Table of Content](#)



63/R/RP/B/8 Residual Phosphate Rotation

Rothamsted Research

Rothamsted Research (1964) *63/R/RP/B/8 Residual Phosphate Rotation ; Yields Of The Field Experiments 1963*, pp 109 - 112 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/B/8.1

RESIDUAL PHOSPHATE ROTATION

(RP)

The long term and residual effects of a number of phosphate fertilisers compared with superphosphate - Great Field IV and Sawyers I 1963, the fourth year.

Rotation: Potatoes, barley, swedes.

Note: Swede tops are ploughed in.

Design: Great Field IV: 1 randomised block of 12 plots per crop.
Sawyers I: 2 randomised blocks of 12 plots per crop.

Area of each plot (acres):

Great Field IV: 0.0193. Area harvested: Potatoes and barley
- 0.0129, swedes - 0.0096.

Sawyers I: 0.0212. Area harvested: Potatoes and barley
- 0.0141, swedes - 0.0106.

Treatments:

Granular superphosphate treatments broadcast in spring before sowing or ridging:-

1. No phosphate.
2. 0.25 cwt P_{205} per acre per year.
3. 0.50 cwt P_{205} per acre per year.
4. 0.75 cwt P_{205} per acre in 1962.
5. 1.50 cwt P_{205} per acre in 1962.

Phosphate fertilisers ploughed in at 3.0 cwt P_{205} per acre in September 1959.

6. Nitrophosphate I (17.1% P_{205} , none water soluble).
7. Nitrophosphate II (18.8% P_{205} , one quarter water soluble).
8. Nitrophosphate III (22.4% P_{205} , half water soluble).
9. Gafsa rock phosphate (28.9% P_{205}).
10. Bessemer basic slag (15.2% P_{205}).
11. Potassium metaphosphate (57.9% P_{205} , 38.8% K_2O).
12. Granular superphosphate (20.4% P_{205}).

Note: To balance the K_2O content of potassium metaphosphate, all the other treatments included 2.0 cwt K_2O per acre as sulphate of potash in autumn 1959.

Basal dressings per acre: Broadcast in spring before sowing or ridging:

N as 'Nitro-Chalk' 21:-

To potatoes: 1.2 cwt, to barley: None on Great Field IV, 0.6 cwt on Sawyers I, to swedes: 0.5 cwt.

K_2O as sulphate of potash:-

To potatoes: 1.0 cwt, to barley: 1.0 cwt, to swedes: 1.0 cwt.

63/B/8.2

Cultivations, etc. (both fields, except as indicated): Ploughed:

Great Field IV - Mar 27, 1963, Sawyers I - Apr 3.

Potatoes: Fertilisers applied, potatoes planted: May 7, 1963.

Earthed up: Great Field IV - June 28, Sawyers I - July 4.

Sprayed with maneb at 1.2 lb in 20 gallons per acre: July 10.

Great Field IV sprayed with copper oxychloride fungicide at 2.3 lb

copper plus 0.35 pints menazon in 20 gallons per acre: Aug 14.

Sawyers I sprayed with copper oxychloride fungicide at 2.3 lb

copper in 40 gallons per acre: Sept 3. Sprayed with undiluted

BOV at 16 gallons per acre: Sept 23. Lifted: Oct 14. Variety:

Majestic.

Barley: Fertilisers applied: Apr 11, 1963. Seed drilled at

2 bushels per acre: Apr 13. Sprayed with mecoprop/2,4-D at

6 pints in 40 gallons per acre: June 12. Combine harvested:

Sept 13. Variety: Proctor.

Swedes: Ground chalk applied to Sawyers I at 23 cwt per acre:

Apr 23, 1963. Fertilisers applied: Sawyers I - May 18,

Great Field IV - May 21. Seed drilled at 1.5 lb per acre:

May 22. Singled: July 3. Lifted: Nov 6. Variety:

Wilhelmsburger.

Note: For details of previous years' results see 'Results of the
Field Experiments' 60/B/9, 61/B/8 and 62/B/8.

Standard errors per plot.

Sawyers I

Potatoes, Total tubers: 0.695 tons per acre or 7.7% (11 d.f.)

Barley, Grain (at 85% dry matter): 1.71 cwt per acre or 4.4%
(11 d.f.)

Swedes, Roots: 2.125 tons per acre or 11.1% (11 d.f.)

63/B/8.3

Summary of Results

Phosphate		<u>Potatoes</u>				<u>Percentage ware (1.5 inch riddle)</u>			
		<u>Total tubers: tons per acre</u>							
		Great Field IV		Sawyers I		Great Field IV		Sawyers I	
		Mean	Increase	Mean	Increase	Mean	Increase	Mean	Increase
				(±0.491) (±0.695)					
None	1	11.26		7.75		95.5		93.7	
	2	12.35	+1.09	9.01	+1.26	93.2	-2.3	93.5	-0.2
	3	13.49	+2.23	8.97	+1.22	94.5	-1.0	94.6	+0.9
	4	10.81	-0.45	8.05	+0.30	95.0	-0.5	94.3	+0.6
	5	12.45	+1.19	8.88	+1.13	93.6	-1.9	93.0	-0.7
	6	13.11	+1.85	9.52	+1.77	93.4	-2.1	94.3	+0.6
	7	12.42	+1.16	9.09	+1.34	93.0	-2.5	94.0	+0.3
	8	12.49	+1.23	9.65	+1.90	93.9	-1.6	94.7	+1.0
	9	12.75	+1.49	9.03	+1.28	94.8	-0.7	94.8	+1.1
	10	12.93	+1.67	9.25	+1.50	96.0	+0.5	95.6	+1.9
	11	12.40	+1.14	9.35	+1.60	94.5	-1.0	94.2	+0.5
	12	11.55	+0.29	9.31	+1.56	95.6	+0.1	93.4	-0.3
Mean		12.33		8.99		94.4		94.1	

		<u>Barley</u>				<u>Straw (at 85% dry matter):</u>			
		<u>Grain (at 85% dry matter):</u>				<u>cwt per acre</u>			
				(±1.21) (±1.71)					
None	1	25.1		35.5		21.9		22.9	
	2	33.9	+8.8	40.1	+4.6	29.6	+7.7	29.3	+6.4
	3	27.9	+2.8	38.5	+3.0	30.6	+8.7	27.8	+4.9
	4	34.2	+9.1	37.7	+2.2	30.6	+8.7	26.7	+3.8
	5	35.5	+10.4	38.3	+2.8	31.1	+9.2	26.0	+3.1
	6	30.3	+5.2	36.9	+1.4	27.0	+5.1	25.2	+2.3
	7	30.6	+5.5	39.6	+4.1	29.6	+7.7	30.1	+7.2
	8	25.8	+0.7	39.9	+4.4	29.9	+8.0	27.7	+4.8
	9	34.5	+9.4	39.7	+4.2	32.4	+10.5	26.6	+3.7
	10	25.8	+0.7	38.9	+3.4	24.4	+2.5	28.9	+6.0
	11	32.0	+6.9	38.2	+2.7	31.3	+9.4	30.6	+7.7
	12	33.4	+8.3	39.4	+3.9	27.2	+5.3	27.4	+4.5
Mean		30.8		38.5		28.8		27.4	

Mean dry matter
% as harvested

78.8

80.1

62.2

75.6

63/B/8.4

Phosphate		Great Field IV		Sawyers I	
		Mean	Increase	Mean	Increase
<u>Swedes, Roots: tons per acre</u>					
		(+1.503)(+2.125)			
None	1	11.81		11.24	
	2	20.65	+8.84	19.87	+8.63
	3	25.28	+13.47	23.24	+12.00
	4	18.20	+6.39	16.06	+4.82
	5	28.75	+16.94	21.66	+10.42
	6	24.40	+12.59	21.47	+10.23
	7	25.56	+13.75	21.51	+10.27
	8	23.89	+12.08	21.20	+9.96
	9	23.66	+11.85	17.24	+6.00
	10	22.18	+10.37	19.89	+8.65
	11	21.44	+9.63	18.69	+7.45
	12	23.80	+11.99	17.49	+6.25
Mean		22.47		19.13	

For explanation of treatment symbols see page 63/B/8.1