


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1963

[Full Table of Content](#)



63/W/WCW/B/11 Weedkiller-cultivation Rotation

Rothamsted Research

Rothamsted Research (1964) *63/W/WCW/B/11 Weedkiller-cultivation Rotation ; Yields Of The Field Experiments 1963*, pp 123 - 125 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/B/11.1

CULTIVATION - WEEDKILLER ROTATION

(WCW)

A comparison of weed control by various cultivation methods and by a pre-emergence weedkiller - Woburn Great Hill I and II 1963, the fourth year.

For history, rotation etc., see 'Numerical Results of the Field Experiments' 60/B/11, 61/B/11 and 62/B/11.

Area of each plot (acres): 0.0482. Area harvested: Potatoes - 0.0107, barley - 0.0115.

Treatments.

Potatoes: All combinations of:-

Initial cultivations: Ploughed (P), rotary cultivated (R), rigid-tine cultivated (T).

Treatments after planting: Normal cultivations (M), prometryne* plus paraquat* applied immediately before crop emergence (S) - duplicate plots.

Barley: All combinations of:-

Initial cultivations: As for potatoes.

Prometryne to potatoes 1962: None (O), after planting (X), after early cultivations (Y).

*Prometryne at 2 lb, plus 0.75 lb paraquat, in 40 gallons per acre.

Basal dressings per acre:

Potatoes: 10 cwt compound fertiliser (17% N, 11% P_{205} , 22% K_{20}).

Barley: 3.5 cwt compound fertiliser (16% N, 9% P_{205} , 9% K_{20}) combine drilled.

Cultivations, etc.:

Potatoes: T plots rigid-tine cultivated twice: Nov 6, 1962. R plots rotary cultivated (one stroke), P plots ploughed: Nov 9. Ground chalk applied at 62 cwt per acre: Feb 20, 1963. P and T plots spring-tine cultivated: Apr 18. Basal dressing applied: Apr 22. P and T plots spring-tine cultivated: Apr 23. R plots rotary cultivated, potatoes machine planted: Apr 24. Ridges rolled: Apr 27. M plots chain harrowed: May 9. Prometryne and paraquat treatment applied: May 24. M plots grubbed: May 25 and June 18. M plots earthed up: June 24. Sprayed with copper oxychloride fungicide at 2.3 lb Cu in 20 gallons per acre: July 25, and again at the same rate, plus 0.35 pints of menazon (against aphids) in

63/B/11.2

30 gallons per acre: Aug 22. Haulm destroyed with diquat at 0.9 pints in 40 gallons per acre: Sept 23. Lifted: Oct 10. Variety: Majestic.

Barley: T plots rigid-tine cultivated on 2 occasions: Nov 6 and 9, 1962. R plots rotary cultivated, P plots ploughed: Nov 9. Ground chalk applied at 62 cwt per acre: Feb 20, 1963. P and T plots spring-tine cultivated, R plots rotary cultivated: Apr 11. Seed drilled at 2.3 bushels per acre: Apr 18. Sprayed with mecoprop/2,4-D at 6 pints in 40 gallons per acre: May 21. Combine harvested: Sept 9. Variety: Proctor.

Standard errors per plot.

Potatoes, total tubers: 1.702 tons per acre or 13.5% (11 d.f.)

Barley, grain (at 85% dry matter): 1.63 cwt per acre or 6.7% (8 d.f.)

63/B/11.3

Summary of ResultsPotatoes

Treatments after planting	Initial cultivation			Mean
	P	R	T	
<u>Total tubers: tons per acre</u>				
M (±1.204)	11.48	11.86	10.89	11.41 (±0.695)
S (±0.851)	13.04	12.54	14.14	13.24 (±0.491)
Mean (±0.695)	12.52	12.31	13.05	12.63

Percentage ware (1.5 inch riddle)

M	81.0	80.5	80.5	80.6
S	82.9	81.4	85.8	83.4
Mean	82.3	81.1	84.0	82.4

Barley

Prometryne to potatoes 1962	Initial cultivation			Mean
	P	R	T	
<u>Grain (at 85% dry matter): cwt per acre</u>				
		(± 1.15)		(± 0.67)
O	24.4	24.4	23.6	24.2
X	25.8	24.8	24.3	25.0
Y	23.6	23.7	23.1	23.5
Mean (± 0.67)	24.6	24.3	23.7	24.2

Mean dry matter % as harvested: 79.4

For explanation of treatment symbols see page 63/B/11.1