


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1963

[Full Table of Content](#)



63/R/IB/C/2 Intensive Barley Growing - Cereals

Rothamsted Research

Rothamsted Research (1964) *63/R/IB/C/2 Intensive Barley Growing - Cereals* ; Yields Of The Field Experiments 1963, pp 133 - 134 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/C/2.1

INTENSIVE BARLEY GROWING EXPERIMENT

(IB)

Little Knott I - 1963, the third year

For treatments etc., see 'Numerical Results of the Field Experiments' 61/C/8.

Area of each plot (acres): 0.0212. Area harvested: 0.0139.

Cultivations, etc.: Ploughed: Sept 26, 1962.

Spring beans: Seed placement drilled at 200 lb per acre:

Apr 9, 1963. Sprayed with demeton methyl at 6 fluid oz in 40 gallons per acre: June 14. Combine harvested: Oct 3.

Variety: Tick 30B.

Oats: Seed combine drilled at 4 bushels per acre: Apr 9, 1963.

'Nitro-Chalk' applied: Apr 18. Sprayed with TBA/MCPA at 4 pints in 40 gallons per acre: May 22. Combine harvested: Sept 9. Variety: Condor.

Spring wheat: Seed combine drilled at 3 bushels per acre:

Apr 11, 1963. 'Nitro-Chalk' applied: Apr 18. Sprayed with TBA/MCPA at 4 pints in 40 gallons per acre: May 22. Combine harvested: Sept 12. Variety: Jufy I.

Barley: Seed combine drilled at 2.5 bushels per acre: Apr 9, 1963.

'Nitro-Chalk' applied: Apr 18. Sprayed with TBA/MCPA at 4 pints in 40 gallons per acre: May 22. Combine harvested: Sept 9. Variety: Proctor.

Winter wheat: Seed combine drilled at 2.5 bushels per acre:

Oct 31, 1962. 'Nitro-Chalk' applied: Mar 13, 1963. Sprayed with TBA/MCPA at 4 pints in 40 gallons per acre: May 22. Combine harvested: Sept 10. Variety: Cappelle.

Potatoes: Basal compound fertiliser applied: Apr 30, 1963. Potatoes machine planted: May 6. Earthed up: June 26. Sprayed with undiluted BOV at 16 gallons per acre: Sept 14. Lifted: Sept 23. Variety: Majestic.

Notes: (1) Yields were only taken for sequences 1, 4, 7, 8 and 9.

(2) For details of the previous years' results see 'Numerical Results of the Field Experiments' 61/C/8 and 62/C/7.

Standard errors per plot. Grain (at 85% dry matter)

Spring wheat (4 and 8): 1.37 cwt per acre or 4.2% (9 d.f.)

Barley (1 and 7): 2.05 cwt per acre or 5.8% (6 d.f.)

63/c/2.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Winter wheat 9

Crop in 1961 Crop in 1962	Spring wheat Winter wheat				
N: cwt per acre	None	0.3	0.6	0.9	Mean
	19.5	23.8	28.0	30.6	25.5

Mean dry matter % as harvested: 77.2

Spring wheat 4 and 8

Crop in 1961 Crop in 1962	Spring wheat Spring wheat				Beans Oats	
N: cwt per acre	None	0.3	0.6	0.9	0.6	Mean
	22.1	27.4	32.9	30.7	37.6	32.9

Mean dry matter % as harvested: 73.0

Barley 1 and 7

Crop in 1961 1962	N: cwt per acre				
	None	0.3	0.6	0.9	Mean
	(±1.45)				(±0.73)
Barley Barley	26.2	33.3	35.9	39.7	33.7
Oats Beans	35.0	41.8	40.5	33.1	37.6
Mean (±1.03)	30.6	37.5	38.2	36.4	35.6

Mean dry matter % as harvested: 73.2