


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1963

[Full Table of Content](#)



63/W/WCT101/DE/1 Carrots - Control of Motley Dwarf Virus

Rothamsted Research

Rothamsted Research (1964) *63/W/WCT101/DE/1 Carrots - Control of Motley Dwarf Virus ; Yields Of The Field Experiments 1963*, pp 181 - 182 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/De/1.1

CARROTS

(Wct 101)

The effect of systemic insecticides on yield through control of motley dwarf virus - Woburn Lansome Field 1963.

Design: A plaid rectangle of 4 rows and 8 columns.

Area of each plot: 0.0212 acres. Area harvested: 0.0016 acres.

Treatments. All combinations of:-

Granular systemic insecticide (to columns): None, 0.75 lb per acre menazon drilled just below the seed.

Early spraying: None, sprayed twice*. (E)

Midseason spraying: None, sprayed twice. (M)

Late spraying: None, sprayed twice*. (L)

The spray used was demeton-methyl at 6 fluid oz in 40 gallons per acre.

*Intended treatments, only one application was made.

Basal dressing: 8 cwt compound fertiliser (10% N, 10% P₂O₅, 18% K₂O) per acre.

Cultivations, etc.: Ploughed: Mar 8, 1963. Basal compound fertiliser applied: Apr 22. Menazon granules applied, seed drilled at 2.25 lb per acre: Apr 23. Demeton-methyl treatments applied: E - May 31, M - June 27 and July 8, L - July 22. Lifted: Sept 12. Variety: Scarlet Intermediate. Previous crops: Barley 1961, sugar beet 1962.

Note: Aphid counts and estimates of virus infection were made.

Standard errors per plot.

Roots: 2.328 tons per acre or 11.3% (14 d.f.)

Tops: 0.995 tons per acre or 14.0% (14 d.f.)

Summary of Results

Systemic Insecticide	Times of spraying								Mean
	-	E	M	EM	L	EL	ML	EML	
<u>Roots: tons per acre</u>									
(±1.646)									
None	21.78	20.31	20.86	22.06	21.83	19.70	21.22	20.31	21.01
Menazon	19.06	18.17	19.56	20.73	20.67	22.25	20.03	22.19	20.33
Mean (±1.164)	20.42	19.24	20.21	21.39	21.25	20.98	20.63	21.25	20.67
Diff. (±2.328)*	-2.72	-2.14	-1.30	-1.33	-1.16	+2.55	-1.19	+1.88	-0.68
<u>Tops: tons per acre</u>									
(±0.704)									
None	7.44	7.42	6.86	7.44	7.70	5.91	7.86	6.31	7.12
Menazon	6.45	5.91	7.44	7.70	6.89	7.70	6.59	7.86	7.07
Mean (±0.498)	6.95	6.67	7.15	7.57	7.29	6.81	7.22	7.09	7.09
Diff. (±0.995)*	-0.99	-1.51	+0.58	+0.26	-0.81	+1.79	-1.27	+1.55	-0.05

63/De/1.2

*For use only in testing the difference of 2 differences.

Times of spraying

E = Early spraying: None, sprayed once

M = Midseason spraying: None, sprayed twice

L = Late spraying: None, sprayed once