

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/R/AR/C/4 Methods of Application of Fertiliser 1963-64 - Wheat

Rothamsted Research

Rothamsted Research (1965) *64/R/AR/C/4 Methods of Application of Fertiliser 1963-64 - Wheat ; Yields Of The Field Experiments 1964*, pp 149 - 151 - DOI:

<https://doi.org/10.23637/ERADOC-1-160>

64/C/4.1

METHODS OF APPLICATION OF FERTILISER 1963 - 64

(AR/W)

Methods of application of fertiliser - Great Knott II 1964, the second year - winter wheat.

Design: 3 x 3 x 3 in 3 blocks of 9 plots each together with 3 additional plots per block.

Area of each plot: 0.0212. Area harvested: 0.0126.

Treatments:

3 x 3 x 3: All combinations of:-

To wheat 1964.

1. NPK: None (F0), 0.66 (F1), 1.32 (F2) cwt N as spring top dressings of 'Nitro-Chalk' each with superphosphate and muriate of potash applied in autumn in the seedbed in the proportion 13 N, 13 P2O5, 20 K2O.

To potatoes 1963.

2. Levels of 13:13:20 to supply (cwt):

N	P2O5	K2O
0.66	0.66	1.02 (L1)
1.32	1.32	2.03 (L2)
2.00	2.00	3.07 (L3)

3. Methods of application: Broadcast (B), placed (P), broadcast and rotary cultivated in (BR).

Additional plots:

To wheat 1964. NPK: F0, F1, F2 as above to plots receiving no treatment in 1963.

Basal dressing: None.

Cultivations, etc.: Chisel ploughed: Oct 24, 1963. PK applied, seed drilled at 3 bushels: Oct 28. 'Nitro-Chalk' applied - 1st half dressing: Apr 1, 1964. Sprayed with mecoprop/2,4-D (Methoxone Extra at 7 pints in 40 gals): Apr 30. 'Nitro-Chalk' applied - 2nd half dressing: May 4. Combine harvested: Aug 27. Variety: Cappelle. Previous crop: Spring beans 1962.

NOTES: (1) For previous year's results see 'Results' 63/C/5.

(2) Green-crop and grain and straw samples were taken for NPK and dry matter determination.

Standard error per plot.

Grain: 2.30 or 4.6% (18 d.f.)

64/C/4.2

SUMMARY OF RESULTS

GRAIN							
	B	P	BR	L1	L2	L3	Mean
		(± 1.33)			(± 1.33)		(± 0.77)
F0	42.2	44.9	47.9	39.5	45.2	50.3	45.0
F1	54.5	53.8	52.4	51.0	55.6	54.1	53.6
F2	53.2	56.4	53.5	54.5	55.2	53.4	54.4
					(± 1.33)		
	B			48.3	51.0	50.7	50.0
	P			48.5	53.1	53.4	51.7
	BR			48.2	51.9	53.6	51.2
		Mean (± 0.77)		48.3	52.0	52.6	50.5 (± 0.44)

Plots untreated in 1963

	F0	F1	F2	Mean
	36.0	49.7 (± 1.33)	54.8	46.8 (± 0.77)

General mean: 49.6

Mean D.M. %: 86.3

64/C/4.3

STRAW

	B	P	HR	L1	L2	L3	Mean
F0	28.2	32.2	33.4	28.0	30.4	35.4	31.3
F1	37.0	39.6	34.9	35.7	39.1	36.7	37.2
F2	43.0	46.5	43.7	45.7	43.9	43.6	44.4
		B		32.8	36.7	38.7	36.1
		P		38.6	40.7	39.0	39.4
		HR		37.9	36.0	38.0	37.3
	Mean			36.5	37.8	38.6	37.3

Plots untreated in 1963

	F0	F1	F2	Mean
	22.6	31.6	41.1	31.8

General mean: 35.9

Mean D.M. %: 92.2