

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/R/WAV/C/18 *Trefoil* and Ryegrass Green Manures - Sugar Beet

Rothamsted Research

Rothamsted Research (1965) *64/R/WAV/C/18 Trefoil and Ryegrass Green Manures - Sugar Beet ; Yields Of The Field Experiments 1964*, pp 185 - 189 - DOI:
<https://doi.org/10.23637/ERADOC-1-160>

64/C/18.1

SUGAR BEET AFTER GREEN MANURES

(WAV)

Effects of trefoil and ryegrass green manures and N - Woburn Great Hill
1964, sugar beet following barley.

Design: 2 replicates of 4 x 2 x 4 in 4 blocks of 16 plots each, with
restricted randomisation.

Area of each plot: 0.0159. Area harvested: 0.0106.

Treatments: All combinations of:-

1. Green manures undersown: None (O), trefoil (T), ryegrass (R),
ryegrass sown with 0.6 cwt N as 'Nitro-Chalk' (RN).
2. Time of ploughing: Nov 28, 1963 (A), Jan 30, 1964 (S).
3. Nitrogen: None (NO), 0.6 (N1), 1.2 (N2), 1.8 (N3) cwt N as
'Nitro-Chalk'.

Basal dressings. Barley: 3 cwt 21:14:14 combine drilled. Green
manures: None. Sugar beet: 5 cwt agricultural salt, 0.45 cwt K2O
as muriate of potash, 2.5 cwt 0:20:20.

Cultivations, etc.: Ploughed: Nov 12, 1962. Barley drilled at 2.25
bushels: Apr 8, 1963. Trefoil undersown at 30 lb and ryegrass at
40 lb: May 6. Barley combine harvested: Sept 9. 'Nitro-Chalk'
applied to RN plots: Oct 11. A plots ploughed: Nov 28. S plots
ploughed: Jan 30, 1964. Salt and muriate of potash applied:
Feb 4. Basal PK applied: Apr 6. 'Nitro-Chalk' treatments applied:
Apr 8. Seed drilled at 5 lb: Apr 14. Singled: May 19 - 27.
Lifted: Oct 14. Varieties: Barley - Proctor, trefoil - English,
ryegrass - English Leafy Italian, sugar beet - Klein E. Previous
crops: Barley 1961, potatoes 1962.

NOTE: The green manures were sampled just before ploughing for
estimates of dry matter and N (roots plus tops).

Standard errors per plot:

Roots (washed):	1.122 or 9.3% (29 d.f.)
Total sugar:	4.98 or 10.4% (29 d.f.)
Tops:	0.922 or 14.4% (29 d.f.)

64/c/18.2

SUMMARY OF RESULTS

	NO	N1	N2	N3	Mean
ROOTS					
(± 0.281)					
Mean	8.55	12.50	13.82	13.26	12.03
(± 0.561)					
O	8.56	13.14	13.69	13.32	12.18
T	8.88	11.85	14.68	13.51	12.23
R	7.88	12.28	13.07	13.89	11.78
RN	8.86	12.74	13.84	12.29	11.93
(± 0.397)					
A	8.34	12.57	13.89	13.21	12.00
S	8.75	12.44	13.75	13.30	12.06
(± 0.397)					
O T R RN					
A	12.12	12.18	11.93	11.78	
S	12.24	12.28	11.63	12.09	

64/C/18.3

	NO	N1	N2	N3	Mean
	SUGAR %				
Mean	20.2	20.4	19.9	18.9	19.8
O	19.8	20.6	20.1	18.8	19.9
T	20.4	20.6	19.7	18.7	19.8
R	20.4	20.2	20.0	19.4	20.0
RN	20.3	20.0	19.9	18.5	19.7
A	20.0	20.5	19.9	18.9	19.8
S	20.4	20.3	19.9	18.8	19.9
	O	T	R	RN	
A	19.8	19.8	20.1	19.6	
S	19.9	19.9	19.9	19.7	

64/C/18.4					
	NO	N1	N2	N3	Mean
TOTAL SUGAR					
(±1.24)					
Mean	34.6	51.0	55.1	50.0	47.7
(±2.49)					
O	34.0	54.2	55.1	50.1	48.3
T	36.3	48.8	57.9	50.5	48.4
R	32.1	49.6	52.2	53.8	46.9
RN	36.0	51.2	55.0	45.7	47.0
(±1.76)					
A	33.4	51.5	55.3	49.9	47.5
S	35.8	50.5	54.8	50.1	47.8
	O	T	R	RN	
(±1.76)					
A	48.0	48.0	47.8	46.2	
S	48.7	48.7	46.1	47.7	

64/C/18.5

	NO	N1	N2	N3	Mean
TOPS					
(± 0.231)					
Mean	3.73	6.00	7.66	8.15	6.38
(± 0.461)					
O T R RN	3.43	6.02	7.03	8.86	6.33
	3.68	5.79	7.91	8.76	6.53
	3.60	5.56	7.47	8.17	6.20
	4.21	6.63	8.21	6.82	6.47
(± 0.326)					
A S	3.74	6.24	7.72	8.09	6.45
	3.73	5.76	7.59	8.21	6.32
(± 0.163)					
	O	T	R	RN	
(± 0.326)					
A S	6.64	6.41	6.31	6.43	
	6.03	6.66	6.08	6.50	