

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1965

[Full Table of Content](#)



65/R/AX/C/10 N and N P K Residues - Barley

Rothamsted Research

Rothamsted Research (1966) *65/R/AX/C/10 N and N P K Residues - Barley* ; Yields Of The Field Experiments 1965, pp 147 - 149 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/C/10.1

N AND NPK RESIDUES

(AX)

An investigation of the relative importance of short term N and PK residues - Fosters West Side, the second year, barley 1965.

Design: 4 randomised blocks of 16 plots (with kale and wheat 1964 on sub-blocks of 8 plots) with interactions involving residual N confounded.

Area of each plot: 0.0193. Area harvested: 0.0128.

Treatments: All combinations of:-

In 1964

1. Spring wheat and kale.
2. N: None (R0), 0.5 (R1), 1.0 (R2), 1.5 (R3) cwt as 'Nitro-Chalk'.
3. PK: None (0), 2.0 cwt P205 and 2.0 cwt K20 as compound (0:20:20) (C).

To barley 1965

4. N: None (N0), 0.33 (N1), 0.66 (N2), 1.00 (N3) cwt N as 'Nitro-chalk'.

Basal applications: Manures: None. Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 34 gals).

Cultivations, etc.: Ploughed: Dec 9, 1964. 'Nitro-Chalk' applied, seed drilled at 156 lb: Mar 29, 1965. Sprayed: May 20. Combine harvested: Sept 1. Variety: Maris Badger.

Standard error per plot. Grain:
Sub plot: 1.67 or 4.9% (14 d.f.)

65/C/10.2

SUMMARY OF RESULTS

	NO	N1	N2	N3	Mean
AFTER WHEAT 1964					
Mean (± 0.59)*	25.7	36.9	37.5	37.1	34.3
		(± 1.18)*			(± 0.59)*
R0	24.5	35.1	37.5	38.8	34.0
R1	22.9	36.0	37.1	38.0	33.5
R2	27.5	38.5	38.2	36.8	35.3
R3	28.0	37.8	37.2	34.9	34.5
		(± 0.83)*			(± 0.42)*
O	26.5	37.6	37.4	37.2	34.7
C	25.0	36.1	37.5	37.1	33.9
AFTER KALE 1964					
Mean (± 0.59)*	21.5	34.5	38.7	38.8	33.3
		(± 1.18)*			(± 0.59)*
R0	19.8	33.7	38.4	40.2	33.0
R1	19.2	34.8	39.0	36.9	32.5
R2	22.7	33.1	39.4	39.4	33.6
R3	24.4	36.3	37.9	38.6	34.3
		(± 0.83)*			(± 0.42)*
O	22.8	34.0	38.4	39.4	33.6
C	20.2	34.9	38.9	38.2	33.1

* For use in comparisons only within the same previous crop

65/C/10.3

Mean over 'after Wheat' and 'after Kale' 1964

	NO	N1	N2	N3	Mean
Mean (± 0.42)	23.6	35.7	38.1	38.0	33.8
		(± 0.83)			(± 0.42)
R0	22.2	34.4	38.0	39.5	33.5
R1	21.0	35.4	38.0	37.4	33.0
R2	25.1	35.8	38.8	38.1	34.4
R3	26.2	37.0	37.6	36.8	34.4
		(± 0.59)			(± 0.29)
O	24.6	35.8	37.9	38.3	34.2
C	22.6	35.5	38.2	37.6	33.5

Mean D.M. %: 78.3