

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1965

[Full Table of Content](#)



65/R/IR/C/14 Irrigation - Potatoes and Spring Beans

Rothamsted Research

Rothamsted Research (1966) *65/R/IR/C/14 Irrigation - Potatoes and Spring Beans ; Yields Of The Field Experiments 1965*, pp 161 - 169 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/C/14.1

IRRIGATION

(IR)

The effect of irrigation on potatoes and spring beans - Great Field I and II 1965.

Design:

Potatoes: A single replicate of $4 \times 2 \times 3 \times 2 \times 2$ in two blocks. Each block is divided into 4 whole plots for the irrigation treatments, and at right angles into 6 strips for varieties and spacing. There is a further split for chitting and N. Spring beans: $4 \times 2 \times 2 \times 2 \times 2$ in two blocks. Each block is divided into 4 whole plots for irrigation, these being divided into 8 sub-plots.

Area of each sub-plot:-

Potatoes: 0.0204.	Area harvested: 0.0080.
Spring beans: 0.0321.	Area harvested: 0.0210.

Treatments: potatoes: All combinations:-

Whole plots. 1. Irrigation: None (O), early (A), late (B), full (C).

To strips of 1/6th plots.

2. Varieties: Majestic (M), King Edward (E).
3. Spacing: Seed 12 (S1), 15 (S2), 18 inches (S3) apart in row.

1/12th plots. 4. Chitting: None (O), seed chitted (Ch).

5. Nitrogen: 0.75 (N1), 1.5 (N2) cwt N as 'Nitro-Chalk'.

Spring beans: All combinations of:-

Main plots. 1. Irrigation: None (O), early (A), late (B), full (C).

Sub-plots. 2. Row spacing: 10.5 (C), 21 inches (W).

3. Seed rate: 200 (L), 300 lb (H).

4. Rates of PK: 400 (F1), 560 lb (F2) (0:20:20).

5. Methods of applying fertiliser: Broadcast (B), placed (P).

Basal applications:

Potatoes: 2.0 cwt P2O5, 4.0 cwt K2O as (0:14:28) rotary cultivated in. Weedkiller: Paraquat at 0.75 lb ion plus linuron at 2 lb in 40 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions.

Spring beans: Manures: None. Weedkiller: Simazine at 1 lb in 40 gals. Insecticide: Menazon at 4 oz in 40 gals.

65/C/14.2

Cultivations, etc.:

Potatoes: Ploughed: Dec 21 - Feb 17, 1964. Basal PK compound applied: Apr 8, 1965. 'Nitro-Chalk' applied: Apr 12. Rotary cultivated, potatoes planted: Apr 15. Sprayed, weed-killer: May 13, fungicide: June 30, July 27, Aug 10. Sprayed with undiluted BCV at 15 gals: Aug 27. Lifted: Oct 21. Previous crops: Barley 1963 and 1964.

Spring beans: Ploughed: Feb 17, 1965. Seed placement drilled for C treatment: Mar 16. Fertiliser broadcast for C treatment: Mar 17. Seed placement drilled and fertiliser broadcast for W treatment: Mar 25. Sprayed, weedkiller: Mar 29, insecticide: June 11. Combine harvested: Oct 6. Variety: Garton's Pedigree Tick. Previous crops: Barley 1963 and 1964.

RAINFALL AND IRRIGATION: INCHES

Week- ending	Rain	Potatoes A and C	Beans A and C
6 May	0.78		
13	0.10		
20	0.52		
27	0.48		
3 June	0.14		0.50
10	0.68		
17	0.94		
24	0.69		
1 July	0.07		
8	0.33	1.00	
15	1.33		
22	0.84		
29	1.58		
5 Aug	0.70		
12	0.03		
19	0.37		
26	0.76		
2 Sept	0.25		
Total	10.59	1.00	0.50

Standard errors per plot.

Potatoes, total tubers:	Pooled whole & strip:	1.017 or 6.4% (9 d.f.)
	1/6 plot:	0.639 or 4.0% (21 d.f.)
	1/12 plot:	1.386 or 8.7% (28 d.f.)
Spring beans, grain:	Whole plot:	0.70 or 2.5% (5 d.f.)
	Sub plot:	2.12 or 7.6% (23 d.f.)

65/c/14.3

SUMMARY OF RESULTS

POTATOES

TOTAL TUBERS

	O & B	A & C	Mean
Mean	16.39	15.55	15.97
	(±0.415)		
M	16.26	15.88	16.07
E	16.53	15.22	15.87
	(1) and (2)		(±0.415)
S1	16.68	16.42	16.55
S2	16.32	14.93	15.62
S3	16.18	15.31	15.74
	(3) and (4)		(±0.519)
O	16.10	14.49	15.29
CH	16.69	16.61	16.65
	(5) and (6)		(±0.200)
N1	16.60	15.50	16.05
N2	16.19	15.61	15.90
	(5) and (6)		(±0.200)

- (1) (±0.184) (3) (±0.226) For use in interaction comparisons only
 (2) (±0.435) (4) (±0.533) For use in all other comparisons
 (5) (±0.461) For use in horizontal and diagonal comparisons
 (6) (±0.283) For use in vertical and interaction comparisons

65/c/14.4

POTATOES

TOTAL TUBERS

	MS1	MS2	MS3	ES1	ES2	ES3
O & B	16.18	16.76	15.84	17.19	15.87	16.52
A & C	16.60	15.63	15.42	16.25	14.22	15.19
	MO	MCH	EO	ECH		
O & B	16.06	16.46	16.14	16.91		
A & C	14.64	17.12	14.34	16.10		
	MN1	MN2	EN1	EN2		
O & B	16.43	16.08	16.76	16.29		
A & C	15.69	16.07	15.30	15.14		
	S10	S1CH	S20	S2CH	S30	S3CH
O & B	16.53	16.84	16.05	16.59	15.72	16.64
A & C	15.50	17.35	14.22	15.64	13.76	16.86
	S1N1	S1N2	S2N1	S2N2	S3N1	S3N2
O & B	16.53	16.84	17.12	15.52	16.16	16.20
A & C	16.24	16.61	15.38	14.47	14.87	15.74
	ON1	ON2	CHN1	CHN2		
O & B	16.43	15.77	16.77	16.60		
A & C	14.51	14.47	16.48	16.74		

65/c/14.5

POTATOES

% WARE

	O & B	A & C	Mean
Mean	97.0	96.4	96.7
M	97.4	97.0	97.2
E	96.7	95.8	96.2
S1	96.8	96.1	96.5
S2	96.8	96.3	96.5
S3	97.5	96.8	97.1
O	96.7	95.6	96.2
CH	97.4	97.1	97.3
N1	97.1	96.3	96.7
N2	97.0	96.4	96.7

65/c/14.6

POTATOES

% WARE

	MS1	MS2	MS3	ES1	ES2	ES3
O & B	96.9	97.3	97.9	96.7	96.3	97.2
A & C	96.9	96.7	97.4	95.3	95.8	96.1
	MO	MCH	EO	ECH		
O & B	97.4	97.4	96.0	97.4		
A & C	96.9	97.1	94.4	97.1		
	MN1	MN2	EN1	EN2		
O & B	97.4	97.4	96.7	96.7		
A & C	97.0	97.0	95.6	95.9		
	S10	S1CH	S20	S2CH	S30	S3CH
O & B	96.7	96.9	96.2	97.4	97.2	97.8
A & C	95.3	97.0	95.7	96.8	95.9	97.6
	S1N1	S1N2	S2N1	S2N2	S3N1	S3N2
O & B	96.7	96.9	96.8	96.8	97.6	97.4
A & C	95.9	96.3	96.3	96.3	96.9	96.7
	ON1	ON2	CHN1	CHN2		
O & B	96.8	96.6	97.3	97.5		
A & C	95.5	95.7	97.1	97.1		

65/C/14.7

SPRING BEANS

GRAIN

	O & B	A & C	Mean
Mean	28.8	27.4	28.1
	(±0.35)		
C	29.8	28.2	29.0
W	27.8	26.6	27.2
	(1) and (2)		(±0.37)
L	29.4	28.0	28.7
H	28.2	26.8	27.5
	(1) and (2)		(±0.37)
F1	28.9	26.7	27.8
F2	28.6	28.0	28.4
	(1) and (2)		(±0.37)
B	29.8	28.2	28.9
P	27.8	26.6	27.2

(1) (±0.51) For use in horizontal and diagonal comparisons

(2) (±0.53) For use in vertical and diagonal comparisons

65/C/14.8

SPRING BEANS

GRAIN

	CL	WL	CH	WH
		(3) and (4)		
O & B	30.3	28.4	29.4	27.0
A & C	28.8	27.2	27.6	25.9
	CF1	WF1	CF2	WF2
		(3) and (4)		
O & B	29.6	28.1	30.0	27.4
A & C	26.8	26.6	29.6	26.6
	CB	WB	CP	WP
		(3) and (4)		
O & B	30.4	29.1	29.2	26.4
A & C	28.6	27.6	27.8	25.5
	IF1	HF1	IF2	HF2
		(3) and (4)		
O & B	29.8	28.0	28.8	28.4
A & C	27.2	26.2	28.8	27.4
	LB	HB	LP	HP
		(3) and (4)		
O & B	30.8	28.8	27.9	27.7
A & C	28.6	27.7	27.4	25.8

(3) (± 0.72) For use in vertical and diagonal comparisons(4) (± 0.75) For use in horizontal and interaction comparisons

65/c/14.9

SPRING BEANS

GRAIN

	F1B	F2B	F1P	F2P
	(3) and (4)			
D & B	30.0	29.5	27.8	27.8
A & C	27.8	28.5	25.6	27.6

(3) (± 0.72) For use in vertical and diagonal comparisons

(4) (± 0.75) For use in horizontal and interaction comparisons