

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1965

[Full Table of Content](#)



65/R/BQ/C/27 Previous Crops and N for Barley - Spring Wheat, Kale and Ryegrass

Rothamsted Research

Rothamsted Research (1966) *65/R/BQ/C/27 Previous Crops and N for Barley - Spring Wheat, Kale and Ryegrass* ; Yields Of The Field Experiments 1965, pp 217 - 219 - DOI:
<https://doi.org/10.23637/ERADOC-1-159>

65/C/27.1

PREVIOUS CROPS X N FOR BARLEY

(BQ)

The effect of previous cropping and nitrogen on the yield of barley -
Stackyard 1965, the 1st year.

Crops in 1965: Spring wheat, kale and ryegrass.

Design for 1966: 3 randomised blocks of 9 plots.

Area of each plot: 0.0321. Area harvested: Spring wheat - 0.0214,
kale - 0.0161, ryegrass - 0.0069.

Treatments applied in 1965: All combinations of:-

1. Cropping: Spring wheat (W), kale (K), Italian ryegrass (G).
2. Nitrogen: None (NO), 1.0 (N2), 2.0 cwt (N4) N as 'Nitro-Chalk'.

NOTE: Barley 1966 will test in addition:

3. Nitrogen: None (NO), 0.5 (N1), 1.0 cwt (N2) as 'Nitro-Chalk'.

Basal applications: 1.0 cwt P2O5, 2.0 cwt K2O as compound (0:14:28).

Weedkillers: To wheat: MCPA at 1.12 lb and dicamba at 0.08 lb in
40 gals, to ryegrass: 0.44 lb 2,4-D butoxyethylester in 40 gals.

Cultivations, etc.: Ploughed: Nov 24, 1964. Rotary cultivated twice:
Mar 15 and 31, 1965. Basal PK applied, N applied: Apr 2.

Spring wheat: Seed drilled at 188 lb: Apr 3, 1965. Sprayed:

May 22. Combine harvested: Sept 20. Variety: Opal.

Kale: Rotary cultivated: May 10, 1965. Seed drilled at 2 lb:

May 12. Harvested: Nov 1. Variety: Thousand Head Canson.

Ryegrass: Seed drilled at 40 lb: Apr 3, 1965. Sprayed: June 2.

Cut 3 times: July 7, Aug 9, Oct 7. Variety: Italian.

NOTE: Samples were taken from each crop at harvest for estimation
of N percentage. Soil samples were taken after harvest for
estimation of mineralisable N.

Standard errors per plot.

Spring wheat, grain:	2.70 or 7.4% (4 d.f.)
Kale, fresh weight:	2.231 or 11.8% (4 d.f.)
Ryegrass, dry matter, 1st cut:	4.83 or 16.9% (4 d.f.)
2nd cut:	2.52 or 14.3% (4 d.f.)
3rd cut:	3.41 or 35.7% (4 d.f.)
Total of 3 cuts:	7.46 or 13.4% (4 d.f.)

65/C/27.2

SUMMARY OF RESULTS

NO	N2	N4	Mean
----	----	----	------

SPRING WHEAT

GRAIN

	(± 1.56)		
28.4	40.4	41.4	36.7

STRAW

29.8	39.3	36.8	35.3
------	------	------	------

Mean D.M.%: Grain 75.9
Straw 78.1

KALE

FRESH WEIGHT

	(± 1.288)		
13.23	19.87	23.66	18.92

65/c/27.3

RYEGRASS
DRY MATTER

NO	N2	N4	Mean
1ST CUT			
14.8	34.5 (±2.79)	36.4	28.6
2ND CUT			
7.7	17.5 (±1.46)	27.7	17.6
3RD CUT			
5.0	9.2 (±1.97)	14.4	9.5
TOTAL OF 3 CUTS			
27.5	61.2 (±4.31)	78.5	55.8

Mean D.M. %: 1st cut 16.0
 2nd cut 19.9
 3rd cut 21.0
 Total of 3 cuts 19.0