

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/BQ/C/18 Previous Crops and N for Barley 1965-66 - Barley

Rothamsted Research

Rothamsted Research (1967) *66/R/BQ/C/18 Previous Crops and N for Barley 1965-66 - Barley ; Yields Of The Field Experiments 1966*, pp 185 - 187 - DOI:
<https://doi.org/10.23637/ERADOC-1-158>

66/c/18.3

GRAIN

	R0				R2				R4			
	NO	N1	N2	NO	NO	N1	N2	NO	N1	N2	NO	N2
	(1) and (2)											
W	10.5	22.7	33.3	16.1	25.7	37.2	21.8	35.1	38.3			
K	22.0	32.8	34.6	21.0	31.9	42.7	21.9	36.3	43.5			
G	9.1	22.9	31.9	8.7	28.8	34.1	17.7	26.0	39.1			

(1) (± 2.27) For use in comparisons involving different combinations of Crop and R

(2) (± 1.77) For use in comparisons involving the same combinations of Crop and R

Mean D.M. %: 85.6

Revised

1000 2000 3000 4000

(1) (11-13) 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000
(2) (15-21) 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000
1 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000
2 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000
3 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000

(3) 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000

10	20	30	40	50	60	70	80	90	100
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000
1000	2000	3000	4000	5000	6000	7000	8000	9000	10000

66/c/19.1

PREVIOUS CROPS & N FOR BARLEY

(BY)

The effect of previous cropping and nitrogen on the yield of
barley - Fosters Corner 1966, the 1st year.

Crops in 1966: Spring wheat, kale and ryegrass.

Design: 3 randomised blocks of 9 plots.

Area of each plot: 0.0321. Area harvested: Spring wheat - 0.0214,
kale - 0.0161, ryegrass - 0.0069.

Treatments applied in 1966: All combinations of:-

1. Cropping: Spring wheat (W), kale (K), Italian ryegrass (G).
2. Nitrogen: None (NO), 1.0 (N2), 2.0 cwt (N4) N as 'Nitro-Chalk'.

NOTE: Barley 1967 will test in addition:-

3. Nitrogen: None (NO), 0.5 (N1), 1.0 cwt (N2) N as 'Nitro-Chalk'.

Basal applications: 9 cwt compound (0:20:20). Weedkiller:
To wheat: Mecoprop/2,4-D (Methoxone Extra at 6 pints in
30 gals).

Cultivations, etc.: Ploughed: Nov 16, 1965. Basal PK applied:
Mar 9, 1966. 'Nitro-Chalk' applied: Mar 16.

Spring wheat: Seed drilled at 170 lb: Mar 16. Weedkiller
applied: May 16. Combine harvested: Sept 3. Variety:
Kloka.

Kale: Seed drilled at 1.5 lb: Apr 29, 1966. Harvested:
Nov 8. Variety: Thousand Head.

Ryegrass: Seed drilled at 40 lb: Mar 18, 1966. Cut 3
times: June 30, Aug 18, Oct 12. Variety: S22 Italian.

Previous crops: Kale and potatoes 1964, barley 1965.

NOTE: Samples were taken from each crop at harvest for estimation
of N percentage. Soil samples were taken on Mar 16 for
estimation of mineralisable N and pH.

Standard errors per plot.

Spring wheat, Dry matter: 1.69 or 3.9% (4 d.f.)

Kale, Fresh weight: 2.590 or 10.8% (4 d.f.)

Ryegrass, Dry matter, 1st cut: 2.99 or 11.1% (4 d.f.)

2nd cut: 1.93 or 7.7% (4 d.f.)

3rd cut: 0.58 or 6.2% (4 d.f.)

Total of 3 cuts: 4.35 or 7.1% (4 d.f.)